



UNITED NATIONS ENVIRONMENT PROGRAMME

# Marine pollution

UNEP Regional Seas Reports and Studies No. 25

Na.86-6475

Note: This document is the reproduction of document A/36/452 submitted to the thirty-sixth session of the United Nations General Assembly. For details see the preface.

For bibliographic purposes this document may be cited as:

UNEP: Marine pollution. UNEP Regional Seas Reports and Studies No. 25. UNEP, 1982.

#### PREFACE

8y resolution 34/103 of 18 December 1979, the United Nations General Assembly requested the Governing Council of UNEP to report to it at its thirty-sixth session, through the Economic and Social Council, on the problems relating to marine pollution—.

As a response to this request, the Executive Director of UNEP prepared this report which was considered, as document UNEP/GC.9/5/Add.3, by the minth session of the Governing Council (Nairobi, 13 - 26 May 1981).

The Governing Council authorized the Executive Director of UNEP to submit the report, on its behalf, to the General Assembly, through the Economic and Social Council (UNEP/GC.9/15, Decision 9/10A, paragraph 6).

The Secretary-General of the United Nations transmitted the report to the thirty-sixth session of the General Assembly as document A/36/452.

The definition of marine pollution long employed by the United Nations, as set forth in the General Principles for Assessment and Control of Marine Pollution used at the United Nations Conference on the Human Environment, is "the introduction by men, directly or indirectly, of substances or energy into the marine environment (including estuaries) resulting in such deletarious effects as harm to living resources, hazards to human health, hindrance to marine activities including fishing, impairment of quality for use of sea water, and

## CONTENTS

		Paragraph <del>o</del>
I	INTRODUCTION	1 - 6
II	SOURCES OF MARINE POLLUTION AND POSSIBLE TRENDS OF CHANGES IN THEIR OCCURRENCE	7 - 22
	Petroleum hydrocarbons	10 - 11
	Halogenated hydrocarbons	12 - 13
	Metals	14 = 15
	Radionuclides	16 - 19
	Persistent solids	20
	Other pollutants	21 - 22
111	EFFECTS OF LIVING RESOURCES, ECOSYSTEMS, HUMAN HEALTH AND AMENITIES	23 - 31
	Living resources	23 - 24
	Ecosystems	25 - 27
	Human health and amenity values	28 - 31
Į	ACTIONS TO ALLEVIATE MARINE POLLUTION PROBLEMS	32
٧	THE ROLE OF UNEP IN ALLEVIATING MARINE POLLUTION	33 - 35
۸ī	RECOMMENDATIONS	36

#### 1 INTRODUCTION

- 1. Problems of marine pollution have not altered greatly in the last decade, but man's perception of them has changed markedly. Improvements in marine environmental quality have been achieved in many coastal areas through pollution control, and severe degradation in some areas has been avoided through the application of stringent restrictions on waste discharges. Regional agreements aiming at the protection of enclosed and semi-enclosed seas, such as the Helsinki, Barcelona and Kuwait Conventions, have entered into force providing the legal basis for co-operation among States on this subject,
- 2. National priorities have shifted since the 1960s and the early 1970s. The acute energy shortages in many parts of the world have necessitated measures to provide new sources of energy, and these have sometimes introduced new pollution problems, and at other times required relaxation of existing strict regulatory controls.
- 3. Protection of coastal waters, in particular those in closed and semi-enclosed seas, against environmental damage deserves high priority. The impact on the marine environment of conventional energy sources, for example through the explaitation of different sections of the continental shelf for fossil fuels, the transport of liquid and gaseous hydrocarbons by tanker and pipeline, and the use of coal for thermal power, must be fully understood. The environmental effects of nuclear power reactors located on rivers discharging into coastal waters, constructed on the coast or mounted on offshore structures must be carefully evaluated before construction proceeds, and these impacts should be minimized by careful selection of sites and development of design to incorporate environmental requirements.
- 4. The main thrust in the 1980s must continue to be towards protection of coastal waters, estuaries and other inland waters, along with restoration of already degraded coastal marine environments. Most of the living resources of the sea are found in these vulnerable coastal zones. Coastal resources could be severely reduced by pollution long before adverse trends can be detected in the open oceans. Human populations are generally growing throughout the world, and pressures on the coastal zones are increasing as a larger proportion of the population moves to the coastal developments for industry and for human habitation can permanently destroy vital estuarine and inshore coastal habitats for marine organisms. The impact can be especially severe in sensitive coastal ecosystems, such as sea-grass beds, mangrove swamps and coral reefs. Often the effects are of a chronic nature, and their impact can only be assessed after long-term studies.
- 5. The choice of alternative energy sources may lead in the next decade or two to utilization of the sea as a source of energy. Tidal power is already being harnessed in significant quantities in at least two coastal areas of the world. Preliminary feasibility studies have demonstrated that energy can also be generated by wave action, coastal winds, salinity gradients and vertical temperature gradients. The utilization of marine biomass (large alose) for methans production

6. Finally, the effects of marine pollution on human health and the quality of life must be fully considered. The elimination of possible transmittal of pathogens through ses-water and seafood must receive high priority. The seathetic quality of coastal waters should not be ignored. Sewage treatment and improved mathods of waste disposal into the sea should be introduced whenever necessary to make the use of coastal regions safer for residents and visitors and more attractive for tourism.

#### 11 SOURCES OF MARINE POLLUTION AND POSSIBLE TRENDS OF CHANGES IN THEIR OCCURRENCE

- 7. The chief sources of marine pollution are land-based and may reach the sea through rivers, direct coastal out-falls, coastal urban and agricultural runoff, and precipitation of atmospheric emissions. Pollutants may also be introduced by shipping and by structures such as oil drilling rigs mounted on the continental shelf.
- 8. Because of the paucity of reliable baseline data on the levels of major marine pollutants in the world oceans, the trend in their occurrence is barely understood. However, intensive regional baseline studies carried out during the last decade in some areas (e.g. the North Atlantic, the Baltic, the Mediterranean) are shedding more light on the situation in the areas studied.
- 9. The major categories of pollutants which may have a long-term impact on the global oceans are: petroleum hydrocarbons, halogenated hydrocarbons, metals, radionuclides and persistent solids.

#### Petroleum hydrocarbone

- 10. Petroleum hydrocarbons reach the sea through various routes, with only a small part of the total coming from ships, even though catastrophic oil spills from large tankers are most dramatic. The total amount of oil introduced into the sea annually has been estimated at 6.113 million tonnes, of which 2.133 million tonnes come from various sources involved in transportation of oil by sea; only 0.2 million tonnes of that is attributable to tanker accidents. Rivers introduce about 1.6 million tonnes annually, while natural seeps are estimated to contribute 0.6 million tonnes of petroleum hydrocarbons annually, the same amount reaching the sea from the atmosphere. The balance of the annual input is contributed by various sources, including coastal refineries and coastal municipal wastes.
- 11. There has been an increase during the last decade in the amount of patroleum hydrocarbons transported on the world oceans. The number and volume of spills have also grown. Nevertheless, the amount of oil introduced into the oceans in 1980 has probably not increased greatly since the early 1970s.

#### Halogenated hydrocarbons

12. Halogenated hydrocarbons in the sea originate largely from terrestrial application of chlorinated hydrocarbon posticides in agriculture and forestry. These synthetic chemicals may reach the sea through agricultural runoff and rivers.

ang<u>raphy and the lighted of the second of t</u>

13. In the northern hemisphere, at least at middle latitudes, the use of auch halogenated hydrocarbons as DOTs and PCBs was curtailed around 1972 (although other peaticides, e.g. toxephene, have taken the place of OOTs) and they have thus generally shown decreased concentrations in estuaries and coastal waters along the North Atlantic and Pacific during the last decade. Their application in tropical and subtropical areas has, however, not substantially abated, and in the southern hemisphere it has increased, particularly in South America. Therefore, it can be expected that concentrations of these compounds may increase in southern hemisphere and low-latitude northern hemisphere sea-waters.

#### <u>Metals</u>

- 14. Pollution by metals arises from various land-based operations such as mining, milling and smelting activities, metal plating and assorted manufacturing processes. Some of the metals may enter the sea through the aquatic route, while a certain proportion reaches the oceans via the atmosphere and is washed out by rain. The amount of iron, manganese, copper, zinc, lead, tin and antimony entering the sea through river discharges is by an order of magnitude higher than the amount discharged through natural geological processes. Smelter emissions may transmit substantial quantities of metals into the atmosphere. The same holds true for coal-burning thermal plants and metallurgical industries. Nevertheless, seewater concentrations of metals in oceanic areas are still regarded essentially as at "background" levels. The problem of mercury is discussed below.
- 15. Lead is the only metal whose natural geochemical cycling has been clearly altered by man, leading to concentrations of lead in the surface waters of some parts of the oceans at above background levels. The use of tetraethyl-lead as an anti-knock agent in gasoline and the emissions from steel plants have increased the flux of lead to the atmosphere by an order of magnitude (0.44 million tonnes per year) over the natural flux. Other metals may exhibit comparatively high local concentrations in nearshors waters, where there may be industrial or urban sources of these metals, but their global budget has not been significantly altered.

## Radionuclides

- 16. Radionuclides have been reaching the sea through fallout from atmospheric testing of nuclear weapons. The annual deposition of fission products from nuclear tests reached a peak in 1963, but since the partial Treaty Banning Muclear Weapon Tests in the Atmosphere, in Outer Space and Under Water of 5 August 1963, this source of radionuclides has been steadily declining, although there have been a few small peaks superimposed on the decreasing trend as a result of atmospheric tests of nuclear weapons by countries which were not signatory to the Treaty.
- 17. Radioactive contributions to the marine environment from peaceful uses of radioactivity have been comparatively small, and reach the sea through rivers or direct coastal discharges. New nuclear power reactors, when operating normally, release little radioactivity into the atmosphere or into nearby waters. Consequently, emissions of radionuclides from nuclear power reactors have not substantially increased during the last decade. However, reactor accidents may lead to unpredictably high releases.
- 18. Low-level radioactive wastes sealed in containers were comped in the Atlantic

arta alemanti di deleggerata in progresse della dispersazzanta il progressioni del personali del presidente del

continued, and is now being conducted under the auspices of the European Nuclear Energy Agency. Ocean dumping of low-level and intermediate-level radioactive wastes from European sources will probably continue under the Convention for the Prevention of Marine Pollution by Dumping from Ships and Aircraft, 15 February 1972 (the Oslo Convention), and the Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter, 29 December 1972 (the London Dumping Convention).

19. As plans for new nuclear power reactors are implemented in many countries to supplement existing energy sources, it is expected that the entry of radioactive materials into the sea will increase. Reprocessing of nuclear fuel from power reactors may also add some radioactive materials to the sea.

## Persistent solids

20. Persistent solids, such as plastics, may have undesirable ecological effects in the sea and hinder maritime operations. In general, much of the persistent plastics and other litter entering the sea from land-based sources comes from garbage disposal and ordinary human refuse. Solid wastes, of which persistent plastics form only a part, are at present being introduced into the oceans in ever increasing amounts. Approximately  $6.4 \times 10^6$  tonnes of shipboard litter are discarded annually into the world's oceans. At present, only 0.7 per cent of this litter is plastic. However, plastics production is doubling every 12 years, so that the amount of plastic litter can be expected to increase substantially unless controls are applied.

#### Other pollutants

- 21. Nearshore marine pollution problems can also arise from a host of other agents that are not included in the above categories. While having perhaps little impact globally, they can have serious consequences in local areas. These are the materials that normally undergo degradation in the natural marine environment, but are often introduced in such quantities as to exceed the assimilative capacity of the local receiving waters. If there is a continuous discharge of such agents, chronic pollution can arise. These materials can be classified as:
  - (e) Dissolved organic substances;
  - (b) Particulate organic matter;
  - (c) Particulate inorganic matter:
  - (d) Soluble inorganic substances, including nutrient constituents;
  - (e) Micro-organisms;
  - (f) Thermal discharges.

Normally, these groups of substances are not considered to have a long-term, wide-ranging impact on the marine environment. They also may not have a serious effect on the nearshore environment, if they are effectively diluted with a large volume of receiving water and dispersed. Unfortunately, this is not always the case.

could arise from such mining operations through disturbance of the bottom and creation of turbidity by excavating nodules or dredging for metalliferous auds. Pollution could also result from the concentrating process. Whether the metals are extracted and concentrated at sea or on land, there would be tailings (pulverized rock) for disposal, which, if discharged into the sea, may create a turbidity problem. Moreover, metals may leach from the fine particles resulting from physical crushing and grinding of nodules followed by chemical treatment.

III EFFECTS OF LIVING RESOURCES, ECOSYSTEMS, HUMAN HEALTH AND AMENITIES

#### Living resources

- 23. Clear-cut effects of pollution on the living resources of the sea have not been unequivocably documented. There are no known cases of pelagic species of fish, for example, where populations have been shown to suffer a decline due strictly to pollution. The effects of fishing intensity, natural environmental factors, and success or failure of a particular year-class because of a combination of favourable or unfavourable conditions, have always dominated the variables that determine the size of a given pelagic fish population. It would be difficult to identify any impact of pollution superimposed on such far-ranging effects of other factors. The only instance of possible affect of a pollutant on the population of a marine organism was reported in the reproductive failure of Baltic Sea seals, which showed high levels of DDTs and PCBs in their tissues.
- 24. The impact of pollution on fisheries resources is felt most in the nearshore zone. This is where the various shellfish stocks, molluscan and crustacean, are located. Anadromous fish species use the estuaries and other parts of the coastal zone as nursery grounds before they proceed out to sea. Adults of both anadromous and catadromous species must swim through estuaries in order to reach their spawning grounds, and many coastal species feed in estuaries. Hence, pollutants in the coastal waters, or developments that disrupt or destroy inshore hebitats, can be disastrous for various coastal fish and invertebrate species. Long before impacts of pollution are recognized in offshore waters, the inshore fisheries resources can be severally reduced. This has happened to many stocks of cysters on both the Atlantic and the Pacific coasts of North America, while in Europe and on the east coast of North America, salmon and sea-run trout have vanished from streams where they were once abundant.

#### Ecosystems

- 25. The impact of pollution on marine ecosystems tends to be slow and insidious, and it is difficult to identify the most sensitive component of the ecosystem. The impact of pollution on Arctic and Antarctic ecosystems can be much more serious than in tropical and subtropical regions.
- 26. Disruption of food chains can occur in coastal waters as a result of pollution or coastal area development. Food chains in estuaries, in particular, are highly specialized and vulnerable to disturbance by man. They are extremely important from a practical point of view when they lead to economically valuable species such as

an karang berandan terma meng<u>alah pransada</u>kan keringgal dibanada dan d**a**ng lebengan dan dangkan <u>banda</u>ng

#### Human health and amenity values

- 28. Human health can be affected in a variety of ways by pollution in the merine environment. The most dramatic incidents are those where deaths or severe illnesses of humans are caused by consumption of contaminated seafood. Mercury, one of the more hazardous metals because of its neurological effect on human consumers when ingested in metallorganic form, has been widely studied in the last decade. Levels of mercury found in certain long-lived fish species in some regions (e.g. the Mediterranean) are up to four times higher (3 mg/kg) than the levels in the same species from open oceans. Nevertheless, it seems that these levels are a natural phenomenon not caused by man-made pollution.
- 29. Sewage effluents, or municipal wastewaters as they are sometimes known, contribute to perhaps the most universal form of marine pollution. Their impact is usually in inshore waters, which are used for growing and/or havesting of shellfish as well as for recreation such as bothing and bosting. The effect of sewage on such filter-feeders as oysters, clams and mussels is well known. These shellfish concentrate bacteris and viruses from sewage in the process of feeding. The consumption of raw or partially cooked shellfish which have been exposed to untrested sewage can thus lead to the transmittal of viral diseases such as hepatitis.
- 30. Sewage pollution in coastal waters affects water-related recreation and amenities. While cases of serious diseases, e.g. typhoid and cholera, due to sea bathing in sewage-polluted waters have never been conclusively proved, there is evidence of transmittal of certain mild afflictions, such as respiratory infections and gastro-enteritis, through this mode. The more undesirable aspect of sewage-pollution is its effect on amenity value of the water and beach. There may be visible turbidity, scum and other floatables on the water surface and beach, and an offensive odour, associated with the sewage discharge.
- 31. Oil spills are another form of pollution which has its undesirable visual and olfactory attributes, adversely affecting seashore residential and recreational areas, and sometimes causing severe set-backs in tourism.

## IV ACTIONS TO ALLEVIATE MARINE POLLUTION PROBLEMS

- 32. The approach to solving marine pollution problems requires a series of steps, which may be summarized as follows:
  - (a) Identification of pollution problems through assessment of their sources, amounts, levels, trends and effects.
  - (b) Development of environmental quality criteria, based on levels of pollutants in sea-water, marine blots and sediments that are not expected to damage marine organisms or ecosystems, or to be harmful to human health.
  - (c) Formulation and promulgation of national, regional and global legislation and regulations to control marine pollution, as an essential instrument of environmental menagement policies.

## Y THE ROLE OF UNEP IN ALLEVIATING MARINE POLLUTION

- 33. UNEP, as the "focal point for environmental action and co-ordination within the United Nations system" (General Assembly resolution 2997 (XXVII)) assists in, and in many cases co-ordinates, the development and implementation at the global, regional and national levels of the steps outlined in paragraph 32 above.
- 54. Specifically, UNEP concentrates on problems of the marine and coastal environment of ten regional seas, through programmes involving the co-operation of more than 188 coastal States and eighteen United Nations bodies and specialized agencies, as well as intergovernmental and non-governmental organizations. These afforts have led to the formal adoption of regional action plans for the protection and development of the Mediterranean (1975), the Red Sea and Gulf of Aden (1976), the Kuwait Action Plan Region (1978), the West African Region (1981), the Wider Caribbean (1981) and for the East Asian Seas (1981). (The action plan for the South-East Pacific may also be adopted in the second half of 1981). In some cases, within the framework of these Government-approved action plans, regional conventions for the protection and development of the marine environment and coastal areas, supplemented by protocols dealing more specifically with a particular source of pollution or environmental problem, have been adopted (Barcelona Convention, 1976; Kuwait Convention, 1978; Abidjan Convention 1981)—
- 35. In addition, UMEP actively supports, contributes to, or is associated with, the efforts of other United Nations and non-United Nations organizations related to the protection of the marine environment, for example through support to the secretariat of the Third United Nations Conference on the Lew of the Sea, GESAMP, GIPME, IGOSS, RIGS, the London Dumping Convention, the Oslo Convention, and the Conventions on the Protection of the Marine Environment of the Baltic Sea Area (Helsinki, 22 March 1974) and on the Prevention of Marine Pollution from Land-Based Sources (Paris, 4 June 1974).

#### VI RECOMMENDATIONS

## 36. It is recommended that:

- (a) the continued assessment of the sources, amounts and affects of marine pollutants should be actively pursued by UNEP as a component of Earthwatch, through support for national, regional and global afforts;
- (b) the work on the development and implementation of regional action plans for the protection of enclosed and semi-enclosed seas, as well as of cosstal waters of regions with defined common problems, should be intensified as the most rational approach to controlling marine pollution at its source through the full mobilization of national resources;

<sup>1/2</sup> For a more up-to-date information see UNEP: Achievements and planned

(c) the trends in marine pollution should be kept under constant review using existing mechanisms (e.g. GESAMP, SCOR, regional action plans) and periodic reports prepared on the state of the marine environment at both regional and global levels.

(d) full support should be given to the activities defined by the Law of the Sea Conference as related to the protection of the marine environment. <u> 1. april graduago i la politica substantamente i la ci</u>

#### PUBLICATIONS IN THE UNEP REGIONAL SEAS REPORTS AND STUDIES SERIES

- No. 1 CMEP: Achievements and planned development of UMEP's Regional Seas Programme and comparable programmes sponsored by other bodies. (1982)
- No. 2 UNIDO/UNEP: Survey of marine pollutants from industrial sources in the West and Central African region. (1982)
- No. 3 UNESCO/UNEP: River inputs to the West and Central African marine environment. (1982)
- No. 4 IMCO/UNEP: The status of oil pollution and oil pollution control in the West and Central African region. (1982)
- No. 5 [AEA/LMEP: Survey of tar, oil, chlorinated hydrocarbons and trace metal pollution in coastal waters of the Sultanate of Oman. (1982)
- No. 6 UM/UNESCO/UNEP: Marine and coastal area development in the East African region. [1982]
- No. 7 UNIDO/UNEP: Industrial sources of marine and coastal pollution in the East African region. (1982)
- No. 8 FAO/UNEP: Marine pollution in the East African region. (1982)
- No. 9 VHO/UNEP: Public health problems in the coastal zone of the East African region. (1982)
- No. 10 IMO/UNEP: Oil pollution control in the East African region. (1982)
- No. 11 IUCN/UNEP:Conservation of coastal and marine ecosystems and living resources of the East African region. (1982)
- No. 12 UNEP: Environmenta) problems of the East African region. (1982)
- No. 13 UNEP: Pollution and the marine environment in the Indian Ocean. (1982)
- No. 14 UNEP/CEPAL: Development and environment in the Wider Caribbean region: A Synthesis. (1982)
- No. 15 UNEP: Guidelines and principles for the preparation and implementation of comprehensive action plans for the protection and development of marine and coastal areas of regional seas. (1982)
- No. 16 GESAMP: The health of the oceans. (1982)
- No. 17 UNEP: Regional Seas Programme: Legislative authority. (1985)
- No. 18 UNEP: Regional Seas Programme: Horkplan. (1982)
- No. 19 Rev. 2. UMEP: UMEP Oceans Programme: Compendium of projects. (1985)
- No. 20 CPPS/INEP: Action Plan for the protection of the marine environment and coastal areas of the South-East Pacific. (1983)
- No. 21 CPPS/UNEP:Sources, levels and effects of marine pollution in the South-East Pacific.

illi orana kan ili orang mengarang kanang kanang kanang kanang pang kang di pang pang pang pang pang pang pang

- No. 23 FAO/UNESCO/IOC/MHO/MMO/IAEA/UNEP: Co-ordinated Mediterranean Pollution Monitoring and Research Programme (NED POL) Phase I: Programme Description. (1983)
- No. 24 UMEP: Action Plan for the protection and development of the marine and coastal areas of the East Asian region. (1983)
- Mo. 25 UNEP: Marine pollution. (1983)
- No. 26 UNEP: Action Plan for the Caribbean environment programme. (1983)
- No. 27 UNEP: Action Plan for the protection and development of the marine environment and coastal areas of the West and Central African region. (1983)
- No. 28 UMEP: Long-term programme for pollution monitoring and research in the Mediterranean (MED POL) Phase II. (1983)
- No. 29 SPC/SPEC/ESCAP/UNEP: Action Plan for managing the natural resources and environment of the South Pacific region. (1983)
- No. 30 UNDIESA/UNEP: Ocean energy potential of the West and Gentral African region. (1983)
- No. 31 A. L. DAHL and I. L. BAUMGART: The state of the environment in the South Pacific. (1983)
- No. 32 UMEP/ECE/UNIDO/FAO/UMESCO/WHO/IAEA: Pollutants from land-based sources in the Mediterranean. (1984)
- Mp. 33 UNDIESA/UNEP: Onshore impact of offshore oil and natural gas development in the West and Central African region. (1984)
- No. 34 UNEP: Action Plan for the protection of the Mediterranean. (1984)
- No. 35 UNEP: Action Plan for the protection of the marine environment and the coastal areas of Bahrain, Iran, Iraq, Kuwait, Cman, Qatar, Saudi Arabia and the United Arab Emirates. (1983)
- No. 36 UMEP/ECLAC: The state of marine pollution in the Wider Caribbean region. (1984)
- No. 37 UNDIESA/UNEP: Environmental management problems in resource utilization and survey of resources in the West and Central African region. (1984)
- No. 38 FAO/LINEP: Legal aspects of protecting and managing the marine and coastal environment of the East African region. (1983)
- No. 39 IUCN/UNEP: Marine and coastal conservation in the East African region. (1984)
- No. 40 SPC/SPEC/ESCAP/UNEP: Radioactivity in the South Pacific. (1984)
- No. 41 UNEP: Socio-economic activities that may have an impact on the marine and coastal environment of the East African region. (1984)
- No. 42 GESAMP: Principles for developing coastal water quality criteria. (1984)
- No. 43 CPPS/UNEP: Contingency plan to combat oil pollution in the South-Fast Parifir in cases

nakan kangan penggangan di perdagan kangan penggan berangan berangan penggan beranggan penggan penggan berang Penggan pengga

- No. 45 GESAMP: Thermal discharges in the marine environment. (1984)
- No. 46 UNEP: The marine and coastal environment of the West and Central African region and its state of pollution. (1984)
- No. 47 UNEP: Prospects for global ocean pollution munitoring. (1984)
- No. 48 SPC/SPEC/ESCAP/UNEP: Hazardous waste storage and disposal in the South Pacific. (1984)

  No. 48/ Appendices SPC/SPEC/ESCAP/UNEP: Hazardous waste storage and disposal in the South Pacific. (1984)
- No. 49 FAC/UNEP: Legal aspects of protecting and managing the marine and coastal environment of the East African region: National Reports. (1984)
- No. 50 IUCN/UNEP: Marine and coastal conservation in the East African region: Mational Reports. (1984)
- No. 51 UMEP: Socio-economic activities that may have an impact on the marine and coastal environment of the East African region: Mational Reports. (1984)
- No. 52 UMEP: Arab co-operation for the protection and development of the marine environment and coastal areas resources of the Mediterranean. (1984)
- No. 53 UMEP: UNEP Regional Seas Programme: the Eastern African Experience. (1984)
- No. 54 UMIDO/UNEP: Contingency planning for emergencies associated with industrial installations in the West and Central African region. (1985)
- No. 55 FAO/UNEP: Marine mammals; global plan of action. (1985)

  No. 55/ Annex FAO/IUCN/INC/UNEP: Marine mammals: global plan of action. (1985)
- No. 56 GESAMP: Cadmium, lead and tim in the marine environment. (1985)

- No. 57 IMO/UMEP: Oil spills and shoreline clean-up on the coasts of the Eastern African region. (1985)
- No. 58 UNEP: Co-operative programmes sponsored by UNEP for the protection of the marine and coastal environment in the wider Indian Ocean region. (1985)
- No. 59 UNEP: Environmental problems of the marine and coastal area of India: Mational Report. (1985)
- No. 60 IUCN/UNEP: Management and conservation of renewable marine resources in the Indian Ocean region: Overview. (1985)
- No. 61 UMEP: Action Plan for the protection, management and development of the marine and coastal environment of the Eastern African region. (1985)
- No. 62 IUCN/UNEP: Management and conservation of renewable marine resources in the South Asian Seas region. (1985)
- No. 63 IUCN/UNEP: Management and conservation of renewable marine resources in the Kuwait

- No. 64 IUCH/UNEP: Management and conservation of renewable marine resources in the Red Sea and Gulf of Aden region. (1985)
- No. 65 IUCH/UMEP: Management and conservation of renewable marine resources in the East Asian Seas region. (1985)
- No. 66 IUCN/UMEP: Management and conservation of renewable marine resources in the Eastern African region. (1985)
- No. 67 UN/UNEP: Coastal erosion in Hest and Central Africa. (1985)
- No. 68 GESAMP: Atmospheric transport of contaminants into the Mediterranean region. (1985)
- No. 69 UNEP: Environment and resources in the Pacific, (1985)

en de la composition La composition de la

- No. 70 BMESCO/ROPME/UPM/UNEP: Proceedings of the Symposium/Workshop on oceanographic modelling of the Kuwait Action Plan (KAP) region. (1985)
- No. 71 IUCN/ROPME/UMEP: An ecological study of the rocky shares on the southern coast of Oman.
  (1985)
- No. 72 IUCH/ROPME/UNEP: An ecological study of sites on the coast of Bahrain. (1985)
- No. 73 SPC/SPEC/ESCAP/UMEP: Ecological interactions between tropical coastal ecosystems. (1985)
- No. 74 UMEP: Environmental problems of the marine and coastal area of Sri Lanka: National Report (1986)
- No. 75 UMEP: Environmental problems of the marine and coastal area of Bangladesh: National Report (1986)
- No. 76 UMEP: Environmental problems of the marine and coastal area of Maldives: National Report (1986)
- No. 77 UMEP: Environmental problems of the marine and coastal area of Pakistan: National Report (1986)
- No. 78 GESAMP: Organosilicons in the marine environment. (1986)
- No. 79 H.I. SHUVAL: Thalassogenic diseases. (1986)
- No. 80 GESAMP: Environmental capacity: an approach to marine pollution prevention. (1986)
- No. 81 UMEP: Action Plan for the conservation of the marine environment and coastal areas of the Red Sea and Gulf of Aden. (1986)
- No. 82 UNEP: Environmental problems of the South Asian Seas region: An overview. (1986)

## Tasued and printed by:



Programme Activity Centre for Oceans and Coastal Areas United Nations Environment Programme

Additional copies of this and other publications issued by the Programme Activity Centre for Oceans and Coastal Areas can be obtained from:

Programme Activity Centre for Oceans and Coastel Areas United Nations Environment Programme