



MEDITERRANEAN ACTION PLAN  
MED POL

---

UNITED NATIONS ENVIRONMENT PROGRAMME



FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS

**EVALUATION OF THE TRAINING WORKSHOPS ON THE STATISTICAL TREATMENT  
AND INTERPRETATION OF MARINE COMMUNITY DATA**

**MAP Technical Reports Series No. 68**

**In cooperation with**



**IOC**

---

**UNEP**  
Athens, 1992

Note: The designations employed and the presentation of the material in this document do not imply the expression of any opinion whatsoever on the part of UNEP, FAO or IOC concerning the legal status of any State, Territory, city or area, or of its authorities, or concerning the delimitation of their frontiers or boundaries.

Note: Les appellations employées dans ce document et la présentation des données qui y figurent n'impliquent de la part du PNUE, de la FAO ou de la COI aucune prise de position quant au statut juridique des Etats, territoires, villes ou zones, ou de leurs autorités, ni quant au tracé de leurs frontières ou limites.

For bibliographic purposes this volume may be cited as:

UNEP/FAO/IOC: Evaluation of the Training Workshops on the Statistical Treatment and Interpretation of Marine Community Data. MAP Technical Reports Series No. 68. UNEP, Athens, 1992.

Pour des fins bibliographiques, citer le présent volume comme suit:

PNUE/FAO/COI: Evaluation des ateliers de formation sur le traitement statistique et l'interprétation des données relatives aux communautés marines. MAP Technical Reports Series No. 68. UNEP, Athens, 1992.

This volume is the sixty-eighth issue of the Mediterranean Action Plan Technical Reports Series.

This series contains selected reports resulting from the various activities performed within the framework of the components of the Mediterranean Action Plan: Pollution Monitoring and Research Programme (MED POL), Blue Plan, Priority Actions Programme, Specially Protected Areas and Regional Marine Pollution Emergency Response Centre for the Mediterranean.

Ce volume constitue le soixante-huitième numéro de la série des Rapports techniques du Plan d'action pour la Méditerranée.

Cette série comprend certains rapports élaborés au cours de diverses activités menées dans le cadre des composantes du Plan d'action pour la Méditerranée: Programme de surveillance continue et de recherche en matière de pollution (MED POL), Plan Bleu, Programme d'actions prioritaires, Aires spécialement protégées et Centre régional méditerranéen pour l'intervention d'urgence contre la pollution marine accidentelle.



## PREFACE

The United Nations Environment Programme (UNEP) convened an Intergovernmental Meeting on the Protection of the Mediterranean (Barcelona), 28 January - 4 February 1975), which was attended by representatives of 16 States bordering on the Mediterranean Sea. The meeting discussed the various measures necessary for the prevention and control of pollution of the Mediterranean Sea, and concluded by adopting an Action Plan consisting of three substantive components:

- Integrated planning of the development and management of the resources of the Mediterranean Basin (management component);
- Co-ordinated programme for research, monitoring and exchange of information and assessment of the state of pollution and of protection measures (assessment component);
- Framework convention and related protocols with their technical annexes for the protection of the Mediterranean environment (legal component).

All components of the Action Plan are interdependent and provide a framework for comprehensive action to promote both the protection and the continued development of the Mediterranean ecoregion. No component is an end in itself. The Action Plan is intended to assist the Mediterranean Governments in formulating their national policies related to the continuous development and protection of the Mediterranean area and to improve their ability to identify various options for alternative patterns of development and to make choices and appropriate allocations of resources.

The Co-ordinated Mediterranean Research and Monitoring Programme (MED POL) was approved as the assessment (scientific/technical) component of the Action Plan.

The general objectives of its pilot phase (MED POL - Phase I), which evolved through a series of expert and intergovernmental meetings, were:

- to formulate and carry out a co-ordinated pollution monitoring and research programme taking into account the goals of the Mediterranean Action Plan and the capabilities of the Mediterranean research centres to participate in it;
- to assist national research centres in developing their capabilities to participate in the programme;
- to analyse the sources, amounts, levels, pathways, trends and effects of pollutants relevant to the Mediterranean Sea;
- to provide the scientific/technical information needed by the Governments of the Mediterranean States and the EEC for the negotiation and implementation of the Convention for the Protection of the Mediterranean Sea against Pollution and its related protocols;
- to build up consistent time-series of data on the sources, pathways, levels and effects of pollutants in the Mediterranean Sea and thus to contribute to the scientific knowledge of the Mediterranean Sea.

MED POL-Phase I initially consisted of seven pilot projects and baseline studies on the monitoring of oil, petroleum hydrocarbons and microbial pollution in sea water, heavy metals and chlorinated hydrocarbons in marine organisms as well as research on the effects of pollutants on organisms, populations, communities and ecosystems. In addition, four related projects were also included to broaden the scope of the programme or to provide ancillary support.

Based on the recommendations made at various expert and intergovernmental meetings, a draft Long-term (1981-1990) Programme for pollution monitoring and Research in the Mediterranean (MED POL-Phase II) was formulated by the Secretariat of the Barcelona Convention (UNEP), in co-operation with the United Nations Agencies which were responsible for the technical implementation of MED POL-Phase I, and it was formally approved by the Second Meeting of the Contracting Parties of the Mediterranean Sea against pollution and its related protocols and Intergovernmental Review Meeting of Mediterranean Coastal States of the Action Plan held in Cannes, 2-7 March 1981.

The general long-term objectives of MED POL-Phase II were to further the goals of the Barcelona Convention by assisting the Parties to prevent, abate and combat pollution of the Mediterranean Sea area and to protect and enhance the marine environment of the area. The specific objectives were designed to provide, on a continuous basis, the Parties to the Barcelona Convention and its related protocols with:

- information required for the implementation of the Convention and the protocols;
- indicators and evaluation of the effectiveness of the pollution prevention measures taken under the Convention and the protocols;
- scientific information which may lead to eventual revisions and amendments of the relevant provisions of the Convention and the protocols and for the formulation of additional protocols;
- information which could be used in formulating environmentally sound national, bilateral and multilateral management decisions essential for the continuous socio-economic development of the Mediterranean region on a sustainable basis;
- periodic assessment of the state of pollution of the Mediterranean Sea.

The monitoring of, and research on, pollutants affecting the Mediterranean marine environment reflects primarily the immediate and long-term requirements of the Barcelona Convention and its protocols, but also takes into account factors needed for the understanding of the relationship between the socio-economic development of the region and the pollution of the Mediterranean Sea.

Individual and collective training is provided for scientists and technicians in techniques (methods) required for their effective participation in monitoring and research envisaged in the framework of MED POL - PHASE II. This assistance is in the form of fellowships, experts,

workshops, seminars, grants for attendance to meetings, etc., and covers training in analytical and sampling techniques, data processing, interpretation of results and various research topics.

As in MED POL-Phase I, the overall co-ordination and guidance for MED POL-Phase II is provided by UNEP as the secretariat of the Mediterranean Action Plan (MAP). Co-operating specialized United Nations Agencies (FAO, UNESCO, WHO, WMO, IAEA, IOC) are responsible for the technical implementation and day-to-day co-ordination of the work of national centres participating in monitoring and research.

The sixty-eighth volume of the MAP Technical Reports Series contains an overall report on the Training Workshops on the Statistical Treatment and Interpretation of Marine Community Data, which took place in the framework of the MED POL programme. The volume also includes the lecture notes in Annex.





TABLE OF CONTENTS

	<u>Page No.</u>
1. Introduction	1
2. Programme of the Workshops	3
3. Evaluation of the Workshops	7
4. Conclusions	8
ANNEX I : List of Participants	11
ANNEX II : Model Programme of the Workshops	17
ANNEX III: Lecture Notes	21

## 1. INTRODUCTION

The Long-term Programme for Pollution Monitoring and Research in the Mediterranean Sea (MED POL - Phase II), which constitutes the scientific/ technical component of the Mediterranean Action Plan, is basically divided into two groups of activities, namely monitoring and research. The research component until 1989 covered twelve topics one of which (research activity I) was concerned with the study of ecosystem modifications in areas influenced by pollutants or where large-scale coastal or inland engineering activities take place. Presently, the research component includes five research areas one of which focuses on the effects of marine pollution on marine life.

A number of research projects has been or is being implemented in the framework of the research component. Final reports on projects are published in the MAP Technical Reports Series.

The FAO/UNEP Meeting on the Effects of Pollution on Marine Ecosystems (Blanes, Spain, 7-11 October 1985) reviewed the work carried out within activity I. Regarding methods of data analysis, the meeting emphasized that no single method of analysis was adequate, and that ecological data could only be fully interpreted with the aid of a range of data analysis techniques. The organization of formal training courses in methods of data analysis was recommended.

Furthermore, the meeting recommended that in view of the wide range of criteria of pollutant effects, and the wide range of methods of study which are in use in different regions, a more rapid progress should be made in their evaluation and the identification of the most useful approaches.

The IOC Programme on Global Investigation of Pollution in the Marine Environment (GIPME) is developing through one of its groups, namely, the IOC/IMO/UNEP Group of Experts on the Effects of Pollutants (GEEP), techniques for quantification of biological effects of contamination including appropriate data analysis and sampling techniques.

As one step in this process, IOC/GEEP organised a series of workshops on the Biological Effects of Pollutants (Oslo, Norway, 11-29 August 1986; Bermuda, 10 September - 2 October 1988; Bremerhaven, Germany, 12-30 March 1990) at which participants were engaged in measuring biological responses in material gathered along a gradient of contamination and in material experimentally exposed to a mixture of contaminants in a mesocosm facility, and then in comparing their findings with chemical determinations carried out simultaneously. In this way, the performance, sensitivity and relative strengths and weaknesses of the biological procedures could be assessed in relation to common environmental gradients. The techniques tested included procedures to measure features of benthic community structure. Both field and experimental studies were subject to rigorous statistical design and analysis, including coding and sampling so that biological analyses were all performed "blind".

In 1987 the Inter-Agency Advisory Committee for the MED POL decided to recommend, as a follow-up activity to the Blanes meeting, the organisation of a training course to help Mediterranean scientists analyse statistically and interpret their data relevant to marine community structures (mainly benthos and plankton).

## Organisational details

In 1988 the first regional FAO/IOC/UNEP Training Workshop on the Statistical Treatment and Interpretation of Marine Community Data took place at the Marine Biological Station in Piran, Yugoslavia (now Slovenia), from 14-24 June. It drew from the experience of the "benthic community studies" component of the IOC workshop in Oslo and the associated development of statistical analysis programmes (software) tested at that Workshop. The training workshop was attended by 26 participants from ten Mediterranean countries (see Annex I for a full list of scientists who benefitted from this series of workshops).

As a large number of applications for participation in the workshop could not be satisfied, it was decided by the relevant UN Agencies to hold two workshops at a national level in the countries from which the majority of applications came from. As a consequence, a training workshop took place in 1989 in Athens, Greece from 18 to 29 September and in 1990 in Split, Yugoslavia (now Croatia) from 26 June to 6 July. The workshop in Athens was locally organised by the National Centre for Marine Research in cooperation with the University of Athens. It took place at the Biology Department of the University of Athens and was attended by 26 participants from all over Greece. The workshop in Split, Yugoslavia (now Croatia) took place at the Institute of Oceanography and Fisheries and was attended by 17 participants from Yugoslavia and 3 from Israel.

The fourth workshop in the series but second at a regional level, took place in Alexandria (Egypt) from 9 to 19 December 1991. It was hosted by the Arab Maritime Transport Academy and was attended by 21 participants from 7 Mediterranean countries. Finally, the training workshop was repeated for a 5th and final time in Israel at a national level. However, this time the workshop was condensed to five and a half days covering only the multivariate aspects. The workshop took place at the University of Bar-Ilan from 14 to 19 June 1992 and was attended by 15 participants from Israel.

Professor J.S. Gray (University of Oslo, Norway) and Dr. K.R. Clarke (Plymouth Marine Laboratory, U.K.) both members of IOC/GEEP were responsible for the scientific programme and the presentation of the lectures. Dr. R.M. Warwick (Plymouth Marine Laboratory, U.K.) joined the team of lecturers for the Athens, Split and Alexandria workshops. They were assisted by Mr. M.R. Carr (Plymouth Marine Laboratory, U.K.) who was responsible for supervising the preparation of the multivariate statistical software; he acted as demonstrator for the practical sessions in all the workshops. Dr. E. Papathanassiou (National Centre for Marine Research, Athens) acted as demonstrator in all workshops except the one in Piran. Mr. B. Reppe (University of Oslo, Norway) and Mr. R.G. Carter (NERC Computer Services, U.K.) acted as demonstrators at the Piran and Athens workshops respectively. Mr. G.P. Gabrielides, FAO Senior Fishery Officer (Marine Pollution) at the Coordinating Unit for the Mediterranean Action Plan was in charge of the overall organization. The information of this section is summarised in Table 1.

Table 1

Summarised information on the workshops

	Date	Place	Number of participants	Lecturers/ Demonstrators
1	14-24 June 1988	Piran, Slovenia	26	Gray, Clarke, Carr, Reppe
2	18-29 September 1989	Athens, Greece	26	Gray, Clarke, Warwick, Carr, Carter, Papathanassiou
3	26 June - 6 July 1990	Split, Croatia	20	Gray, Clarke, Warwick, Carr, Papathanassiou
4	9-19 December 1991	Alexandria, Egypt	21	Gray, Clarke, Warwick, Carr, Papathanassiou
5	14-19 June 1992	Tel-Aviv, Israel	15	Carr, Papathanassiou

## 2. PROGRAMME OF THE WORKSHOPS

The purpose of the 10-day training workshops was to expound, by means of lectures and a thorough practical programme, some of the wide range of statistical tools that have become available in the last few years for analysing community data (in the form of species abundance and biomass), arising in studies of biological effects of pollutants. The methods covered ranged from "classical" univariate statistics applied to population abundances, diversity indices etc., to multivariate clustering and ordination analyses, and other graphical techniques, applied to large arrays of samples/species data.

The emphasis was on the practical application of statistical analyses to a range of data sets from the literature, principally involving benthic communities examined along gradients in space or time, and to data sets on species abundances that the participants were encouraged to bring with them to the workshop.

Practical computations were performed on IBM/PC compatibles, using both the commercial statistics package STATGRAPHICS (covering principally univariate statistics) and the package PRIMER (Plymouth Routines In Multivariate Ecological Research), a suite of PC programmes written at the Plymouth Marine Laboratory, U.K., covering mainly multivariate statistics specific to community analyses and not available in commercial packages. The lectures and practical sessions drew from the experience of the benthic community studies components of the IOC/GEEP workshops on Biological Effects of Pollutants mentioned above.

Two categories of software were prepared: general data-handling programmes for manipulating species/samples arrays (or any other arrays), and specific statistical software (mainly multivariate) for analysing community data on abundance or biomass of a set of species over a series of samples.

The programmes have been written specifically for IBM-compatible PCs. The work was carried out by Mr. M.R. Carr, Mr. R.G. Carter and several students (Jane Addy, Ralph Bruno, Robert Pritchard and Martin Budge) under the supervision of Mr. Carr and Dr. Clarke of the Plymouth Marine Laboratory, United Kingdom.

The minimum hardware requirements for running the programmes are a machine compatible with an IBM PC XT (or preferably, PC AT or PS/2) with 640K memory, a hard disk, a maths co-processor, an EGA graphics board (if it is required to view plots on the screen), a dot-matrix printer (for hard copy results listings and for low-resolution plots) and an IBM or other flat-bed plotter compatible with a Hewlett Packard 7475 or 7470 (if publication-quality hard-copy plots are required).

The programmes prepared are the following:

#### Data manipulation programmes

- AGGREG - Uses an aggregation file giving details of the genus/family etc. to which a species belongs. The input data file is a species abundance or biomass file in standard format. The rows (species) of this file are pooled to produce an output file in standard format with the same number of columns (samples) but a reduced number of rows. These rows will now represent genus/family etc.
- COLTOT - Adds or averages selected columns. The input file is in standard format, and the output is a standard format file containing the same number of rows.
- CONFMT - Converts between different file formats. It can convert between STANDARD, DIF, CSV and TSV.
- FILEFORM - Used to produce 'readable' data. Many output data files contain data in scientific format, for example 1.00000E+02 (equivalent to 100). Input and output are standard format files.
- FILEINFO - File information programme. Used to check for unexpected characters (keying errors) in the data file; provides information on the size of the file (number of rows and columns). Input is a standard format file.
- MERGE - Merges two or more files either side-by-side or end-to-end. All files are in standard format.
- QCKSTD - Converts from quick format to standard format. Various error checks are provided.

- REDUCE - Selects a subset of columns and rows. The input file is in standard format, and the output file is a subset of the input file in standard format.
- SWAP - Transposes a data file, i.e. rows become columns and columns become rows. Input and output are standard format data files.

#### Statistical analysis programmes

- ANOSIM - Analysis of Similarities. This is used to test for differences between groups of samples. Input is a similarity or dissimilarity matrix.
- CASWELL - Caswell's neutral model. Input is a standard format data file.
- CLUSTER - Performs hierarchical agglomerative clustering. The input data file can be a species/samples file, or a similarity/ dissimilarity matrix. A wide variety of options are available covering data selection, standardisation, transformations, different similarity measures and different link options. The output takes the forms of a results file, a similarity matrix, a plot file for input to DENPLOT, and an optional command file for input to MDS.
- CONPLOT - Plots a sample (or species) configuration using the configuration file obtained from PCA or MDS. Environmental data can also be superimposed (as a symbol of varying size) on the sample positions in the ordination.
- DENPLOT - Plots a dendrogram using the dendrogram file output by the CLUSTER programme.
- DIVERSE - Calculates a suite of diversity indices for a species/ samples data file. The input file and the output file are in standard format. In the output file, rows are samples, and columns are the different diversity indices.
- DOMPLOT - Comparisons of dominance curves. Graphical output in the form of dominance curves is produced. Input is a species-samples data file.
- GEOPLOT - Plots number of species against individuals per species in geometric size classes, (i.e. 1,2-3,4-7 etc. individuals per species as size classes 1,2,3 etc.). Input is a species-samples data file.
- MDS - Nonmetric multidimensional scaling programme. Input files are the similarity matrix and MDS command file from the CLUSTER programme. The programme outputs a results file and a configuration file.

- PCA - Principal Components Analysis. Input is a data file in standard format with columns as the variables, and row as the replicates (samples). The programme outputs a results file and a configuration file.
- SIMPER - Examines the contribution of individual species to the Bray-Curtis similarity measure. Input is a standard format data file.

The workshops included a set of introductory lectures and practical sessions (refresher course) on basic (univariate) statistics, so that all participants would be at comparable levels when undertaking the second (multivariate) part of the course. The emphasis given here was not to statistical theory but rather to the practicalities that marine ecologists face. Questions treated covered such problems as: how many samples? or how large a sample should one take?, what are the problems of sub-sampling?, what are the consequences of organisms being clustered rather than randomly distributed in space? etc. To answer these questions requires an understanding of some basic statistical terms such as variance, standard deviation and standard error, confidence limits, transformations etc. The introductory course covered these topics and then demonstrated how to compare univariate samples using Student's 't' test and the analysis of variance, the relationships between samples using regression and correlation analyses and finally non-parametric statistics.

The lecture material for the second part of the course was presented in the order of the main steps in a field study of changes in community structure (through time or in space), namely:

- a) the use of multivariate methods (clustering and ordination) to represent graphically the similarities between species abundances (or biomass) observed in a set of samples;
- b) the demonstration of statistically significant differences in species composition between several sites (or the same site at several times) - this is a necessary pre-requisite to further analyses attempting to explain those differences;
- c) the construction of univariate indices (e.g. diversity, indicator species) and distributional plots (e.g. individuals amongst species curves, abundance-biomass comparisons) which indicate levels of disturbance or "stress" at sites;
- d) the relation of both univariate and multivariate faunal descriptions to gradients of chemical contamination and background environmental variables.

The practical sessions allowed the participants to apply the methods described in the accompanying lectures, on published data sets chosen to illustrate changes in benthic community composition at macrofaunal and meiofaunal levels, resulting from contaminant impact by sewage sludge dumping, pulp mill effluent, oil spills etc.

Throughout the second half of the workshop, periods of time were scheduled for participants to enter and analyse their own data sets, using the new computing tools provided by the course. In the last two or so days, the emphasis was on completing at least a first attempt at analysis and interpretation of all data sets brought to the course, and these were successfully concluded by the end of the course. A model programme of the workshops appears as Annex II.

Comprehensive written material was prepared for all participants at the workshop, consisting of detailed lecture notes on the statistical methods covered, including the results of their use on real data sets. The lecture notes distributed are attached to this report as Annex III.

### 3. EVALUATION OF THE WORKSHOPS

The evaluation of the workshops was based on an anonymous questionnaire which was filled in by all participants at the end of the workshops. The questions covered:

- i) the adequacy and accuracy of the pre-course information,
- ii) the arrangements for accommodation and course facilities,
- iii) the quality and understandability of the lectures and other presentations,
- iv) the quality of the accompanying written material,
- v) the overall importance and success (or otherwise) of the course.

The most obvious feature of the results, in all cases, was the very high level of satisfaction that participants expressed with virtually all aspects of the course; the overall response was in fact very positive and enthusiastic.

Questions on prior notification of the course content, levels of expertise expected etc attracted excellent ratings except in one of the national workshops in this series. The institution facilities were considered good or acceptable. Computing facilities varied depending on the available PCs at each hosting Institution. However, every effort was made to make available one PC for every two participants and one printer for every two PCs. The configuration varied from XT-compatibles to PS/2; maths co-processors were necessary in all cases to speed up the operations. The participants seemed to prefer machines of a more powerful nature while the lecturers preferred an ample number of machines (at least one for every two participants) of basically the same type and set up in the same way. Another point of interest was the computing support from local staff, when occasional hardware or installation problems were encountered, which was truly excellent in most of the cases. Installation and operation of the programmes was straightforward and comparatively trouble-free.

Some of the participants thought that the computer programmes were not easy to use. This partly results from the fact that they had previously no experience of PCs but also, one suspects, reflects the fact that those with some previous experience (the majority) were more used to a Windows-type interface (a GUI - graphical user interface) which is becoming the "norm" in mass-market PC software.



The Introductory Statistics lectures, covering univariate statistical techniques such as t-tests, ANOVA, regression etc were regarded as very important by nearly two-thirds of the participants, and useful by all. The STATGRAPHICS package, used to perform these analyses, was also considered an essential part of the course by nearly all people. In the sessions of analysing the participants' own data, several univariate analyses were important, particularly higher-way analysis of variance and regression/correlation analyses.

The multivariate statistics lectures were clearly very well received. Both parts of the written lecture notes were also clearly very satisfactory. The multivariate programmes covered by the PRIMER package were considered as fulfilling their data requirements "well" (one-quarter said "averagely"). This, along with other answers, reflects the careful selection of participants most of whom had strong interest in analysing data sets of the type covered in the courses.

In a few instances, participants thought that the course had packed too much material into too short a time and in certain cases, language was identified as a barrier in absorbing so much material in a limited period.

Some of the comments made by participants were, that:

- a) FAO/UNEP should assist laboratories to purchase a PC and copies of the software packages
- b) the workshop should last longer so that sufficient time would be available for running programmes and for revising what had been learnt, at the end of each day.
- c) FAO/UNEP should organise additional workshops on topics such as modelling of ecosystems and time series
- d) feedback with the lecturers/demonstrators was excellent, with the discussions on individuals' data sets being particularly useful.

#### 4. CONCLUSIONS

Unquestionably, this series of workshops was ambitious in the scope and quantity of its material but, by any standards of assessment, it must be regarded as a substantial success. These workshops enabled 108 Mediterranean scientists to acquaint themselves with the recent techniques in analysing statistically and interpreting marine community data.

For both parts of the course, the lectures, written material, and programmes and examples presented in the practical sessions, were clearly highly regarded, and the whole course considered to be very well-structured.

Language difficulties were not the prominent feature anticipated, and though the participants had widely differing past experience of statistical methods, the courses seemed to settle quickly into a speed and level of exposition fitting the median requirements.

In all but a couple of instances, the participants were exceptionally well-motivated, very keen, prepared to work long and taxing days, and able to take on board some very complex concepts. Though the general level of previous experience of statistics and microcomputing,

and the facility with spoken and written English, were variable, nonetheless all participants seemed to feel that they had acquired relevant knowledge and experience from the two-week period. The maintenance of such a high level of interest must be attributed principally to the relevance of the workshop material to the data analysis needs of the participants. An important factor here was the practical nature of the workshop (in keeping with the underlying philosophy of IOC/GEEP that advances in understanding of the biological effects of pollutants come through practical application of methodologies rather than purely theoretical discussion of them). An important motivating element was the allowance of time within the programme for participants to analyse their own data sets; by the end of the workshop all participants who had brought data were able to make at least an initial analysis and interpretation of (possibly a subset of) their data.

Several interesting discussions of participants' own research data were initiated and it is anticipated that this should result in an improved quality of reporting and publication of their work. The lecturers felt that not only had they made good contacts during the courses but that mutual research interests were explored so that greater collaboration could be expected in the future.

Lastly, there was a widespread (and unanticipated) call from the participants for a follow-up meeting to discuss experiences in using these statistical analyses, and possibly to be updated on any further research or software developments, perhaps arising from the further planned development and practical testing activities of IOC/GEEP.

**ANNEX I**

**LIST OF PARTICIPANTS**

- 1) *ABELSON A.*, Department of Zoology, Tel-Aviv University, Ramat-Aviv, Israel
- 2) *AGIOVLASSITI O.*, Ministry of Agriculture, Department of Aquaculture, Athens, Greece
- 3) *ANGEL D.*, Israel Oceanographic and Limnological Research Ltd. National Center for Mariculture, Eilat, Israel
- 4) *AVCIN A.*, Marine Biological Station, Piran, Slovenia
- 5) *AVIV R.*, Israel Limnology Oceanography Research, Haifa, Israel
- 6) *BAKALEM A.*, Institut des Sciences de la Mer et de l'Aménagement du Littoral (ISMAL), Alger, Algeria
- 7) *BARANOVIC A.*, Institute of Oceanography and Fisheries, Split, Croatia
- 8) *BASSO D.*, Università degli Studi di Milano, Dipartimento di Scienze della Terra, Sezione di Geologia e Paleontologia, Milano, Italy
- 9) *BEI F.*, National Centre for Marine Research, Athens, Greece
- 10) *BEN-ELIAHU N.*, Life Sciences Institute, Department of Systematics, Ecology and Evolution, The Hebrew University of Jerusalem, Jerusalem, Israel
- 11) *BEN-ZION M.*, Department of Chemistry, Bar-Ilan University, Ramat-Gan, Israel
- 12) *BRIKNER I.*, Department of Life Sciences, Bar-Ilan University, Ramat-Gan, Israel
- 13) *CATSIKI A.V.*, National Centre for Marine Research, Athens, Greece
- 14) *CHAFFAI HAMZA A.*, Laboratoire de l'Environnement et de la Physiologie des poissons, Département de Biologie, Faculté des Sciences de Sfax, Sfax, Tunisia
- 15) *CHRISTAKI U.*, National Centre for Marine Research, Athens, Greece
- 16) *CHRISTOU E.*, National Centre for Marine Research, Athens, Greece
- 17) *DANIELIDIS D.*, Department of Biology, University of Athens, Athens, Greece
- 18) *DEL PIERO D.*, Laboratory of Marine Biology, Department of Biology, University of Trieste, Trieste, Italy
- 19) *DOLEV Y.*, Department of Ecology, Systematics and Evolution, The Hebrew University of Jerusalem, Jerusalem, Israel

- 20) *DULCIC J.*, Institute of Oceanography and Fisheries, Split, Croatia
- 21) *EISA S.A.*, National Institute of Oceanography and Fisheries, Ministry of Scientific Research and Technology, Alexandria, Egypt
- 22) *EL-EZZ S.A.*, National Institute of Oceanography and Fisheries, Ministry of Scientific Research and Technology, Alexandria, Egypt
- 23) *EL-KOMI M.*, National Institute of Oceanography and Fisheries, Ministry of Scientific Research and Technology, Alexandria, Egypt
- 24) *EL-RASHIDY H.*, Alexandria University, Department of Oceanography, Alexandria, Egypt
- 25) *EL-SAYED RAMADAN Sh.*, National Institute of Oceanography and Fisheries, Alexandria, Egypt
- 26) *EL-SHERIF Z.*, National Institute of Oceanography and Fisheries, Ministry of Scientific Research and Technology, Alexandria, Egypt
- 27) *EL-ZIANI I.S.*, Marine Biology Research Centre, Tripoli, Libya
- 28) *FANUKO N.*, Marine Biological Station, Piran, Slovenia
- 29) *FARSTEI B.*, Interuniversity Institute of Eilat, Eilat, Israel
- 30) *FERNANDEZ de PUELLES M.L.*, Spanish Institute of Oceanography, Balears Laboratory, Palma de Mallorca, Spain
- 31) *FONDA UMANI S.*, Laboratory of Marine Biology, Department of Biology, University of Trieste, Trieste, Italy
- 32) *FRAGOPOULOU N.*, Faculty of Science, Laboratory of Zoology, Department of Biology, University of Patras, Patras, Greece
- 33) *FRASCHETTI S.*, Institute for Marine Environmental Sciences, University of Genoa, Genova, Italy
- 34) *FURLAN L.*, Laboratory of Marine Biology, Department of Biology University of Trieste, Trieste, Italy
- 35) *GAL G.*, Interuniversity Institute of Eilat, Eilat, Israel
- 36) *GATENO D.*, Department of Ecology, Systematics and Evolution, The Hebrew University of Jerusalem, Jerusalem, Israel
- 37) *GHASEM A.S.*, Marine Biology Research Centre, Tripoli, Libya
- 38) *GLAMUZINA B.*, Institute of Oceanography and Fisheries, Split, Croatia
- 39) *GOLANI D.*, Department of Zoology, The Hebrew University of Jerusalem, Jerusalem, Israel
- 40) *GOREN R.*, Department of Zoology, Tel-Aviv University, Tel-Aviv, Israel

- 41) *HADJICHRISTOPHOROOU M.*, Department of Fisheries, Ministry of Agriculture and Natural Resources, Nicosia, Cyprus
- 42) *IGNATIADES L.*, Institute of Biology, National Research Center "Democritus", Athens, Greece
- 43) *JAKLIN A.*, "Rudjer Boškoviæ" Institute, Center for Marine Research Rovinj, Croatia
- 44) *JASPRICA N.*, Biological Institute, Dubrovnik, Croatia
- 45) *JUG DUJAKOVIC J.*, Institute of Oceanography and Fisheries, Split, Croatia
- 46) *JUSTIC D.*, Department of Zoology, University of Zagreb, Zagreb, Croatia
- 47) *KATSILIERI Ch.*, National Centre for Marine Research, Athens, Greece
- 48) *KARAKASSIS I.*, Department of Biology, University of Crete, Crete, Greece
- 49) *KARYDIS M.*, Department of Environmental Studies, University of the Aegean, Mytilene, Greece
- 50) *KOCATAS A.*, Institute of Hydrobiology, Faculty of Science, EGE University, Bornova-Izmir, Turkey
- 51) *KORAY T.*, Institute of Hydrobiology, Faculty of Science, EGE University, Bornova-Izmir, Turkey
- 52) *KRALJEVIC M.*, Institute of Oceanography and Fisheries, Split, Croatia
- 53) *LAKKIS S.*, Conseil National de la Recherche Scientifique, Centre de Recherches Marines, Jounieh, Lebanon
- 54) *LEGAKIS A.*, Department of Biology, University of Crete, Crete, Greece
- 55) *LUCIC D.*, Biological Institute, Dubrovnik, Croatia
- 56) *MAHMOUD A.*, Alexandria University, Department of Oceanography, Alexandria, Egypt
- 57) *MALEJ A.*, Marine Biological Station, Piran, Slovenia
- 58) *MARASOVIC I.*, Institute of Oceanography and Fisheries, Split, Croatia
- 59) *MARSIC J.*, Institute of Oceanography and Fisheries, Split, Croatia
- 60) *MILANI L.*, Laboratory of Marine Biology, Trieste, Italy
- 61) *MILLIOU H.*, Zoological Laboratory, University of Athens, Athens, Greece
- 62) *MIZRAHI L.*, Department of Life Sciences, Bar Ilan University, Ramat-Gan, Israel

- 63) *MOROVIC M.*, Institute of Oceanography and Fisheries, Split, Croatia
- 64) *MOZETIC P.*, Marine Biological Station, Piran, Slovenia
- 65) *MUTLU E.*, Institute of Marine Sciences, Middle East Technical University, Erdemli, Turkey
- 66) *NAHHAS R.*, Marine Research Centre, Latakia, Syria
- 67) *NEUBERGER-CYWIAKL.*, Department of Life Sciences, Bar Ilan University, Ramat-Gan, Israel
- 68) *NICOLAIDOU A.*, Zoological Laboratory, University of Athens Greece
- 69) *PAGOOU K.*, National Centre for Marine Research, Athens, Greece
- 70) *PANAYOTIDIS P.*, National Centre for Marine Research, Athens, Greece
- 71) *PANCUCCI-PAPADOPOULOU M.A.*, National Centre for Marine Research, Athens, Greece
- 72) *PAPASSISSI Ch.*, Zoological Laboratory, Department of Biology, University of Athens, Athens, Greece
- 73) *PAPATHANASSIOU E.*, National Centre for Marine Research, Athens, Greece
- 74) *PETRAKIS G.*, National Centre for Marine Research, Athens, Greece
- 75) *PISANTY S.*, Ministry of Agriculture, Department of Fisheries, Israel
- 76) *PLAOT I.*, Interuniversity Institute of Eilat, Eilat, Israel
- 77) *PLETIKOSIC M.*, Institute of Oceanography and Fisheries, Split, Croatia
- 78) *POLITOU Ch.Y.*, National Centre for Marine Research, Athens, Greece
- 79) *RAMDANI M.*, Institut des Pêches Maritimes CASA, Casablanca, Morocco
- 80) *RAVLIÆB.*, Institute of Oceanography and Fisheries, Split, Croatia
- 81) *SARËEVIÆM.*, Institute of Oceanography and Fisheries, Split, Croatia
- 82) *SARTOR P.*, Department of Environmental and Territorial Science, Section Marine Biology, Pisa, Italy
- 83) *SBRANA M.*, Department of Environmental and Territorial Science, Section Marine Biology, Pisa, Italy
- 84) *SCHLICHTER M.*, Israel Oceanographic and Limnological Research Ltd., Kinneret Limnological Laboratory, Tiberias, Israel
- 85) *SHAPIRO J.*, Ministry of Agriculture, Kinneret Fishery Laboratory, Tiberias, Israel

- 86) *SHOKRY H.*, National Institute of Oceanography and Fisheries, Ministry of Scientific Research and Technology, Alexandria, Egypt
- 87) *SIKAVARA E.*, Department of Biology, University of Crete, Crete, Greece
- 88) *SIMBOURA N.*, National Centre for Marine Research, Athens, Greece
- 89) *SIOKOU-FRANGOU I.*, National Centre for Marine Research, Athens, Greece
- 90) *SISKO M.*, Institute of Biology, University of Ljubljana, Ljubljana, Slovenia
- 91) *SOLIC M.*, Institute of Oceanography and Fisheries, Split, Croatia
- 92) *SONIN O.*, Ministry of Agriculture, Fishing Technology Unit, Kiryat Motzkin, Israel
- 94) *TAFAS T.*, Section of Ecology and Systematics, Department of Biology, University of Athens, Athens, Greece
- 95) *TAKVAM A.*, Institute of Oceanography and Fisheries, Split, Croatia,
- 96) *THOMAS G.H.*, National Institute of Oceanography and Fisheries, Ministry of Scientific Research and Technology, Alexandria, Egypt
- 97) *TRAVIZI A.*, "Rudjer Boškoviæ" Institute, Center for Marine Research Rovinj, Croatia
- 98) *TURKI S.*, Institut National Scientifique et Technique d'Océanographie et de Pêche, Tunis, Tunisia
- 99) *TUSNIK P.*, Marine Biological Station, Piran, Slovenia
- 100) *VAGO R.*, Department of Life Sciences, Bar-Ilan University, Ramat-Gan, Israel
- 101) *VASSILOPOULOU V.*, National Centre for Marine Research, Athens, Greece
- 102) *VILIËIÆD.*, Biological Institute, Dubrovnik, Croatia
- 103) *VOREADOU C.*, Department of Biology, University of Crete, Crete, Greece
- 104) *ZACHARIADOU T.*, National Centre for Marine Research, Athens, Greece
- 105) *ZAGHLOUL F.*, National Institute of Oceanography and Fisheries, Ministry of Scientific Research and Technology, Alexandria, Egypt
- 106) *ZAGOUZI S.W.*, Marine Biology Research Centre, Tripoli, Libya
- 107) *ZAHTILA E.*, "Rudjer Boškoviæ" Institute, Center for Marine Research Rovinj, Croatia
- 108) *ZENETOS A.*, National Centre for Marine Research, Athens, Greece





## **ANNEX II**

### **MODEL PROGRAMME OF THE WORKSHOPS**

#### **First day**

Registration of participants

Opening of the Workshop

I.1 The precision of the mean and confidence limits by J.S. Gray

Coffee break

Introduction to PC-DOS and STATGRAPHICS by M.R. Carr

Lunch break

Practical session on PC-DOS and introduction to STATGRAPHICS

#### **Second day**

I.2 Spatial dispersion of populations and transformations of data by J.S. Gray

Practical session on I.1 and I.2

Coffee break

Practical session on I.2

I.4 Comparing samples: t-test and paired t-test by J.S. Gray

Lunch break

Practical session on I.4

Discussion of arrangements for analysing participants' own data sets

#### **Third day**

I.4 Analysis of variance by J.S. Gray

Practical session on I.4

Coffee break

Practical session on I.4

I.5 Regression and correlation by J.S. Gray

Lunch break

Practical session on I.5

Introduction to entering or reformatting own data sets by E. Papathanassiou

Practical session on entering or reformatting own data

#### Fourth day

I.6 Non-parametric methods by J.S. Gray

Practical session on I.6

Coffee break

General lecture on biological effects and monitoring of pollutants by J.S. Gray

Lunch break

Practical session on own data sets (univariate analysis only)

Question and answer session on basic (univariate) statistics

#### Fifth day

II.1 A framework for studying changes in community structure by R.M. Warwick

II.2 Multivariate methods: measures of similarity of species abundance/biomass between samples by K.R. Clarke

Coffee break

Practical session on lecture II.2 (by hand)

II.3 Multivariate methods: hierarchical clustering by K.R. Clarke

Lunch break

Practical session on lecture II.3 (by hand)

Introduction to multivariate software by M.R. Carr

Practical session on lectures II.2 and II.3 (on computer)

Sixth day

II.4 Multivariate methods: ordination of samples by Principal Components Analysis by K.R. Clarke

Continued practical session on lectures II.2 and II.3

Coffee break

II.5 Multivariate methods: ordination of samples by Multi-Dimensional Scaling (MDS) by K.R. Clarke

Lunch break

Practical session on lectures II.4 and II.5

Seventh day

II.8 Univariate and distributional methods: diversity measures, dominance curves and other graphical analyses by R.M. Warwick

Practical session on lecture II.8

Coffee break

Continued practical session on lecture II.8

II.10 Species aggregation by R.M. Warwick

Lunch break

Practical session on own data sets

Eighth day

II.6/7 Testing for differences between groups of samples, and species contributions by K.R. Clarke

Coffee break

Practical session on lecture II.6/7

Lunch break

II.9 Transformations by K.R. Clarke

Practical session on own data sets

Ninth day

- II.11      Linking community analyses to environmental variables by K.R. Clarke
- Practical session on lecture II.11
- Coffee break
- II.12      Causality: community experiments in the field and laboratory by R.M. Warwick
- II.13      Data requirements for biological effects studies: which components and attributes of the biota to examine?
- Lunch break
- Practical session on own data sets

Tenth day

- II.14      Relative sensitivities and merits of univariate, graphical/ distributional and multivariate techniques by R.M. Warwick
- Practical session on own data sets
- Coffee break
- Practical session on own data sets
- Arrangements for obtaining and mounting multivariate software by M.R. Carr
- Lunch break
- Discussion of participants' own data results
- Question and answer session on multivariate/graphical methods

Note:      The numbering of lectures corresponds to the order of material presented in the Workshop notes: the Part I lecture notes (I.1 to I.6) by Prof. Gray deal with basic (univariate) statistics and the Part II notes (II.1 to II.14) by Drs Clarke and Warwick deal with multivariate community analyses. The other (unnumbered) lectures detailed above are not covered in the formal lecture notes but hand-outs will be available with the lecture, in some instances. A separate set of notes by Mr Carr covers the practical details of using the computer programs for multivariate analysis, developed at the Plymouth Marine Laboratory.

**ANNEX III**  
**LECTURE NOTES**



## INTRODUCTION

This annex is divided into two parts. Part I contains the six introductory lectures and Part II the lectures on multivariate analysis. No lecture notes were available for the informal lectures which did not constitute the core of the workshop but hand-outs were made available. A separate set of notes covering the practical details of using the computer programmes for multivariate analysis was distributed.

The following lectures which deal with some basic and important statistical concepts and their practical application are contained in Part I.

- Lecture 1: The precision of the mean and confidence limits
- Lecture 2: Spatial dispersion of populations and transformations of data
- Lecture 3: Sampling and sub-sampling
- Lecture 4: Comparing samples: 't' test, paired 't' test and analysis of variance
- Lecture 5: Regression and correlation analyses
- Lecture 6: Non-parametric methods

The following lectures which are principally concerned with graphical and multivariate statistical analysis of community data are contained in Part II.

- Lecture 1: A framework for studying changes in community structure
- Lecture 2: Multivariate methods: measures of similarity of species abundance/biomass between samples
- Lecture 3: Multivariate methods: hierarchical clustering
- Lecture 4: Multivariate methods: ordination of samples by Principal Component Analysis (PCA)
- Lecture 5: Multivariate methods: ordination of samples by Multi-Dimensional Scaling (MDS).
- Lecture 6: Multivariate methods: testing for differences between groups of samples
- Lecture 7: Multivariate methods: species analyses
- Lecture 8: Univariate and distributional methods: diversity measures, dominance curves and other graphical analyses
- Lecture 9: Transformations
- Lecture 10: Species removal and aggregation

Lecture 11: Linking multivariate and univariate community analyses to environmental variables

Lecture 12: Causality: community experiments in the field and laboratory

Lecture 13: Data requirements for biological effects studies: which components and attributes of the biota to examine?

Lecture 14: Relative sensitivities and merits of univariate, graphical/distributional and multivariate techniques.



## **PART I**

## Lecture 1: THE PRECISION OF THE MEAN AND CONFIDENCE LIMITS

There are a few terms that must be understood from the beginning.

$\bar{x}$  - the sample mean  
 $\mu$  - the true population mean

The sample mean is simply the average of a series of samples. For example let us take 5 x 0.1 m<sup>2</sup> samples of a population of bivalves on a beach. We obtain the following data:

7,11,12,8,13  $\bar{x} = 10.2$  and the sample size (n)= 5

We first calculate the variance ( $s^2$ )

$$\begin{aligned}\text{This is } s^2 &= (X^2 - (\bar{X})^2/n) / (n - 1) \\ &= (547 - (51^2/5)) / (4) \\ s^2 &= 6.70\end{aligned}$$

This gives another term the STANDARD DEVIATION (s)

where  $s = \sqrt{s^2}$

$$s = 2.588$$

Often we wish to estimate the total population in a given area and the precision with which we estimate that population.

e.g.  $\bar{x} = 10.125$  taken from a sample area of 100 cm<sup>2</sup>. The total area is 300,000 cm<sup>2</sup>.

Therefore, the population estimate is

$$300,000 * 10.125/100 = 30,375$$


The true population mean ( $\mu$ ) has only one value but the sample mean ( $\bar{x}$ ) has many values

e.g. 9.51, 10.74, 9.82, 10.20, 10.125

and these could all be used to estimate the total population size. If we took enough samples all these estimates could be arranged in a frequency distribution and would give us a normal distribution centered around ( $\mu$ ) the true population mean.

This gives us one of the major rules in statistics; the so-called Central Limit Theorem which states that:

"The means of large random samples from the same population are approximately normally distributed with mean equal to the population mean ( $\mu$ ) and variance near the population variance ( $\sigma^2$ )."



In the above definition the term large is used. In statistical terms large means a sample size of over 30 samples, so-called large sample statistics. Often in marine biology it is impossible to take such large samples and we must be aware that in most cases we operate on SMALL SAMPLE STATISTICS which are NOT necessarily those that are in your P.C. or mainframe computer!

Above, we have used another term, variance. We have the sample variance ( $s^2$ ) and the population variance ( $\sigma^2$ ). Often in statistics one uses the standard deviation called ( $s$ ) for a sample and  $\sigma$  for a population.

But in studies of populations (and samples of populations) there is another term that is often confused with the standard deviation, namely, standard error. This term in relation to populations (and samples) refers strictly to the standard error of the mean.

In the above example we had a series of estimates of the true population mean ( $\mu$ ) which gives :

Standard deviation of sample means called standard error (s.e.)

$$s.e. = \sqrt{\frac{\sigma^2}{n}}$$

More usually, this is estimated from samples as:

$$s.e. = \sqrt{\frac{s^2}{n}} \quad \text{or} \quad \frac{s}{\sqrt{n}}$$

This latter term estimates the error in  $\bar{x}$  as an estimator of  $\mu$ . This is usually written as :  $\bar{x} \pm \text{s.e.}$

For example if  $n = 80$ ,  $\bar{x} = 10.125$ ,  $s^2 = 8.5918$

$$\begin{aligned} \text{s.e.} &= \sqrt{(8.5918/80)} \\ &= 0.3277 \end{aligned}$$

For the first 8 counts only  $\bar{x} = 10.5$ ,  $s^2 = 12.857$

$$\text{s.e.} = 1.268 \text{ i.e. } 4 \times \text{the s.e. of } 80 \text{ counts.}$$

The standard error thus estimates the precision which the sample mean ( $\bar{x}$ ) is an estimate of the true population mean ( $\mu$ ). Only when interested in this estimate should one use standard error. More often, one is interested in an estimate of the variability in the sample mean and here one should use confidence limits. These are usually written as 95 or 99% confidence limits. The limits show the range within which one can be 95 or 99% certain that the true population mean lies.

We will deal with both Large Sample and Small Sample statistics.

#### CONFIDENCE LIMITS FOR LARGE SAMPLES ( $n = > 30$ ).

In a normal distribution 95% of the values lie within 1.96 s.d.'s of the true population mean. Therefore, 95% of the sample means lie within 1.96 s.e.'s of the population mean.

$$\text{i.e. } \bar{x} - 1.96 \text{ s.e. to } \bar{x} + 1.96 \text{ s.e.}$$

$$\text{or } \bar{x} - t/s^2/n \text{ to } \bar{x} + t/s^2/n$$

Here we use the statistical table 't' when the population variance ( $\sigma^2$ ) is unknown and is estimated by the sample variance ( $s^2$ ). For 't', degrees of freedom (d.f. =  $n-1$ ) where  $n = \text{infinity}$  't' = 1.96. For d.f. = 30 't' = 2.04.

$$\bar{x} = 10.125, s^2 = 8.5918, n = 80, 't' \text{ for d.f. } 79 = 1.99$$

$$\text{s.e.} = 0.3277$$

$$\bar{x} = 10.125 \pm 0.3277$$

$$95\% \text{ c.l.} = \bar{x} - 1.99(0.3277) \text{ to } \bar{x} + 1.99(0.3277)$$

$$= 10.125 - 0.6251 \text{ to } 10.125 + 0.6251$$

$$= 9.4730 \text{ to } 10.7770$$

For total population estimate:

$$300,000(9.4730)/100 = 28,419$$

to  $300,000(10.7770)/100 = 32,331$

We can be 95% certain that the true population mean lies between 28,419 and 32,331.

SMALL SAMPLES ( $n < 30$ ).

Here we cannot assume that a normal distribution holds. Instead we use an estimate based on the Poisson fraction:

$$\bar{x} \pm \frac{t_{0.05} s}{\sqrt{n-1}}$$

e.g.  $\bar{x} = 11.273$ ,  $s^2 = 7.415$ , ( $s = 2.723$ ),  $n = 10$ ,

$$t_{0.05,9} = 2.262$$

$$\bar{x} = \pm 2.262 (2.723) / \sqrt{9}$$

$$\bar{x} = \pm 2.0531$$

$$= 9.220 \quad \text{to} \quad 13.325$$

## Lecture 2: SPATIAL DISPERSION OF POPULATIONS AND TRANSFORMATIONS OF DATA

There are three basic types of dispersal:

1. random
2. regular
3. contagious (or aggregated)

These groupings can overlap, for example a contagious distribution can result from randomly distributed individuals with regularly distributed individuals in each group. The investigation of patch structure and patch size becomes important.

The 3 types of dispersal can be characterized statistically and simply since in:

random distributions: variance ( $s^2$ ) =  $\bar{x}$  - Poisson

regular distribution:  $s^2 < \bar{x}$  - Positive binomial

contagious distribution:  $s^2 > \bar{x}$  - Negative binomial

The mathematical distributions on the r.h.side are those which can be applied to the respective distributions. In fact the positive and negative binomial models are just one of many types that could be fitted.

### 1. Random distribution.

Statistically it is not in fact possible to test for this distribution! Yet one of the requirements before being able to do statistical tests is that one has a normal (i.e. random) distribution. One simply has to use a test and say that the hypothesis of randomness is not disproved.

A random distribution results from a) chance effects or b) the influence of a single environmental factor. Usually in nature environmental factors do not affect populations randomly and there are a multiplicity of factors which act in concert. So in nature random distributions are in fact rare.

Randomness is often produced by inefficient sampling (e.g. wrong sample size where quadrat is much larger or smaller than the average size of a patch and the patches are randomly distributed then the population will appear randomly distributed). Similarly, if the population density is low it is extremely hard to detect patches. As populations get older they tend to become more random as patches are split up. Here randomness may apply.

Tests for randomness.

The Index of Dispersion  $I = s^2 (n-1) / \bar{x}$

This is tested for significance by reference to the Chi square table.

Example 1: Counts 14, 15, 12, 7, 8, 14, 11, 14, 10, 9, 10

$$\bar{x} = 11.273 \quad s^2 = 7.415 \quad n = 11$$

$$s^2 (n - 1) / \bar{x} = 7.415 (10) / 11.273 \\ = 6.578$$

From chi square table  $p_{0.05} (10) = 18.387$

Conclusion: we cannot reject the hypothesis that the counts have a random distribution.

Example 2: Counts 98, 22, 72, 214, 67

$$\bar{x} = 94.6 \quad s^2 = 5202.8 \quad n = 5$$

$$I = 5202.8 (4) / 94.6 \\ = 219.99$$

Chi square  $p_{0.05} (4) = 9.488$

Conclusion: we reject the hypothesis that the counts come from a random distribution. As the variance / mean ratio is much greater than one a contagious distribution occurs.


Often one wishes to know the size of the patches. One practical way to do this is to use quadrats of increasing sample size. The maximum variance will be found when patch size and quadrat size are equal.

Morisita's index (M).

Here one must keep doubling quadrat size and then compare the ratio of the index (M) of the smaller to the larger quadrat.

$$M = n (\Sigma(x^2) - \Sigma x) / (\Sigma x)^2 - \Sigma x$$

Ratio = M for quadrat size q / M for quadrat size 2q



In the above example patches are at 8q cm<sup>2</sup> with larger patches at 256q cm<sup>2</sup>.

Many other indices of aggregation have been produced and Elliott (1971) has a good coverage of this topic.

Random distributions are rather rare in nature. Patchiness seems to be the rule. This has consequences for quantitative statistical analyses of most field data as there are a series of rules that data sets must conform to before one can apply statistical techniques. These are that:

1. The data follow a normal distribution
2. The variance of the sample is independent of the mean
3. Components of the variance should be additive.


For most biological data patchiness is the general pattern so that rule 1 does not apply and usually the variance increases with the mean. Rule 3 is particularly important when applying the analysis of variance, which we will treat later.

One way of overcoming the problems above that the rules are broken is to TRANSFORM the raw data. There are a number of different transformations that are commonly applied such as the square root transformation or log transformation. These will be dealt with more fully under the multivariate analysis section. Probably the most widely used is the log transformation. Often in biological sampling there are frequently zeros in the data set. One cannot take the log of 0 so in such cases a transformation is used called  $\log_{10}(n + 1)$ , where 1 is added to every number in the set so that when transformed the log of 1 is 0.

The log transformation usually achieves two things a) it normalizes the data and b) it renders the variance independent of the mean. That this is achieved should be tested, but is rarely done and is simply assumed. Let us examine an example.




Samples	A	B	C	D	E	F
	0	3	1	6	7	12
	2	1	5	1	2	7
	1	1	2	5	6	10
	0	1	0	7	9	15
	0	4	2	4	5	9
	1	0	5	1	2	6
	1	1	2	6	7	13
	0	4	1	5	6	11
	1	3	3	3	4	8
	1	3	4	3	3	7
	0	5	1	5	5	10
	2	3	3	3	3	8
	1	2	2	4	6	11
	0	2	4	3	4	8
	0	1	0	8	8	14
	2	1	3	4	5	9
	3	2	4	2	2	6
	0	2	4	2	3	7
	1	2	3	4	4	9
	1	0	6	2	1	5
$\bar{x}$	0.85	2.05	2.75	3.90	4.60	9.25
$s^2$	0.77	1.84	2.83	3.67	4.78	7.57



Here mean and variance are equal but variance increases with mean (rule 2 broken) so we must transform

	A	B	C	D	E	F
$\text{Log}_{10} (n + 1)$	0.22	0.43	0.53	0.65	0.71	0.99
$s^2_y$	0.04	0.05	0.06	0.03	0.03	0.02



So variance and mean are no longer proportional.

Confidence limits where log transformation is used.

The mean of a transformed count ( $\bar{y}$ ) =  $\Sigma \log x / n$  and the variance  $s_y^2$  is calculated in the usual way but on the log-transformed data

95% confidence limits are  $\bar{y} \pm t s_y^2 / n$

But when one back transforms the confidence limits become \*and / not + and -.

This gives a GEOMETRIC (or derived) MEAN which is always smaller than the arithmetic mean.

e.g. Counts 98, 22, 72, 214, 67

$$\bar{x} = 94.60 \quad s^2 = 5202.80$$

$$\text{Log transformed } \bar{y} = 1.8695 \quad s_y^2 = 0.1268$$

From the 't' table  $t_{0.05} (4) = 2.776$

$$\begin{aligned}
 \text{Confidence limits are } \bar{y} &= \pm t (\sqrt{s_y^2 / n}) \\
 &= 1.8695 \pm 2.776 \sqrt{(0.1268/5)} \\
 &= 1.8695 \pm 0.4419 \\
 &= 1.4276 \text{ to } 2.3114
 \end{aligned}$$

Antilogs give  $\bar{y} = 74.05$  with 95% c.l. 26.77 to 204.83

Alternatively  $74.05 \times / 2.77 = 27$  to 205

e.g.  $\text{Log}_{10} (\bar{x} + 1)$  Counts 0, 3, 9, 10  $\bar{x} = 5.5$

$$\bar{y} = 0.6609 \quad s_y^2 = 0.2333 \quad t_{0.05} (3) = 3.182$$

$$\begin{aligned}\bar{y} &= \pm t \quad (s_y^2 / n) \\ &= 0.6609 \pm 3.182 \sqrt{(0.2333 / 4)} \\ &= 0.6609 \pm 0.7685 \\ &= -0.1076 \text{ to } 1.4295\end{aligned}$$

Geometric mean = antilog  $\bar{y} - 1 = 4.58 - 1 = 3.58$

95% c.l. (antilog  $-0.1076$ )  $-1 = < 0$

(antilog  $1.4295$ )  $-1 = 26.88 - 1 = 25.88$

Geometric mean = 3.58 with c.l. from 0 to 25.88!

One should note that percentages are not normally distributed and where one has percentages between 0 and 30 or/and 70 and 100 a transformation is necessary. Physiological data often contains percentage data and only rarely have I seen transformations properly employed.

The transformation to be used here is the ANGULAR (OR ARCSINE) TRANSFORMATION.

The transformation involves replacing the percentage (p) by the angle whose sine is p. Tables are to be found in most statistical books e.g. Snedecor and Cochran (1967), Rohlf and Sokal (1981).

For example 10% is transformed to 18.44, 15% to 22.79, 30% to 33.21 etc.

Lecture 3: SAMPLING AND SUB-SAMPLING

Most sampling methods used in marine habitats are aimed to give an estimate of the size of a given population or populations, (e.g. the plankton or benthos where it is not possible to count all individuals). But how does one take the sample? The simplest method is to take a RANDOM sample but is this the most efficient method? The answer is "it is not" but let me try and demonstrate why.

Let us imagine we cover a whole area with 6 samples and find the following counts

Sample	A	B	C	D	E	F	
Nos.	1	2	4	6	7	16	Total = 36

If we draw only 3 samples from this what are our population estimates?

Samples	Sample Total (T)	Estimate of popn. (x) (T * 2)	Estimate of error (36 - x)
ABC	7	14	-22
ABD	9	18	-18
ABE	10	20	-16
ABF	19	38	2
ACD	11	22	-14
ACE	12	24	-12
ACF	21	42	6
ADE	14	28	-8
ADF	23	46	10
AEF	24	48	12
BCD	12	24	-12
BCE	13	26	-10
BCF	22	44	8
BDE	15	30	-6
BDF	24	48	12
BEF	25	50	14
CDE	17	34	-2
CDF	26	52	16
CEF	27	54	18
DEF	29	58	22
Mean	18	36	0

As a measure of the accuracy of sampling we can use the MEAN SQUARE ERROR (M.S.E.).

$$\begin{aligned} \text{This is M.S.E.} &= \Sigma(\text{error estimate})^2 / \text{sample size (n)} \\ &= (22^2 + 18^2 + 16^2 + \dots + 22^2) / 20 \\ &= 3504 / 20 = 175.2 \end{aligned}$$

This gives a standard error of  $\sqrt{175.2} = 13.2$

i.e  $36 \pm 13.2$  (or 37%)

The sampling plan adopted (3 random samples) is not therefore very efficient.

If we know something about the populations we can improve accuracy. Say we knew that F would give higher values than the other samples. Wherever F occurs in a sample we get much higher values. So the strategy is to divide the area into two STRATA, stratum 1 (S1) with F and stratum 2 (S2) without F. The tactic now is then to always include F but take the other two samples at random.

Sample		Sample total in Stratum 2(T2)	Estimate (16 + 2.5 T2)	Error of Estimate
S2	S1			
AB	F	3	23.5	-12.5
AC	F	5	28.5	- 7.5
AD	F	7	33.5	- 2.5
AE	F	8	36.0	0
BC	F	6	31.0	- 5.0
BD	F	8	36.0	0
BE	F	9	38.5	2.5
CD	F	10	41.0	5.0
CE	F	11	43.5	7.5
DE	F	13	48.5	12.5
Mean			36.0	0

The estimate for S1 total is always correct: 16. The estimate for S2: there are 2 out of 5 samples therefore we multiply T2 by 2.5.

M.S.E. is now  $487.5 / 10 = 48.75$

standard error = 7.0 (or 19% of total)

So the s.e. is much improved from 37% to 19%. In general STRATIFIED RANDOM sampling is much the preferred strategy. It does require however that one know something about the area or populations being sampled. This implies that one should first do preliminary surveys before setting out on detailed quantitative sampling programmes.

An example showing how one plans a stratified random sampling programme for a benthic survey follows. We plan to sample an area of 200 m<sup>2</sup> with a grab taking an area of 0.05 m<sup>2</sup>. Potentially therefore, there are 200/0.05 = 4,000 sampling units within the area. We do a preliminary survey and find that the bottom is very heterogeneous. Since we know nothing about the benthic fauna we suspect that grain size variations could be important so we map the sediment. Then we want to sample with equal intensity on each type of bottom. This is called PROPORTIONAL ALLOCATION of samples. Let us plan to give an even coverage of 10% to each area i.e. 40 samples total, a not unreasonable number.

We find gravel (n1) covers 1000 sampling units, coarse sand (n2) 500, sand (n3) 1500, fine sand (n4) 800 and mud (n5) 200, totalling 4000 sampling units.

We then allocate our 40 samples in proportion

$$n1 = 1000 * 40 / 4000 = 10 \text{ samples}$$

$$n2 = 500 * 40 / 4000 = 5 \text{ samples}$$

$$n3 = 1500 * 40 / 4000 = 15 \text{ samples}$$

$$n4 = 800 * 40 / 4000 = 8 \text{ samples}$$

$$n5 = 200 * 40 / 4000 = 2 \text{ samples}$$

As to the placement of samples within the area ideally we divide up the whole area give each potential sampling unit a number and pick the numbers from random number tables from a book of statistical tables.

Another method of allocating samples within a stratified random approach is called OPTIMAL ALLOCATION where one takes more samples where there is high variability. As a simple method one can allocate samples according to the variability of the standard error. Let us take an example:

In this example the area was divided up first according to sediment types using methods similar to those shown above and a preliminary sampling done. The total number of animals found in this preliminary survey taking 7 replicates per station were:

Stratum	Station	Replicates						
		A	B	C	D	E	F	G
1	1	1020	1180	1300	2100	980	900	1050
	2	390	490	210	360	220	310	150
	3	140	440	360	150	490	1070	920
2	4	140	150	190	160	150	180	140
	5	420	950	350	150	180	330	150
3	6	370	420	700	100	200	190	220
	7	620	1390	380	450	480	2600	870

Stratum	Station	A	B	C	D	E	F	G
4	8	390	430	110	440	110	180	160
	9	40	20	350	60	80	20	50
	10	150	140	660	320	240	880	1660
	11	730	670	470	340	930	370	410
5	12	1380	1410	1190	2710	1600	1290	530
	13	1620	320	1550	760	1250	1990	270
	14	1850	2060	1090	2410	1520	220	1620

Using Statgraphics calculate the means, standard deviation and standard errors for this data. This gives:

Stratum	Samples size (n)	Mean	s.d	s.e
1	21	677.62	504.62	110.12
2	14	260.00	218.32	58.35
3	14	642.14	654.90	175.03
4	21	309.05	381.59	83.27
5	28	1162.86	686.68	129.77
Total				556.54

Calculate each s.e. as a proportion of the Total s.e. and use this proportion to calculate sampling allocation per stratum. Here it is assumed that a total of 65 samples can be taken in the next survey.

Stratum	Proportion of Total s.e.	No. of samples/stratum
1	0.198	13
2	0.104	7
3	0.315	20
4	0.149	10
5	0.232	15

We can then calculate how effective this sampling system has been, compared with random sampling. First run an analysis of variance (using Statg) on the data testing within strata variance compared with between strata variance. This gives:

Source of variation	Sum of Squares	d.f.	Mean Square	F ratio
Between strata	11902293	4	2975573.3	10.27
Within strata	26931369	93	289584.6	
Total	38833662	97	400347.0	

The pooled standard deviation within strata  $s_w$  is:

$$s_w = \sqrt{289584.6}$$

$$s_w = 538.13$$

$$\begin{aligned} \text{Estimated s.e. of } Y_{st} &= s(Y_{st}) \\ &= s_y / \sqrt{n} \quad n = 98 \\ &= 538.13 / \sqrt{98} \\ &= 54.36 \end{aligned}$$

With purely random sampling  $S_y = s / \sqrt{n}$

$$\begin{aligned} S_y &= \sqrt{400347} / \sqrt{98} \\ &= 63.91 \end{aligned}$$

Therefore the stratified sampling reduces the s.e. by  
 $((63.91 - 54.36) * 100) / 63.91 \%$   
 $= 14.9\%$

This is a big change and again illustrates the advantages of stratified sampling.

#### Size of sample.

In general, small sample sizes are better than large ones because:

- i) more small units can be taken with the same counting effort.
- ii) more samples gives a greater number of degrees of freedom for statistical tests and therefore, a reduction in error.
- iii) many small samples in a given area will cover more ground and be more representative than few large samples.

But size reduction must not go too far otherwise edge effects will occur where the population is underestimated due to the disturbance of the edge of the sampler. So a compromise is necessary. Do NOT assume however, that a given grab or plankton net is the appropriate size for a given population simply because it is available. Most grabs were developed as fractions of 1 m<sup>2</sup> and this may be a quite inappropriate size for a given population.

#### Number of samples.

As we have established most species are not randomly or regularly distributed but aggregated. If only a small number of samples is taken from an aggregated population then the population estimate will be highly inaccurate.

One of the simplest methods to determine the number of samples that should be taken is to take 5 samples and calculate the mean and variance, take 5 more and calculate the mean and variance for all ten samples and repeat until the mean and variance are stable. The minimum number of samples where this is achieved is the correct number.

An alternative is to decide on an acceptable error of one's estimate of the population mean and use this in the following equation:

Let us assume that 10% error is acceptable and call this proportion (0.1) D. The number of samples that should be taken (n) is:

$$\begin{aligned} n &= s^2 / (0.1)^2 \bar{x}^2 \\ &= 100 s^2 / \bar{x}^2 \text{ for a 10\% accepted error.} \end{aligned}$$



Example: counts: 14,15,12,7,8,14,11,14,10,9,10

$$s^2 = 7.42 \quad \bar{x} = 11.273$$

$$n = 100 * 7.42 / (11.273)^2 \\ = 5.82 \text{ i.e. 6 samples}$$

For an aggregated distribution:

Counts: 98,22,72,214,67

$$s^2 = 5202.8 \quad \bar{x} = 94.60$$

$$n = 100 * 5202.8 / (94.6)^2 \\ = 58$$

This is an enormous number of samples and is clearly impractical. So accept a lower error estimate e.g. 20%

$$n = 25 * 5202.8 / (94.6)^2 \\ = 14.53$$

---

### Sub-sampling.

Frequently with plankton samples one must sub-sample to reduce the amount of material obtained to reasonable numbers. There are a number of commercial plankton splitters on the market (and indeed there are some for splitting meiobenthos samples). One must test that these samplers are in fact making random splits of the sample.

Say that one has 4 l of concentrated plankton and that he takes 5 x 50ml subsamples and obtains the following data. Is it a random split?

$$20,25,25,30,40 \quad s^2 = 57.5 \quad \Sigma x = 140 \quad n = 5$$

In a random distribution  $s^2 / \bar{x} = 1$

$$\text{We use the Index of Dispersion test } I = s^2 (n-1) / \bar{x} \\ = 57.5 * 4 / 28 \\ = 8.2$$

Chi square for  $p_{0.05}$  4 d.f. = 9.49

As the value is less than the tabulated one we cannot reject the hypothesis that the sample comes from a random distribution.

We have sampled  $5 * 50 \text{ ml} = 250$  from 4000 ml i.e 1/16th

The estimated numbers of animals in the 4000ml is:

$$16 * 140 = 2240$$

To obtain 95% confidence limits we look up in a table of confidence limits for a Poisson variable (Biometrika 1959 46, 441-453 copy in appendix)

For 140 we find  $\pm 23$

So the population estimates are  $23 * 16 = 368$

Giving  $2240 \pm 368$ .

If the sub-samples do NOT fit a random distribution then one cannot estimate the numbers in the original sample. In my experience many plankton splitters do not in fact give reliable splits. SO BE WARNED!.

#### Comparing efficiency of a sampler.

Often one wants to know whether the observed catches of a sampler are equally efficient within acceptable limits.

The  $H_0$  is that the samplers are equally efficient.

Example: Samplers 1 2 3 4 5  
Counts 6, 8, 16, 5, 18  $\Sigma = 53; n = 5$

$$\text{Expected count} = 53 / 5 = 10.6$$

$$\begin{aligned} \text{Chi}^2 &= (\text{Observed} - \text{Expected})^2 / \text{Expected} \\ &= (6-10.6)^2 / 10.6 + \dots + (18-10.6)^2 / 10.6 \\ &= 13.51 \end{aligned}$$

$$\text{d.f.} = n-1 = 4, p0.05_{(4)} = 9.49$$

Conclusion: As the calculated value is greater than the Table value we reject  $H_0$  that the samplers are equally efficient. This type of test has many variants and is widely used.

Lecture 4: COMPARING SAMPLES: 't' TEST, PAIRED 't' TEST AND ANALYSIS OF VARIANCE

Often one wants to compare the variability between two samples. A simple and illustrative test is the Coefficient of Variation (C).

$$\text{Where } C = s (100) / \bar{x}\%$$

The coefficient of variation is scaled for differences in mean and is a widely used descriptive parameter.

Another commonly used test is that of comparing two means. The null hypothesis ( $H_0$ ) is that the two means come from the same population and that the means are within the error for that population. It is usual to assume a 5% error due to chance.

In all statistical methods there are two types of error that one can make Type I and Type II errors.

TYPE I ERROR - where one rejects  $H_0$  when it is true

TYPE II ERROR - where one accepts  $H_0$  when it was false

All statistical tests are prone to both types of error and there is a greater chance of making one type of error than another in each test. The ideal test is one where the probability of rejecting  $H_0$  when true is small and the likelihood of rejecting  $H_0$  when false is large. Both errors are reduced by increasing the number of degrees of freedom in a test.

Before doing any quantitative statistical test we must make sure that our three primary rules hold. When comparing two means with the 't' test there should be similar variances. If the variances are significantly different then we cannot validly test if the means are significantly different or not.

Here we use the variance ratio test 'F'

where  $F = s_1^2 / s_2^2$  where  $s_1$  is always the largest of the two.

Let us test  $s_1^2 = 8.865$   $n_1 = 60$ ,  $s_2^2 = 7.465$   $n_2 = 80$

$$F = 8.855 / 7.465 = 1.11862$$

Look up the F ratio table for  $n-1 = 59$  and 79 d.f.

$$p_{0.05} \ 60,120 = 1.48$$

Conclusion: Since our value is < the table value we cannot reject the hypothesis that the variances come from the same population and so we can test for differences between means using the 't' test.

Student's 't' test.

a) For large samples from normal distributions.

$$\text{Here 't'} = \bar{x}_1 - \bar{x}_2 / \sqrt{(s_1^2 / n_1 + s_2^2 / n_2)}$$

$$\text{e.g. } \bar{x}_1 = 10.125 \quad s_1^2 = 7.465 \quad n_1 = 80$$

$$\bar{x}_2 = 12.245 \quad s_2^2 = 8.855 \quad n_2 = 60$$

$$\text{'t'} = 12.245 - 10.125 / \sqrt{(7.465/80 + 8.855/60)}$$

$$= 4.3194$$

$$\text{d.f} = n_1 + n_2 - 2 = 80 + 60 - 2 = 138$$

$$\text{'t'} \text{ p}0.05_{(138)} = 1.96$$

Conclusion: Since our calculated value is greater than the table value we reject the hypothesis that the two means come from the same population.

Small samples from contagious distributions.

Example:

$$\bar{x}_1 = 4, 5, 8, 14, 14, 15, 15, 19, 28, 36$$

$$\bar{x}_2 = 2, 4, 5, 7, 12$$

$$\bar{x}_1 = 15.80 \quad s_1^2 = 99.07 \quad n_1 = 10$$

$$\bar{x}_2 = 6.00 \quad s_2^2 = 14.50 \quad n_2 = 5$$

Clearly the variance increases with the mean so that we must transform the data. Let us assume that a  $\log_n$  transformation is adequate. Now we obtain:

$$\bar{y}_1 = 0.602, 0.699, 0.903, 1.146, 1.146, 1.176, 1.176, 1.279,$$

$$1.447, 1.556$$

$$\bar{y}_2 = 0.301, 0.602, 0.699, 0.845, 1.079.$$

$$\bar{y}_1 = 1.0638 \quad s_1^2(y) = 0.2747$$

$$\bar{y}_2 = 0.7052 \quad s_2^2(y) = 0.2887$$

Firstly test the variances:

$$F = 0.2887 / 0.2747 = 1.050$$

$$p_{0.05} \text{ } 9,4 = 8.90$$

Conclusion: Since the calculated value is less than the table value we conclude that the variances are similar and we can proceed with a 't' test.

$$t = 1.113 - 0.705 / \sqrt{(0.0914/10) + (0.0833/5)}$$
$$= 2.497$$

$$d.f = n_1 + n_2 - 2 = 13$$


$$p_{0.05}(13) = 2.16$$

Conclusion: Since the calculated value is greater than the table value at  $p_{0.05}$  we reject  $H_0$  that the means come from the same population and the means are therefore, significantly different.

Making paired comparisons.

Often two sets of data vary over seasons and one is interested not in comparing the overall means but in seeing if there is a significant overall difference, where the null hypothesis is that there is no significant difference between pairs.

The figure illustrates a typical data set.



The data are shown below in tabular form for calculating the Paired 't' test.

Month	No. of individuals		Difference	
	Spp A	Spp B	(D)	D <sup>2</sup>
Jan	12	11	1	1
Mar	56	63	-7	49
Jun	125	107	18	324
Sept	87	78	9	81
Dec	34	36	-2	4
		Total	19	459

$$\bar{D} = 19/5 = 3.8$$

$$\begin{aligned} S_D &= \sqrt{((\sum D^2 - (\sum D)^2 / n) / (n - 1))} \\ &= \sqrt{((459 - 19^2 / 5) / 4)} \\ &= 9.83 \end{aligned}$$

$$\begin{aligned} \bar{S}_D &= S_D / \sqrt{n} \\ &= 9.83 / \sqrt{5} \\ &= 4.396 \end{aligned}$$

$$\begin{aligned} t &= \bar{D} / \bar{S}_D \\ &= 3.8 / 4.396 \\ &= 0.8644 \end{aligned}$$

$$p_{0.05(4)} = 2.776$$

Conclusion: Since the calculated value is less than the tabular value we cannot reject H<sub>0</sub>: there is no difference between species.

### Analysis of variance.

More often than not one is interested in comparing more than two means and here one should use the analysis of variance rather than test two and two means by themselves. The analysis of variance (anova) is one of the most used and robust statistical tests devised. But it requires that the data sets comply to the three rules a) samples normally distributed b) variance independent of the mean and c) components of the variance additive. This latter criterion has probably been a bit of a mystery but now all will be revealed in that the ANOVA test breaks down the sources of variance into their components on the assumption that the components are additive.

Example

Sample	A	B	C	D
	98	12	86	2
	22	13	12	5
	72	46	49	12
	214	38	33	3
	67	49	72	19
$\bar{x}$	94.60	31.60	50.40	8.20
$s^2$	5202.80	320.30	878.30	51.70

Clearly the variance increases with the mean so that we must transform. Use the  $\log_{10}$  transformation.

Sample	A	B	C	D
	1.991	1.079	1.935	0.301
	1.342	1.114	1.079	0.699
	1.857	1.663	1.690	1.079
	2.330	1.580	1.519	0.477
	1.826	1.690	1.857	1.277
Total	9.346	7.126	8.080	3.835
$\bar{x}_s$	1.8692	1.4252	1.6160	0.7670
$s^2$	0.1268	0.0918	0.1158	0.1163

Now the variance is independent of the mean.

Calculate:

1. The Grand Total  $Y = 9.346 + 7.126 + \dots + 3.835 = 28.387$
2. Sum of squared obs.  $= 1.991^2 + 1.342^2 + \dots + 1.277^2 = 45.625$
3. Sum of squared group totals / n  
 $= (9.346^2 + 7.126^2 + \dots + 3.835^2) / n$   
 $= 43.626$
4. Grand Total squared / Total Sample Size (Correction term)  
 $= (28.387)^2 / 20 = 40.291$
5. Sum of squares (Total)  $= (2) - (CT) = 45.625 - 40.291 = 5.3340$
6. S.S. (Groups)  $= (3) - (CT) = 43.626 - 40.291 = 3.3350$
7. S.S. (Within)  $= SS (Total) - SS (Groups) = 5.3340 - 3.3350 = 1.9990$

ANOVA TABLE

Source of Variation	d.f.	SS	MS	F
Between groups	3	3.3350	1.1117	8.9007
Within groups	16	1.9990	0.1249	
Total	19	5.3340		

$$p_{0.05}(3,16) = 3.24$$

$$p_{0.01}(3,16) = 5.29$$

Conclusion: Since the calculated value for F (8.9007) is greater than the table value for  $p_{0.01}$  we conclude that  $H_0$  must be rejected and there are significant differences between samples.

This analysis does NOT however tell us which means are significantly different. We can use a test based on means called the Least Significant Difference (LSD) test where:

$$LSD = t_{(0.05)} \text{ d.f. } (2/n(MS_{\text{within}}))$$

$$\text{Giving: } 2.12 \quad (2/4(0.1249))$$

$$LSD = 0.5297$$

Thus means differing by more than 0.795 are significant at the 0.05 level.

Ranking the means we have: 1.8692, 1.6160, 1.4252, 0.7670

Only D(0.7670) is significantly lower than the other means.

(0.7670+0.5297=1.2568, which is lower than any of the other means and the difference between the other means is less than 0.5297).

TWO WAY ANOVA

This design can be easily extended to a TWO or THREE-WAY ANOVA. Both are easy to calculate but take time. To illustrate the point let us use data on the oxygen consumption of two species of limpet in three concentrations of seawater. The question is there a significant difference in oxygen consumption between species and how does this vary with salinity.

Oxygen figures are in  $\mu\text{g O}_2 / \text{mg dry wt.}/\text{min}$  @ 22°C, n = 8.

FACTOR A SPECIES

FACTOR B SEAWATER	<u>Acmaea scabra</u>	<u>Acmaea digitalis</u>	Total
100%	7.16	8.26	6.14
	6.78	14.00	3.86
	13.60	16.10	10.00
	8.93	9.66	11.60
			5.49
			5.80



		84.49	59.43	143.92
75%		5.20 13.20	4.47 4.95	
		5.20 8.39	9.90 6.49	
		7.18 10.40	5.75 5.44	
		6.37 7.18	11.80 9.90	
		63.12	58.70	121.82
50%		11.11 10.50	9.63 14.50	
		9.74 14.60	6.38 10.20	
		18.80 11.10	13.40 17.70	
		9.74 11.80	14.50 12.30	
		97.39	98.61	196.00
		245.00	216.74	461.74

- 1) Grand Total = 461.74
- 2) Sum of obs.squared =  $(7.16)^2 + (6.78)^2 \dots (12.30)^2 = 5065.1530$
- 3) Sum of squared group totals / sample size of groups  
 $= ((84.49)^2 + (59.43)^2 \dots (98.61)^2) / 8 = 4663.6317$
- 4) Sum of squared column totals / sample size of column  
 $= ((245.00)^2 + (216.74)^2) / 24 = 4458.3844$
- 5) Sum of squared row totals / sample size of row  
 $= ((143.92)^2 + (121.82)^2 + (196.00)^2) / 16 = 4623.0674$
- 6) Grand Total squared / total sample size = C.T.  
 $= (461.74)^2 / 48 = 4441.7464$
- 7)  $SS_{total} (2) - (6) = 5065.1530 - 4441.7464 = 623.4066$
- 8)  $SS_{subgroupd} (3) - (6) = 4663.6317 - 4441.7464 = 221.8853$
- 9)  $SS_A (4) - (6) = 4458.3844 - 4441.7464 = 16.6380$
- 10)  $SS_B (5) - (6) = 4623.0674 - 4441.7464 = 181.3210$
- 11)  $SS_{AB} (8)-(9)-(6) = 2221.8853-16.6380-181.3210 = 23.9263$
- 12)  $SS_{error} = (7) - (8) = 623.4066 - 221.8853 = 401.5213$

Source of variation	df	SS	MS	F
Subgroups	5	221.8853	44.377	
Between species (A)	1	16.6380	16.638	1.74 n.s
Between salinities(B)	2	181.3210	90.660	9.48 ***
Species X salinities (AB)	2	23.9263	11.963	1.25 n.s
Error (within group)	42	401.5213	9.560	
Total	47	623.4066		

$$F_{0.05(1,42)} = 4.07, \quad F_{0.05(2,42)} = 3.22$$

$$F_{0.001(2,42)} = 8.18$$

Conclusion: oxygen consumption does not differ significantly between species, but does between salinities for both species.

This test does not however, tell which means are significantly different from each other. One can use the Least Significant Range (LSR) test.

$$LSR = Q_{0.05(df)} \quad (MS \text{ within} / n)$$

Q is a factor from Tables called the STUDENTIZED RANGE

$$Q_{0.05(2,42)} = 2.858$$

$$LSR = 2.858 \quad (9.560 / 16)$$

$$= 2.2092$$

Arrange means in ascending (or descending) order

	100%	75%	50%
$\bar{x}$	71.96	60.91	98.00


As all the difference between means are greater than the calculated significance level of 2.209 we conclude that all are significantly different.

## Lecture 5: REGRESSION AND CORRELATION ANALYSES

In regression analysis we relate one variable, the dependent (Y) to another, the independent (X). Such techniques are used when one wishes to know

- a) if Y depends on X
- b) to predict Y knowing X
- c) the shape of the relationship between Y and X

Example: Linear Regression analysis



$$\Sigma X^2 = 16125 \quad \Sigma Y^2 = 101341 \quad \Sigma XY = 40155$$

$$(\Sigma X)^2 / n = 15125; \quad (\Sigma Y)^2 / n = 99405; \quad (\Sigma X)(\Sigma Y) / n = 38755$$

$$\Sigma x^2 = (\Sigma X)^2 - (\Sigma X)^2/n = 16125 - 15125 = 1000$$

$$\Sigma y^2 = (\Sigma Y)^2 - (\Sigma Y)^2/n = 101341 - 99405 = 1936$$

$$\Sigma xy = \Sigma XY - (\Sigma X)(\Sigma Y)/n = 40155 - 38755 = 1380$$

$$b = \Sigma xy / \Sigma x^2 = 1380 / 1000 = 1.38$$

$$\hat{Y} = \bar{Y} + b(X - \bar{X}) = 141 + 1.38(X - 55)$$

$$= 65.1 + 1.38x$$

Substitute two values in the equation

$$\text{Let } X = 50, Y = 65.1 + 1.38(50) \\ = 134.1$$

$$\text{Let } X = 70, Y = 65.1 + 1.38(70) \\ = 161.7$$

But how good a fit is this line to the data points?

Here we will calculate the 95% confidence intervals for the line.

Firstly, calculate the deviations from the line

$$d_{y.x} = Y - \bar{Y}$$

For all points this is:

$$\begin{aligned}d_{y.x} &= \Sigma y^2 - (\Sigma xy)^2 / \Sigma x^2 \\ &= 1936 - (1380)^2 / 1000 \\ &= 31.60\end{aligned}$$

Since there are two variables X and Y this deviation has n-2 d.f. = 3.

$$\begin{aligned}S_{y.x}^2 &= 31.60 / 3 \\ &= 10.53\end{aligned}$$

The mean square deviation  $S_{y.x} = \sqrt{10.53} = 3.245$

Now we need to calculate the sample S.D. of the regression coefficient b, i.e.  $S_b$

$$\begin{aligned}S_b &= S_{y.x} / \sqrt{\Sigma x^2} \\ &= 3.245 / \sqrt{1000} \\ &= 0.1026\end{aligned}$$

The significance of the regression can be tested using a 't' test in the usual way where  $t = b / S_b$

$$\begin{aligned}&= 1.38 / 0.1026 \\ &= 13.5\end{aligned}$$

$$p0.05_{(3)} = 3.182$$

Conclusion: The calculated value is greater than the table value. We conclude that since the regression coefficient is significantly greater than the error term there is a significant fit to the data.

Now we must calculate the confidence limits:

$$\begin{aligned}\text{For } b \text{ 95\%} &= t0.05_{(3)} S_b \\ &= 3.182 (0.1026) \\ &= 0.3265\end{aligned}$$

$$b = 1.38 \pm 0.3265$$

$$\begin{aligned} \text{The s.e. of } Y &= S_{yx} \sqrt{\left(\frac{1}{n}\right) + \left(\frac{x^2}{\sum x^2}\right)} \\ &= 3.245 \sqrt{\left(\frac{1}{5} + \frac{x^2}{1000}\right)} \\ &= \sqrt{10.30} \sqrt{0.2 + 0.001x^2} \\ &= 2.06 + 0.0103 x^2 \end{aligned}$$

---

$$\begin{aligned} \text{For } X = 35 \quad x &= X - \bar{x} = 35 - 55 = -20 \\ S_y &= \sqrt{(2.06 + 0.0103 (-20)^2)} \\ &= 2.4859 \end{aligned}$$

---

$$\begin{aligned} \text{For } X = 55, \quad x &= 55 - 55 = 0 \\ S_y &= \sqrt{(2.06 + 0.0103 (0)^2)} \\ &= 1.4352 \end{aligned}$$

---

$$\begin{aligned} \text{For } X = 75, \quad x &= 75 - 55 = 20 \\ S_y &= \sqrt{(2.06 + 0.0103 (20)^2)} \\ S_y &= 2.4859 \end{aligned}$$

---

95% confidence limits are:

$$Y - t_{0.05} S_y \quad \text{to} \quad Y + t_{0.05} S_y$$

$$\begin{aligned} X = 35 \\ \hat{Y} &= Y + bx \\ &= 141 + 1.38(-20) \\ &= 113.4 \end{aligned}$$

$$\begin{aligned} t_{0.05} S_y &= 3.182 (2.4869) \\ &= 7.913 \\ &= 113.4 \pm 7.913 \end{aligned}$$

---

$$\begin{aligned} X = 55 \\ \hat{Y} &= Y + bx \\ &= 141 + 1.38 (0) \\ &= 141 \end{aligned}$$

$$\begin{aligned} t_{0.05} S_y &= 3.182(1.4352) \\ &= 4.567 \\ &= 141 \pm 4.567 \end{aligned}$$

$$\begin{aligned} X = 75 \\ \hat{Y} &= Y + bx \\ &= 141 + 1.38 (20) \\ &= 168.6 \\ &= 168.6 \pm 7.913 \end{aligned}$$

So the confidence limits are asymmetrical, with greater chance of error the further one is from the mean value.

Curvilinear regressions.


For many biological phenomena the relationship between the two variables is not linear but curvilinear. For example the initial phases of growth of populations of bacteria doubling in size at each time interval. Or data on weight increases over time. The regression now is:

$$\text{Weight (W)} = (A)(B^x), \text{ where A and B are constants.}$$

One can use a log transformation to obtain a linear relationship where

$$\text{LogW} = \text{logA} + (\text{logB})X$$

The plots are shown below.



The data are:

Age in days X	Dry weight W (gm)	Log <sub>10</sub> W Y
6	0.029	-1.538
7	0.052	-1.284
8	0.079	-1.102
9	0.125	-0.903
10	0.181	-0.742
11	0.261	-0.583
12	0.425	-0.372
13	0.738	-0.132
14	1.130	0.053
15	1.882	0.275
16	2.812	0.449

The calculations are as in the example for the linear regression using here X and Y, the log transformed data.

$$\text{We obtain } Y = 0.1959X - 2.689$$

One of the most interesting parameters from a relationship such as that above is the relative rate of increase

$$W = Ae^{cx}$$

where  $e = 2.718$  the base of the natural logarithm series.

$$\begin{aligned} \text{This gives us } \text{Log}_e W &= (\text{log}_{10} W)(\text{log}_e 10) \\ &= 2.3026 \text{ log}_{10} W \end{aligned}$$

In terms of the equation above we get:

$$\begin{aligned} W &= (2.3026)(0.1959) \\ &= 0.451 \text{ gm per day per gm} \end{aligned}$$

We must also backtransform the equation

$$\text{Log } W = 0.159X - 2.689$$

$$\text{antilog } -2.689 = \text{antilog } (0.311 - 3) = 0.00205$$

$$\text{Giving: } W = (0.00205)e^{0.451X}$$

---

### Correlation

This is related to regression analysis and shows the degree to which two variables are correlated together. Here there is no dependent and independent variable rather  $X_1$  and  $X_2$  instead of X and Y.

Example

Brothers height  $X_1$  71 68 66 67 70 71 70 73 72 65 66  $\bar{x}_1 = 69$

Sisters height  $X_2$  69 64 65 63 65 62 65 64 66 59 62  $\bar{x}_2 = 64$

$n = 11$ ,  $\sum x_1^2 = 74$ ,  $\sum x_2^2 = 66$ ,  $\sum x_1 x_2 = 39$

Remember:

$$\sum x_{1,2} = \sum X_1^2 - (\sum X_1)^2 / n \quad \text{etc}$$

$$r = \sum x_1 x_2 / \sqrt{(\sum x_1^2)(\sum x_2^2)}$$

$$= 39 / \sqrt{(74)(66)}$$

$$= 0.558$$

$$\text{d.f.} = n - 2 = 11 - 2 = 9 \quad p_{0.05(9)} = 0.602$$

Conclusion: The value is less than the table value. We cannot reject  $H_0$  that there is no correlation between brothers height and sisters height.



## Lecture 6: NON-PARAMETRIC METHODS

So far we have been dealing with analysis of quantitative data. I believe that as far as possible one should use the parametric methods discussed before. There is a tendency within biology today to use the argument that we do not know if such and such a data set is normally distributed nor do we know the relationship between the variance and mean, particularly with small samples with which we usually work. So the decision is taken almost exclusively to use non-parametric statistics as these do not require distributions to be normal, nor variance be independent of the mean. I believe that this is often a mistaken belief in that one misses much of the potential of the data that can only be revealed by a proper parametric analysis.

It must be said, however, that non-parametric methods are in some cases as efficient as parametric ones, save when the analyses are complex such as ANOVA and regression analyses.

Let us examine a few of the most commonly used:

### A non-parametric 't' test the MANN-WHITNEY 'U' test.

This is a highly useful alternative to the 't' test and with almost equal precision. The null hypothesis is that two independent random samples come from the same populations having the same parent distribution and the same means.

Example: Sample 1  $n_1 = 5$  Counts 2,4,5,7,12

Sample 2  $n_2 = 10$  Counts 4,5,8,14,14,15,15,19,28,36

The counts are arranged in rank order from lowest to highest. Now substitute ranks for each count giving an average rank for equal numbers.

This gives:

$$n_1 = 1, 2.5, 4.5, 6, 8$$

$$n_2 = 2.5, 4.5, 7, 9.5, 9.5, 11.5, 11.5, 13, 14, 15$$

Sum the ranks for each sample:

$$R_1 = 22$$

$$R_2 = 98 \quad \text{Check that } R_1 + R_2 = (n_1 + n_2)(n_1 + n_2 + 1)/2$$

Calculate test statistics  $U_1$  and  $U_2$ :

$$U_1 = n_1 n_2 + (n_2(n_2 + 1)/2) - R_2$$

$$U_2 = n_1 n_2 + (n_1(n_1 + 1)/2) - R_1$$

$$U_1 = 50 + (110/2) - 98 = 7$$

$$U_2 = 50 + (30/2) - 22 = 43 \quad \text{Check } U_1 + U_2 = 50 = n_1 n_2$$

Refer to Table of U for  $n_1 = 5$   $n_2 = 10$  at  $p0.05_{(5,10)} = 8$

Conclusion: Here we use an unusual decision technique in that the calculated value must be SMALLER than the table value to be significant.

We find that the smallest (U) value = 7 and since this is smaller than the table value 8, we reject the hypothesis that the two means come from the same population and can conclude that the mean of sample 2 is significantly higher than that of sample 1.

Non-parametric ANOVA: Kruskal-Wallis test.

Here the null hypothesis is that the means come from the same population.

Example: Let us take the same data as used in the parametric test. Firstly, we arrange the data in ascending order within samples.

- Samples stn. 1) 98, 22, 72, 214, 67
- stn. 2) 12, 13, 46, 38, 49
- stn. 3) 86, 12, 49, 33, 72
- stn. 4) 2, 5, 12, 3, 19

Now rank all the above in ascending order:

		Total	$n_i$	$(R_i^2/n_i)$	
stn. 1)	19, 9, 16.5, 20, 15	$R_1$	79.5	5	1264.05
stn. 2)	5, 7, 12, 11, 13.5	$R_2$	48.5	5	470.45
stn. 3)	18, 5, 13.5, 10, 16.5	$R_3$	63	5	793.80
stn. 4)	1, 3, 5, 2, 8	$R_4$	19	5	72.20

---

Total    R= 210    N= 20    2600.50

Calculate K statistic

$$\begin{aligned}
 K &= (12 / (N(N + 1))) (\sum(R_i)^2 / n_i) - 3(N + 1) \\
 &= (12 / 20(21)) (2600.5) - 3(21) \\
 &= 11.3
 \end{aligned}$$

Refer to Tables of  $\chi^2$  for  $v = i - 1$  d.f. = 3  
 $p0.05_{(3)} = 7.81$ ;  $p0.01_{(3)} = 11.2$

Conclusion: Since the calculated value is greater than the table value at  $p0.01$  we reject the hypothesis (with 99% certainty) that the means are from the same population.

Non-parametric Two-Way ANOVA: The Friedman test.

This test is used in COMBINATION with the Kruskal-Wallis test under defined conditions. The number of counts in each sample must be the same and each count must belong to one sample and one group, where the group can represent different bottom types, different samplers or different workers. In the previous example let us assume that the four samples were each counted by five different workers and we are therefore, interested to know if there is also a significant difference between workers. The  $H_0$  is that there is no significant difference between workers.

Example: 4 sample  $i = 4$ , 5 workers  $n = 5$

	Worker				
	1	2	3	4	5
Sample 1	98	22	72	214	67
Sample 2	12	13	46	38	49
Sample 3	86	12	49	33	72
Sample 4	2	5	12	3	19

Firstly arrange samples by rank in each row:

Sample 1	4	1	3	5	2
Sample 2	1	2	4	3	5
Sample 3	5	1	3	2	4
Sample 4	1	3	4	2	5

---

Total	11	7	14	12	16
	$R_1$	$R_2$	$R_3$	$R_4$	$R_5$

$$\Sigma R_n = 60 \quad \Sigma R_n^2 = 766$$

$$S = \Sigma R_n^2 - ((\Sigma R_n)^2 / n)$$

$$= 766 - (60^2) / 5 = 46$$

$$n = 5, i = 4$$

So we must calculate a  $\chi^2$  value where:

$$\chi^2 = 12S / (in(n + 1))$$

$$= 12(46) / 4 * 5(6)$$

$$= 4.6$$

$$\chi^2 \text{ for } v = n-1 \text{ d.f.} = 4 \text{ for } p0.05_{(4)} = 9.49$$

Conclusion: As the calculated value is less than that of the Table value we cannot reject the null hypothesis that there was no significant difference between workers.

Non-parametric Correlation tests.

1. Spearman's rank correlation.

Here we first simply rank the two sets of data.

Example:	Sample 1	Sample 2	(Difference) <sup>2</sup>
	4	4	0
	1	2	1
	6	5	1
	5	6	1
	3	1	4
	2	3	1
	7	7	0
		Total	8

$$\begin{aligned}
 \text{Calculate } r_s &= 1 - (6\sum d^2) / (n(n^2 - 1)) \\
 &= 1 - 6(8) / 7(49-1) \\
 &= 0.857
 \end{aligned}$$

This value is tested against the Table value for the Correlation Coefficient (r). d.f. = n - 1 = 6 p0.05<sub>(6)</sub> = 0.707.

Conclusion: As the calculated value is greater than the Table value we reject the H<sub>0</sub> that there is no correlation between the two data sets.

This test is calculated to be 90% as efficient as the parametric test.

2. Kendall's tau.

Here the two ranks are set alongside each other:

Sample 1	Sample 2	No. of ranks < pivotal rank
1	2	1
2	3	1
3	1	0
4	4	0
5	6	1
6	5	0
7	7	0
Total (Q)		3

$$\begin{aligned}\text{Tau} &= 1 - (4(Q)) / (n(n - 1)) \\ &= 1 - (4*3 / 42) \\ &= 0.714\end{aligned}$$

We look up the Table of the Correlation Coefficient (r) for d.f. 6 and  $p_{0.05} = 0.707$

Conclusion: Since the calculated value is greater than the Table value we reject  $H_0$  that there is no correlation between the two data sets.

#### References:

**General texts.** One of the following three is required:

Snedecor, G.W. and W.G. Cochran (1980), Statistical Methods. 7th ed. Iowa State Univ. Press, Ames, Iowa 507 pp.

Sokal, R.R. and F.J. Rohlf (1981), BIOMETRY, 2nd ed. W.H. Freeman & Co., San Francisco, 859 pp.

Zar, J.R. (1984), Biostatistical Analysis, Prentice-Hall.

**Statistical tables:** One set is needed

Snedecor & Cochran and Zar include tables in their books. Rohlf and Sokal has separate (and expensive) tables.

Rohlf, F.J. and R.R. Sokal (1984), Statistical Tables. W.H. Freeman

Neave, H.R. (1978), Statistical Tables Allen & Unwin. (Cheap!)

**Excellent book on analysis of field samples:**

Elliott, J.M. (1971), Statistical Analysis of samples of Benthic Invertebrates. Fresh Water Biological Association, Windermere, U.K.

**PART II**

## LECTURE 1

### A FRAMEWORK FOR STUDYING CHANGES IN COMMUNITY STRUCTURE

#### STAGES

- 1) REPRESENTING COMMUNITIES (graphical description of faunal relations).
- 2) DISCRIMINATING SITES on the basis of faunal composition (e.g. spatial: control v. impacted, temporal: before v. after impact).
- 3) DETERMINING LEVELS OF "STRESS" or disturbance in communities.
- 4) LINKING WITH ENVIRONMENTAL VARIABLES (e.g. correlating to contaminants)
- 5) ESTABLISHING CAUSALITY of link to contaminants.

#### TECHNIQUES

UNIVARIATE - diversity indices

- indicator species abundance

DISTRIBUTIONAL - "ABC" curves (k-dominance)

- distn. of individuals amongst species

MULTIVARIATE - triangular matrix of similarities between samples, leading to:

- hierarchical classification (CLUSTER)
- multidimensional scaling (MDS)
- principal component analysis (PCA)

#### UNIVARIATE TECHNIQUES

#### EXAMPLES

#### STAGES

	Diversity indices	Indicator species
1) REPRESENTING COMMUNITIES	Means $\pm$ confidence intervals (CIs for each site)	
2) DISCRIMINATING SITES	One-way analysis of variance (ANOVA)	
3) DETERMINING STRESS LEVELS	By reference to historical data, e.g. ultimately a decrease in diversity	initial increase in "opportunistic" species
4) LINKING TO ENVIRONMENT	Regression techniques	

5) ESTABLISHING CAUSALITY

Mesocosm or field experiments with controlled dosing of contaminants. All entries above apply, e.g. now significant discrimination of "sites" (=treatments) demonstrates that contaminant causes biological effect.

IOC/GEEP WORKSHOP ON BIOLOGICAL EFFECTS OF POLLUTANTS

OSLO 1986: MACROFAUNAL DATA, Gray et al. (1988)




Fig. 1.1 Frierfjord and Langesundfjord, Norway. Benthic community sampling sites (A-G) for the Oslo Workshop



Four 0.1 m<sup>2</sup> Day grab samples taken at 6 sites (A-E, G), sieved at 1 mm, and counts/biomass recorded of 110 species identified.

Table 1.1

Macrofaunal abundance matrix (part), numbers per 0.1 m<sup>2</sup>.

Species	A				B			
<i>Cerianthus lloydi</i>	0	0	0	0	0	0	0	0
<i>Halicryptus sp.</i>	0	0	0	1	0	0	0	0
<i>Onchnesoma</i>	0	0	0	0	0	0	0	0
<i>Phascolion strombi</i>	0	0	0	1	0	0	1	0
<i>Golfingia sp.</i>	0	0	0	0	0	0	0	0
Holothuroidea	0	0	0	0	0	0	0	0
Nemertina, indet.	12	6	8	6	40	6	19	7
Polychaeta, indet.	5	0	0	0	0	0	1	0
<i>Amaena trilobata</i>	1	1	1	0	0	0	0	0
<i>Amphicteis gunneri</i>	0	0	0	0	4	0	0	0
Ampharetidae	0	0	0	0	1	0	0	0
<i>Anaitides groenlandica</i>	0	0	0	0	1	0	0	0
<i>Anaitides sp.</i>	0	0	0	0	0	0	0	0

Table 1.2

Macrofaunal biomass matrix (part), mg per 0.1 m<sup>2</sup>.

Species	A				B			
<i>Cerianthus lloydi</i> /10	0	0	0	0	0	0	0	0
<i>Halicryptus sp.</i>	0	0	0	26	0	0	0	0
<i>Onchnesoma</i>	0	0	0	0	0	0	0	0
<i>Phascolion strombi</i>	0	0	0	6	0	0	2	0
<i>Golfingia sp.</i>	0	0	0	0	0	0	0	0
Holothuroidea	0	0	0	0	0	0	0	0
Nemertina, indet./10	1	41	391	1	5	1	2	1
Polychaeta, indet.	9	0	0	0	0	0	0	0
<i>Amaena trilobata</i>	144	14	234	0	0	0	0	0
<i>Amphicteis gunneri</i>	0	0	0	0	45	0	0	0
Ampharetidae	0	0	0	0	0	0	0	0
<i>Anaitides groenlandica</i>	0	0	0	7	11	0	0	0
<i>Anaitides sp.</i>	0	0	0	0	0	0	0	0

UNIVARIATE: REPRESENTATION AND DISCRIMINATION




Fig. 1.2 Frierfjord macrofauna. Means and 95% confidence intervals for two indices. a) Number of species (S); b) Shannon diversity (H')

DISTRIBUTIONAL TECHNIQUES

EXAMPLES

"ABC" curves (k-dominance curves)	Distribution of individuals amongst species
---	---

STAGES

- |    |                              |  |
|----|------------------------------|--|
| 1) | REPRESENTING<br>COMMUNITIES  | Curves for each site<br>(or preferably replicate)  |
| 2) | DISCRIMINATING<br>SITES      | ANOSIM (Analysis of Similarities)<br>test on "distances" between every<br>pair of curves   |
| 3) | DETERMINING<br>STRESS LEVELS | Biomass curve   Species abundance<br>drops below   distribution is<br>numbers curve   less "smooth"<br>when subject   with disturbance<br>to disturbance |
| 4) | LINKING TO<br>ENVIRONMENT    | Possible for univariate summary<br>statistics by regression  |
| 5) | ESTABLISHING<br>CAUSALITY    | Mesocosm or field dosing<br>experiments. Entries above apply   |

ORGANIC ENRICHMENT OF BENTHOS - Pearson (1975)

LOCH LINNHE (SCOTLAND) MACROFAUNA - discharges started in 1966,  
increased 1970, decreased 1972.



DISTRIBUTIONAL: REPRESENTATION AND STRESS DETERMINATION




Fig. 1.4 Loch Linnhe site 34. (A) Shannon diversity. (B)-(L) ABC curves for 1963-73: biomass (x), numbers ( $\square$ ). Warwick (1986)

DISTRIBUTIONAL: REPRESENTATION AND STRESS DETERMINATION




Fig. 1.5 Frierfjord macrofauna, sites A-E,G. Number of species against number of individuals per species in geometric classes (I = 1 individual per species, II = 2-3 ind. per spp., III = 4-7, IV = 8-15 etc.). Gray *et al.* (1988)

MULTIVARIATE TECHNIQUES

EXAMPLES

Hierarchical clustering		MDS ordination		PCA ordination
-------------------------	--	----------------	--	----------------

STAGES

REPRESENTING COMMUNITIES

Dendrogram of replicates

Configuration of replicates (often 2-D)

DISCRIMINATING SITES

ANOSIM test on triangular matrix of similarities  
Similarity percentage breakdown (SIMPER) gives species responsible

Multinormal tests (e.g. Wilks'  $\Lambda$ ), but often invalid

DETERMINING STRESS LEVELS

Not appropriate

LINKING TO ENVIRONMENT

Visual (superimposing environmental variables on faunal ordinations). Finding subset of environmental variables whose ordination "best" matches the faunal ordination.

ESTABLISHING CAUSALITY

Mesocosm or field dosing experiments. Use above techniques - significance in discriminating "sites" (=treatments) establishes causality.

MULTIVARIATE: REPRESENTATION

Table 1.4

Frierfjord macrofauna counts. Similarities (Bray-Curtis coefficient, after  $\sqrt{v}$  transformation) between every pair of replicates (sites A-C only).

	A1	A2	A3	A4	B1	B2	B3	B4	C1	C2	C3	C4
A1	-											
A2	61	-										
A3	69	60	-									
A4	65	61	66	-								
B1	37	28	37	35	-							
B2	42	34	31	32	55	-						
B3	45	39	39	44	66	66	-					
B4	37	29	29	37	59	63	60	-				
C1	35	31	27	25	28	56	40	34	-			
C2	40	34	26	29	48	69	62	56	56	-		
C3	40	31	37	39	59	61	67	53	40	66	-	
C4	36	28	34	37	65	55	69	55	38	64	74	-




Fig. 1.6 Frierfjord macrofauna. Dendrogram for hierarchical clustering (group-average link) of 4 replicates from 6 sites, using above similarities



MULTIVARIATE: REPRESENTATION AND DISCRIMINATION




Fig. 1.7 Frierfjord macrofauna. Non-metric MDS ordination (in 2-D) of the 4 replicates from each of sites A-E and G, from Table 1.4 similarities




Fig. 1.8 Loch Linnhe macrofauna. PCA ordination (in 2-D) of the 11 years abundance data, omitting the less-common species

MULTIVARIATE: LINKING TO ENVIRONMENT




Fig. 1.9 Frierfjord macrofauna. Values of four environmental variables: (a) water depth, (b) sediment grain size, (c) metal and (d) PAH concentrations in sediment, superimposed on the abundance-based MDS

NUTRIENT ENRICHMENT MESOCOSM EXPERIMENT

- Gee et al. (1985)

Meiofaunal abundances under 2 dosing regimes, Solbergstrand facility (NIVA), Norway

Table 1.5

Copepod numbers (nematodes not shown) from 4 boxes for each treatment (high, low and no additions of powdered *Ascophyllum nodosum*).

	Control				Low dose				High dose			
	C1	C2	C3	C4	L1	L2	L3	L4	H1	H2	H3	H4
Copepoda, Harpacticoida												
Ectinosomidae												
<i>Halectinosoma gothiceps</i>	-	-	1	1	16	23	8	16	-	1	-	-
Tachidiidae												
<i>Danielssania fusiformis</i>	1	1	1	1	1	3	8	5	1	-	-	3
Tisbidae												
<i>Tisbe</i> sp. 1 (gracilis group)	-	-	-	-	-	-	-	-	2	27	119	31
<i>Tisbe</i> sp. 2 (graciloides?)	-	-	-	-	45	22	39	25	6	-	3	32
<i>Tisbe</i> sp. 3	-	-	-	-	86	83	88	-	5	29	-	20
<i>Tisbe</i> sp. 4	-	-	-	-	151	249	264	87	8	-	-	34
<i>Tisbe</i> sp. 5	-	-	-	-	129	-	-	115	4	-	1	40
Diosaccidae												
<i>Typhlamphiascus typhlops</i>	4	2	2	4	5	8	4	3	-	-	-	-
<i>Bulbamphiascus imus</i>	1	-	-	2	-	-	-	-	-	-	-	-
<i>Stenhelia reflexa</i>	3	1	-	1	2	-	-	-	-	-	-	-
<i>Amphiascus tenuiremis</i>	1	-	-	-	-	-	2	6	-	-	-	-
Ameiridae												
<i>Ameira parvula</i>	-	-	-	-	4	2	3	2	2	-	1	2
<i>Proameira simplex</i>	-	-	-	-	-	2	-	5	-	-	-	-
Paramesochridae												
<i>Leptosyllus paratypicus</i>	-	-	1	-	-	-	-	-	-	-	-	-
Cletodidae												
<i>Enhydrosoma longifurcatum</i>	2	2	1	2	3	1	-	-	-	-	-	-
Laophontidae												
Unidentified copepodite	-	-	-	-	-	-	1	-	-	-	-	-
Ancorabolidae												
<i>Ancorabolis mirabilis</i>	3	-	4	4	2	18	3	3	27	3	1	-
Unidentified Copepodites	-	-	1	-	1	1	1	3	-	-	-	-

MULTIVARIATE: ESTABLISHING CAUSALITY




Fig. 1.10 Mesocosm meiofauna (nutrient enrichment). MDS ordination of abundances from 4 replicate boxes from 3 treatments: circles = control, squares = low dose, triangles = high dose. (Gee et al., 1985)

DATA TRANSFORMATION AND SPECIES SELECTION/AGGREGATION

Some techniques may need TRANSFORMATION of the raw abundances/biomass (or derived statistics) for:

- a) validity of assumptions for statistical analysis (e.g. normality, constant variance);
- b) balancing contributions of rare/abundant species.

Some techniques may be possible with data on SELECTED (more dominant) species or data AGGREGATED to higher taxonomic levels, thus minimising identification time.

<u>TECHNIQUE</u>	<u>EXAMPLES</u>	<u>TRANSFORMATION</u>	<u>SELECTION/ AGGREGATION</u>
UNIVARIATE	Diversity indices	Counts: No Index: Possibly	No
	Indicator species	Yes (on counts/biomass)	Yes
DISTRIBUTIONAL	ABC curves	Possible but not usual	Possible
	Ind. among species	No	No
MULTIVARIATE	Cluster	Usual (log or 4th root) on counts/biomass	Possible
	MDS	MDS transforms similarities	Possible
	PCA	also, to ranks.	Needed

## LECTURE 2

### MULTIVARIATE METHODS: MEASURES OF SIMILARITY OF SPECIES ABUNDANCE/BIOMASS BETWEEN SAMPLES

DATA MATRIX: A  $p$  (species)  $\times$   $n$  (samples) array of scores (counts or biomass). The  $n$  samples might consist of a number of replicates (possibly only one) at each of a number of sites or times.

SIMILARITY COEFFICIENT: Measures the similarity ( $S$ ) of the community structure between any pair of samples (thus SAMPLE SIMILARITIES), using:

- a) absolute numbers (or biomass) of each species,
- b) relative numbers (or biomass), i.e. STANDARDISE the scores, to reflect only species COMPOSITION (%),
- c) only presence or absence of each species.

$S$  is usually defined in the range (0, 1) or (0, 100%).

$S = 1$  (or 100%) means samples are totally similar,  
 $S = 0$  means samples are totally dissimilar.

SIMILARITY MATRIX: This is a set of similarity coefficients, calculated between every pair of samples and laid out in a lower triangular array.

Similarity matrices are the basis for many clustering and ordination techniques (REPRESENTATION) and associated tests (DISCRIMINATION), which:

- a) discriminate sites or times (similarities between replicates at a site  $>$  similarities between sites)
- b) cluster sites (similarities within groups of sites  $>$  similarities between groups)
- c) allow gradation of sites (site A has similarities with B, and B has with C, but A and C less similar).

SPECIES SIMILARITY MATRIX: A matching triangular array of similarities between every pair of species, in terms of patterns of occurrence across the samples.

Many different ways to assess similarity (because data is multi-species). One of the most useful in ecology is:

BRAY-CURTIS COEFFICIENT: (Bray and Curtis, 1957).  
Similarity between jth and kth samples is:

$$S_{jk} = 100 \left\{ 1 - \frac{\sum_{i=1}^P |y_{ij} - y_{ik}|}{\sum_{i=1}^P (y_{ij} + y_{ik})} \right\} \quad (2.1)$$

$$= 100 \frac{\sum_{i=1}^P 2 \min(y_{ij}, y_{ik})}{\sum_{i=1}^P (y_{ij} + y_{ik})}$$

where  $y_{ij}$  = score (count or biomass) for  $i$ th species in  $j$ th sample ( $i=1,2,\dots,p$ ;  $j = 1,2,\dots,n$ ).

Example: Loch Linnhe macrofauna (Pearson, 1975).

Table 2.1

(a) Abundance (untransformed) for some selected species and years from site 34 data. (b) Resulting Bray-Curtis similarity matrix.

(a) Year:	64	68	71	73	(b)				
(Sample:	1	2	3	4)	Sample	1	2	3	4
Species					1	-			
<i>Echinoca</i>	9	0	0	0	2	8	-		
<i>Myrioche</i>	19	0	0	3	3	0	42	-	
<i>Labidopl.</i>	9	37	0	10	4	39	21	4	-
<i>Amoeana</i>	0	12	144	9					
<i>Capitella</i>	0	128	344	2					
<i>Mytilus</i>	0	0	0	0					

- 1) Note  $S = 0$  if the two samples have no species in common (e.g. 1 and 3 above).
- 2) A scale change in  $y$  (e.g. biomass changed from mg per  $m^2$  to per  $cm^2$ ) does not change  $S$ .
- 3) "Joint absences" also have no effect on  $S$  (as is desirable), e.g. can omit species 6 in the table.

With "raw" counts (or biomass),  $S$  gives too much weight to large scores, so a  $\log(1+y)$  or  $\sqrt{y}$  transform is often applied, before computing  $S$ .

Example: Loch Linnhe macrofauna,  $\sqrt{}$  transformation

Table 2.2

(a)  $\sqrt{}$ - transformed abundances for 4 years.  
 (b) Resulting Bray-Curtis similarity matrix.

(a) Year: (Sample: Species	64 1	68 2	71 3	73 4)	(b) Sample	1	2	3	4
1	1.7	0	0	0	1	-			
2	2.1	0	0	1.3	2	26	-		
3	1.7	2.5	0	1.8	3	0	68	-	
4	0	1.9	3.5	1.7	4	52	68	42	-
5	0	3.4	4.3	1.2					
6	0	0	0	0					

CANBERRA COEFFICIENT: Lance and Williams, 1967.  
 Similarity between samples j and k is:

$$S_{jk} = 100 \left\{ 1 - p^{-1} \sum_{i=1}^p \frac{|y_{ij} - y_{ik}|}{(y_{ij} + y_{ik})} \right\} \quad (2.2)$$

It gives a more equal contribution from each species (so tends to be overdominated by rarer ones).

CORRELATION COEFFICIENT: Product-moment correlation

$$r_{jk} = \frac{\sum_i (y_{ij} - \bar{y}_{.j}) (y_{ik} - \bar{y}_{.k})}{\sqrt{[\sum_i (y_{ij} - \bar{y}_{.j})^2 \cdot \sum_i (y_{ik} - \bar{y}_{.k})^2]}} \quad (2.3)$$

is not a similarity (it can be <0). Use:

$$S_{jk} = 50 (1 + r_{jk}) \quad (2.4),$$

but note that S increases with more joint absences.



### PRESENCE/ABSENCE DATA

Many similarity coefficients have been proposed based on (0,1) data arrays (Sneath and Sokal, 1973). For comparing samples j and k let:

a = number of species present in both samples,  
b+c = number present in one sample and not the other,  
d = number absent from both samples.

"SIMPLE MATCHING" COEFFICIENT:

$$S_{jk} = 100.(a+d)/(a+b+c+d) \quad (2.5)$$

Note that this is a function of joint absences (d).

JACCARD'S COEFFICIENT:

$$S_{jk} = 100.a/(a+b+c) \quad (2.6)$$

SORENSEN (OR DICE) COEFFICIENT:

$$S_{jk} = 100.2a/(2a+b+c) \quad (2.7)$$

This is simply BRAY-CURTIS applied to (0,1) data.

McCONNAUGHEY COEFFICIENT (McConnaughey, 1964):

$$S_{jk} = 100[a(2a+b+c)]/[2(a+b)(a+b)] \quad (2.8)$$

### RECOMMENDATION:

- 1) Use coefficient not dependent on joint absences.
- 2) Similarities from raw counts (or biomass) are too dominated by common (or large) species, but
- 3) Reduction to presence/absence loses too much useful information, so recommend use:
- 4) BRAY-CURTIS on  $\sqrt{y}$  or  $\log(1+y)$  transformed data.
- 5) Standardise scores if non-comparable sample volumes used, or if "patchiness" makes compositional change more relevant than fluctuations in absolute counts.

SPECIES SIMILARITIES: These are computed from the same data array but between any pair of species (rows i,l say) across all samples (columns).

BRAY-CURTIS: 
$$S'_{il} = 100 \left\{ 1 - \frac{\sum_{j=1}^n |y_{ij} - y_{lj}|}{\sum_{j=1}^n (y_{ij} + y_{lj})} \right\} \quad (2.9)$$

However:

- 1) Similarities between rare species have little meaning ( $S'$  usually 0) and should be omitted from any species clustering or ordination.
- 2) Standardisation (not transformation) of  $y$  needed:

$$y_{ij}^* = 100 y_{ij} / (\sum_{k=1}^n y_{ik}) \quad (2.10),$$

(before computing  $S'$ ), so two species in strict ratio across samples are "perfectly similar".

<u>Example</u>	<u>Counts</u>					→	<u>Similarities</u>			
Sample:	1	2	3	4	5		Species	1	2	3
Species							1	-		
1	2	0	0	4	4	→	2	33	-	
2	10	0	0	20	20		3	20	7	-
3	0	4	4	1	1					
						↓				
						Standardise				
Species							1	-		
1	20	0	0	40	40	→	2	100	-	
2	20	0	0	40	40		3	20	20	-
3	0	40	40	10	10					

CORRELATION coefficients are more appropriate for species similarity, since they incorporate scale changes, but the location changes are undesirable.

**RECOMMENDATION:** For species similarities, use BRAY-CURTIS on standardised scores. Remove rarer species (never >3%, say, of total score in any sample).

### DISSIMILARITY COEFFICIENTS

These are important in constructing ordinations, in which dissimilarities ( $\delta$ ) between pairs of samples are turned into distances ( $d$ ) between sample locations on a "map". ( $\delta$  therefore >0, of course).

Similarities can easily become dissimilarities, by:

$$\delta = 100 - S \quad (2.11),$$

e.g. for BRAY-CURTIS:

$$\delta_{jk} = 100 \cdot \frac{\sum_{i=1}^p |y_{ij} - y_{ik}|}{\sum_{i=1}^p (y_{ij} + y_{ik})} \quad (2.12)$$

so  $\delta=0$ : no dissimilarity,  $\delta=100$ : total dissimilarity.


Other dissimilarity measures, based on distances:

EUCLIDEAN DISTANCE:

$$d_{jk} = \sqrt{[\sum_{i=1}^p (y_{ij} - y_{ik})^2]} \quad (2.13)$$

MANHATTEN (or CITY-BLOCK) DISTANCE:

$$d_{jk} = \sum_{i=1}^p |y_{ij} - y_{ik}| \quad (2.14)$$



[METRICS: Euclidean and Manhattan measures, (2.13) and (2.14), are called distances or metrics because they obey the triangle inequality, i.e. for any three samples, j,k,r:

$$d_{jk} + d_{kr} \geq d_{jr} \quad (2.15)$$

Note: Bray-Curtis dissimilarity does not satisfy the triangle inequality, so should not be called a "metric". However, many useful dissimilarities are also not metrics (e.g. squared Euclidean distance, giving dissimilarities of the same rank order as Euclidean distance, i.e. identical MDS ordinations).

CONCLUDE: Unnecessary to insist that dissimilarities are true "metrics".]

Where necessary (e.g. for input to clustering), distance ( $d$ ) can be conveniently converted to similarity ( $S$ ) by:

$$S = 100/(1 + d) \quad (2.16),$$

and, using (2.11), distance ( $d$ ) turned to dissimilarity ( $\delta$ ) by

$$\delta = 100d/(1 + d) \quad (2.17).$$

So,  $d = 0$  gives  $\delta = 0$ ,  $S = 100$ , and  $d \rightarrow \infty$  gives  $\delta \rightarrow 100$ ,  $S \rightarrow 0$ .

However, note that EUCLIDEAN (or MANHATTEN) distance is the same if a species is absent in both samples or is present in both at the same abundance; this is undesirable. (Same problem as that of similarities based on correlation being dependent on joint absences.) So:

**RECOMMENDATION:** For clustering or MDS of species counts/biomass, use Bray-Curtis dissimilarities, after suitable transformation, rather than Euclidean (or Manhattan) distances.




Fig. 2.1 Stages in a multivariate analysis based on (dis)similarity coefficients

LECTURE 3

MULTIVARIATE METHODS: HIERARCHICAL CLUSTERING

Table 3.1

Frierfjord macrofauna counts. Similarities (Bray-Curtis coefficient, after  $\sqrt{}$  transformation) between every pair of replicates (sites A-C only).

	A1	A2	A3	A4	B1	B2	B3	B4	C1	C2	C3	C4
A1	-											
A2	61	-										
A3	69	60	-									
A4	65	61	66	-								
B1	37	28	37	35	-							
B2	42	34	31	32	55	-						
B3	45	39	39	44	66	66	-					
B4	37	29	29	37	59	63	60	-				
C1	35	31	27	25	28	56	40	34	-			
C2	40	34	26	29	48	69	62	56	56	-		
C3	40	31	37	39	59	61	67	53	40	66	-	
C4	36	28	34	37	65	55	69	55	38	64	74	-

Seeing structure in a similarity matrix is difficult - a graphic representation of relations is needed:

CLUSTER ANALYSIS Clustering (or classification) aims to find "natural groupings" of samples such that samples within a group are more similar than samples in different groups. Use clustering to:

- 1) Distinguish sites (or times) - replicates within sites fall in the same cluster;
- 2) Partition sites (or times) into groups;
- 3) Define species assemblages (spp. co-occur at sites)

Hundreds of clustering methods exist (Everitt, 1980), some operating on (dis)similarities, some on raw data. Cormack (1971) warns against indiscriminate use: "availability of ... classification techniques has led to the waste of more valuable scientific time than any other 'statistical' innovation".

Five classes of clustering methods can be defined:

- 1) Hierarchical, 2) Optimising, 3) Mode seeking, 4) Clumping and 5) Miscellaneous techniques.

Here consider only one (sub)class, which recognises that clustering can occur at several levels.

**HIERARCHICAL AGGLOMERATIVE CLUSTERING:** The n samples are successively fused into groups, starting with samples with the highest mutual similarities then gradually lowering the similarity level at which groups are fused, and ending in a single cluster. (DIVISIVE clustering is the opposite sequence). Process represented by a tree diagram or DENDROGRAM.

DISTINGUISHING SITES: Frierfjord macrofauna counts.




Fig. 3.1 Frierfjord macrofauna counts. Dendrogram for hierarchical clustering (using group-average linking) of 4 replicates from each of sites A-E,G, using Bray-Curtis similarity matrix (Table 3.1)

GROUPING TIMES: Loch Linnhe macrofauna - subset. After  $\sqrt{}$  transformation, data array and bray-Curtis similarity matrix are:

Year:	64	68	71	73	Sample	1	2	3	4
Sample:	1	2	3	4	1	-			
Species					2	25.6	-		
<i>Echin.</i>	1.7	0	0	0	3	0.0	67.9	-	
<i>Myrio.</i>	2.1	0	0	1.3	4	52.2	<u>68.1</u>	42.0	
<i>Labid.</i>	1.7	2.5	0	1.8					
<i>Amaea.</i>	0	1.9	3.5	1.7					
<i>Capit.</i>	0	3.4	4.3	1.2					
<i>Mytil.</i>	0	0	0	0					

Samples 2 and 4 have the highest similarity,  $S(2,4)$ , so they form the first group.


Their similarity to (say) sample 1 defined in one of 3 ways:

Sample	1	2 & 4 fused	3
1	-		
2&4	38.9	-	
3	0.0	<u>55.0</u>	-

↓ (2&4) & 3 fused

Sample	1	2&3&4
1	-	
2&3&4	<u>25.9</u>	-

- a) SINGLE LINKAGE:  $\max\{S(1,2), S(1,4)\}$  (=52.2)
- b) COMPLETE LINKAGE:  $\min\{S(1,2), S(1,4)\}$  (=25.6)
- c) GROUP AVERAGE LINK:  $[S(1,2) + S(1,4)]/2$  (=38.9)  
 (Average weighted by number of samples in groups fused, e.g.  $S(1,2&3&4) = (2 \times 38.9 + 1 \times 0)/3 = 25.9$ ).




Note:

- 1) Samples need to be reordered for clear presentation of the dendrogram (so there are no crossing lines).
- 2) The order of samples on the x axis is not very meaningful (think of a dendrogram as a "mobile").

- 3) Here clustering imposes a (somewhat arbitrary) grouping on what is essentially a continuum (clean (1), impacted (2 and 3) and some recovery (4)), so:
- 4) Small changes in similarities can have larger effects on picture (e.g. reverse S(2,3) & S(2,4)).

DISSIMILARITIES: Exactly converse operations needed for a dissimilarity matrix, i.e. fuse samples with lowest dissimilarity, take minimum dissimilarity in single linkage, maximum in complete linkage.

LINKAGES: These three options are best visualised for an example with only 2 species and dissimilarity defined simply from Euclidean distance.



Group average is mean of all 12 intergroup distances.

Explains why alternative names for the linkages are:

- "NEAREST NEIGHBOUR" = single linkage
- "FURTHEST NEIGHBOUR" = complete linkage

Note: Though single linkage has some nice theoretical properties (e.g. clustering only a function of rank order of similarities), it has a tendency to give chains of linked samples rather than clear groups; group average linking is usually preferable.



Example: Bristol Channel (UK) zooplankton, April 1974, 57 sites X 24 species, Collins and Williams (1982).




Fig. 3.2 Bristol Channel sampling sites 1-29, 31-58




Fig. 3.3 Bristol channel. Dendrogram for hierarchical clustering of 57 sites (group average linking of Bray-Curtis similarities on  $\sqrt{\text{abundance}}$ )

## RECOMMENDATIONS

Hierarchical clustering (with group average linking) on sample (dis)similarity matrices can be useful, especially to delineate discrete communities at differing sites (or groups of sites).

It is less useful (and can be misleading) for a gradation in community structure across sites or times; ordination is preferable for this (see lectures 4 and 5).

Clustering is best used in conjunction with an ordination (even for discrete communities), for example, by superimposing clusters on the sample ordination plot.

## LECTURE 4

### MULTIVARIATE METHODS: ORDINATION OF SAMPLES BY PRINCIPAL COMPONENTS ANALYSIS (PCA)

**ORDINATIONS:** These are techniques for MAPPING the SAMPLES in a low number of dimensions (usually 2) such that the DISTANCE between samples attempts to reflect (DIS)SIMILARITY in community structure. (No guarantee that the attempt will succeed, if the relationships between the samples are complex, i.e. the structure is essentially "high-dimensional".)

Again there are many methods, for example:

PRINCIPAL COMPONENTS ANALYSIS (PCA, e.g. Chatfield & Collins, 1980),

PRINCIPAL CO-ORDINATES ANALYSIS (PCoA, Gower, 1966),

DETRENDED CORRESPONDENCE ANALYSIS (DECORANA, Hill & Gauch, 1980),

NON-METRIC MULTIDIMENSIONAL SCALING (MDS, e.g. Kruskal & Wish, 1978).


Here we consider only PCA (a simple but rather limited method) and MDS (a more complex algorithm but simple in concept and very generally applicable).

#### PRINCIPAL COMPONENTS ANALYSIS

STARTING POINT is the original DATA MATRIX (rather than a similarity matrix). The data array is thought of as defining the positions of samples in relation to axes representing the full set of species (one axis for each species). The samples are thus POINTS in a very HIGH-DIMENSIONAL SPACE, so it helps to visualise the process by considering an example in which there are only two species, i.e. each sample is a point in 2-dimensions.


#### Example:

	Sample:	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>	<u>8</u>	<u>9</u>
Abundance	Sp.1:	6	0	5	7	11	10	15	18	14
	Sp.2:	2	0	8	6	6	10	8	14	14




(This is an ORDINATION already - of 2-d data in 2-d, thus perfectly summarising all the relationships between samples).


For a 1-d ordination (i.e. a genuine ordering of samples) could take just one variable (Sp.1, say):



but this is poorer approximation to relations between samples than given by a (perpendicular) PROJECTION onto the line of "best fit" in the 2-d plot:



This is 1st PC AXIS; PC2 AXIS is PERPENDICULAR to this



PC AXES (full set) are simply a ROTATION of original species axes. Refer samples to (PC1, PC2) rather than (Sp.1, Sp.2) axes because may be able to DISPENSE WITH PC2, giving an ordination in 1-d.

Biggest differences between samples take place along PC1, and this is an equivalent definition of PC1 - the axis along which VARIANCE IS MAXIMISED.

Example: Add a third species to previous example.

	Sample:	1	2	3	4	5	6	7	8	9
Abundance	Sp.1:	6	0	5	7	11	10	15	18	14
	Sp.2:	2	0	8	6	6	10	8	14	14
	Sp.3:	3	1	6	6	9	11	10	16	15

Samples are now points in 3-d and there are 3 PC axes, again a rotation of the 3 species axes, such that:

PC1: Axis which MAXIMISES VARIANCE of points PROJECTED PERPENDICULARLY onto it.

PC2: Constrained to be perpendicular to PC1, again chosen to maximise variance along this axis.

PC3: Perpendicular to PC1 and PC2.

The new variables (PCs) are then just LINEAR COMBINATIONS of the old ones (species), such that PC1, PC2, PC3 are UNCORRELATED.

Here, the three PCs are:

$$\begin{aligned} \text{PC1} &= 0.62 \times \text{Sp.1} + 0.52 \times \text{Sp.2} + 0.58 \times \text{Sp.3} \\ \text{PC2} &= -0.73 \times \text{Sp.1} + 0.65 \times \text{Sp.2} + 0.20 \times \text{Sp.3} \\ \text{PC3} &= 0.28 \times \text{Sp.1} + 0.55 \times \text{Sp.2} - 0.79 \times \text{Sp.3} \end{aligned} \quad (4.1)$$

Letting  $\text{var}(\text{PC}_i)$  = variance of samples on  $i$ th PC axis,  
 $\text{var}(\text{Sp.}_i)$  = variance on  $i$ th species axis ( $i=1,2,3$ ):

$$\sum_i \text{var}(\text{PC}_i) = \sum_i \text{var}(\text{Sp.}_i) \quad (4.2)$$

so % OF (original) VARIANCE EXPLAINED by  $i$ th PC is:

$$\text{var}(\text{PC}_i) / \sum_i \text{var}(\text{PC}_i) \quad (4.3).$$

Here PC1 explains 93%, PC2 6% and PC3 1% of variance. Little variability (information) in PC3. Ignore it, so

PCA ORDINATION: The PC1 and PC2 axes give a 2-d ordination plane (of "best fit" to the sample points) and points are projected perpendicularly onto this from the higher PCs (just PC3 here). In this case, the 2-d ordination is almost a perfect summary of the 3-d data (the sample points lie near to a plane in the original 3-d species space).

**HIGHER-DIMENSIONAL DATA:** Typically, there are many more species (say 30+) but the approach is identical. Samples are points in the 30-d (say) species space; the "best-fit" 2-d plane is found and samples projected onto it to get the 2-d PCA ordination. Success is measured by the % of the variability explained by the first 2 of the 30 PCs.

**COMPUTATION:** Construction of PCs requires derivation of eigenvalues and vectors of a  $p \times p$  matrix ( $p$  = no. of species), e.g. Chatfield and Collins, 1980 (note: knowledge of matrix algebra essential). Problems if  $p$  is large (compared with no. of samples), so:

**EXCLUDE LESS-COMMON SPECIES:** These distort ordination badly (even if the matrix operations are possible). E.g. for Loch Linnhe data, the PCA ordination (Fig. 4.1) excludes species making up <3% of total counts at any site, leaving 29 species from 115.

**TRANSFORM REMAINING ABUNDANCES (/BIOMASS)** before applying PCA, to avoid over-dominance by the very common species. E.g. in Loch Linnhe data, *Capitella* counts go over 4000; Fig. 4.1 uses  $\sqrt{\quad}$  transform.

Example: Loch Linnhe macrofauna (site 34, 1963-1973).




Fig. 4.1 Loch Linnhe abundances. 2-d PCA ordination of samples from 11 years; PC1 (x axis) and PC2 (y axis) account for 57% of total sample variability

SCALE AND LOCATION CHANGES: Data often NORMALISED (after any transform). For each species subtract the mean (across sites) and divide by the standard deviation. Equivalently, extract eigenvalues of the correlation rather than the covariance matrix, i.e. CORRELATION-BASED PCA rather than COVARIANCE-BASED PCA. Essential if variables have different scales (units) or widely differing ranges. Not the case here (after transform at least) so less necessary (but was done in Fig. 4.1).

### PCA STRENGTHS

- 1) CONCEPTUALLY SIMPLE.
- 2) COMPUTATIONALLY STRAIGHTFORWARD, provided the number of species is reduced (usually drastically), and it can then cope with an unlimited number of samples.
- 3) ORDINATION AXES potentially have some meaning, as simple LINEAR COMBINATIONS of the species (though these are rarely readily interpretable in practice).

### PCA WEAKNESSES

- 1) LITTLE FLEXIBILITY in defining relations between samples - in effect "dissimilarities" are simple Euclidean distances in the species space. The only flexibility comes from transformation of the species axes.
- 2) Does NOT do a very good job of PRESERVING these DISTANCES (dissimilarities) in the 2-d ordination - samples that are far apart in the full space can end up coincident on the 2-d "best fit" plane, e.g. projected onto it "from opposite sides".

Example: Nematodes from Solbergstrand mesocosm experiment, GEEP Workshop (Warwick et al., 1988).




Fig. 4.2 Mesocosm nematodes. Correlation-based PCA of 16 samples: 4 replicate boxes from each of 4 treatments. (C=control, L=low, M=medium and H=high levels of diesel oil and Cu, water dosed for 11 weeks). 26 species retained (usual >3% dominance criterion) - log(1+count) transform applied. PC1 accounts for 23% of variability, PC2 15%

Strong suggestion of H replicates separating out but note low % of variability explained, so ORDINATION UNRELIABLE. (MDS gives more realistic picture - see Fig. 5.5).



## LECTURE 5

### MULTIVARIATE METHODS: ORDINATION OF SAMPLES BY MULTI-DIMENSIONAL SCALING (MDS)

#### OTHER ORDINATION METHODS

PRINCIPAL CO-ORDINATES ANALYSIS (PCoA; Gower, 1966; Everitt, 1978): Also referred to as "CLASSICAL SCALING". Overcomes inflexibility of PCA by allowing WIDER RANGE of DISSIMILARITY definitions; essentially converts these to distance and does a PCA (so still subject to same PCA weakness of poor distance preservation). PCA thus a special case of PCoA, with dissimilarity = Euclidean distance.

DETRENDED CORRESPONDENCE ANALYSIS (DECORANA; Hill and Gauch, 1980): Relaxes another constraint of PCA, that of linear combinations of species. Allows CURVILINEAR COMPONENT AXES and can have effect of straightening out "horseshoe" ordinations. But:

MDS offers arguably the GREATEST FLEXIBILITY, in the sense of (lack of) assumptions made about the data.

#### NON-METRIC MULTIDIMENSIONAL SCALING (MDS, e.g. Kruskal and Wish, 1978)

STARTING POINT is the (DIS)SIMILARITY MATRIX between samples (i.e. the relevant sample relationships). In fact, the ordination depends only on the RANKS of similarities in the triangular matrix, so is conceptually simple:


MDS attempts to construct a SAMPLE "MAP" (in a given number of dimensions, e.g. 2-d) using information of the form "Sample 1 is closer to Sample 4 (in species composition) than it is to Samples 2 or 3".

Example: Loch Linnhe macrofauna - subset (√√ counts)

Year:	64	68	71	73	
Sample:	1	2	3	4	
Species					Similarities
<i>Echin.</i>	1.7	0	0	0	
<i>Myrio.</i>	2.1	0	0	1.3	
<i>Labid.</i>	1.7	2.5	0	1.8	
<i>Amaea.</i>	0	1.9	3.5	1.7	
<i>Capit.</i>	0	3.4	4.3	1.2	
<i>Mytil.</i>	0	0	0	0	
Sample	1	2	3	4	
1	-				
2	25.6	-			
3	0.0	67.9	-		
4	52.2	68.1	42.0	-	

Rank dissimilarities

	Sample	1	2	3	4
	1	-			
	2	5	-		
	3	6	2	-	
	4	3	1	1	-



MDS  
plot

NOTE:

- 1) MDS plot can be arbitrarily SCALED, LOCATED, ROTATED or INVERTED; it gives positions of samples relative to each other.
- 2) Not difficult here to place 4 points in 2-d with interpoint distances preserving the rank order dissimilarities exactly. Usually not possible and there will be some distortion or STRESS between (ranked) dissimilarities and corresponding distances in the plot (even in a higher-dimensional ordination).

Example: R. Exe nematodes (Field et al., 1982)




Fig. 5.1 Exe nematodes. 2-d MDS ordination of 19 sites, from bray-Curtis similarities on  $\sqrt{\sqrt{}}$  transformed abundances (182 species)

MDS ALGORITHM - an iterative process

- 1) SPECIFY NUMBER OF DIMENSIONS for MDS plot (= m).
- 2) CONSTRUCT STARTING "MAP" of n samples; this could be result of (say) a PCA ordination or simply a random set of points (in m-dimensions).
- 3) REGRESS INTERPOINT DISTANCES  $\{d_{jk}\}$  from this map on the corresponding dissimilarities  $\{\delta_{jk}\}$ . Can be
  - a) LINEAR (or CURVILINEAR) regression - METRIC MDS; or, more usually
  - b) MONOTONIC (increasing) regression - NON-METRIC MDS (Fig. 5.2).




Fig. 5.2 Exe nematodes. "Shepard diagram" of distance ( $d$ ) in MDS plot (Fig. 5.1) against dissimilarity ( $\delta$ ) in Bray-Curtis matrix.  
o = actual distance ( $d_{jk}$ ),  
(\* =  $\geq 2$  coincident points),  
 $\Delta$  = fitted monotonic regression ( $\hat{d}_{jk}$ ).  
Stress (= 0.053) is a measure of scatter about the regression line

- 4) MEASURE GOODNESS-OF-FIT of the regression by:

$$\text{STRESS} = \frac{\sum_j \sum_k (d_{jk} - \hat{d}_{jk})^2}{\sum_j \sum_k d_{jk}^2} \quad (5.1)$$

where  $\hat{d}_{jk}$  = distance given by the fitted regression line for dissimilarity  $\delta_{jk}$ .

Stress = 0 if the distances preserve the rank order of the dissimilarities  $\{\delta\}$ .

Stress is large if the current map is poorly related to the dissimilarities  $\{\delta\}$ .

- 5) PERTURB CURRENT SAMPLE POSITIONS on the map, in direction decreasing the stress, using a STEEPEST DESCENT algorithm.
- 6) REPEAT STEPS 3 TO 5 (regress  $d$  on  $\delta$ , measure stress, perturb points) until no further reduction in stress is possible.

NOTE:

- a) The algorithm is an ITERATIVE PROCEDURE so could converge to a LOCAL MINIMUM rather than a global minimum of the stress function.

Also possible to get DEGENERATE SOLUTIONS where most samples collapse to the same point, or to the vertices of a triangle, or are strung out round a circle.

REPEAT FOR DIFFERENT RANDOM STARTING CONFIGURATIONS to confirm that gives same solution (with lowest stress value) several times - this is then very likely the GLOBAL MINIMUM (though not guaranteed).

- b) Unlike PCA, a 2-d MDS plot is NOT A PROJECTION of the 3-d plot. Still useful to do the 3-d MDS and use first 2 axes as the start for 2-d MDS - also useful to compare 2-d and 3-d stress values.

ADEQUACY OF MDS REPRESENTATION

- 1) STRESS VALUE: This increases with increasing number of samples and decreasing dimension of the plot, but roughly speaking, in 2-d:

STRESS < 0.05 implies excellent representation,  
< 0.1 good,  
< 0.2 still useful, but  
> 0.3 little better than random points.

(An alternative formula with a different denominator, "STRESS2", is preferred by some, but it increases the likelihood of finding local minima and is not recommended for routine use).

- 2) SHEPARD DIAGRAM: Scatter in this is measured by the stress value (low in Fig. 5.2, stress = 0.053, implying good MDS representation). Diagram also aids detection of "OUTLYING" POINTS and ERRORS in individual dissimilarities.
- 3) CONNECTION OF SIMILAR SAMPLES: Distortion in an MDS plot seen by connecting points whose similarities are in the top 10% or 20% (say) of values in the similarity matrix.
- 4) MINIMUM SPANNING TREE (MST): A similar idea - all points in the MDS plot are joined by a SINGLE CONNECTED LINE (which branches but is not allowed to form a closed loop) such that the sum of dissimilarities along this line is minimised; distortion is indicated by connections which look out of keeping with the distances in the plot (see Gower and Ross, 1969, for MST algorithm).
- 5) SUPERIMPOSITION OF GROUPS FROM CLUSTER ANALYSIS: The combination of clustering and ordination can be very effective.

Example: Exe nematodes, 19 sites (182 species)




Fig. 5.3 Exe nematodes. Dendrogram (group average linking, Bray-Curtis similarities on  $\sqrt{v}$  - abundance). 4 groups of sites separated by 15% similarity cut-off; 8 groups by a 30% (to 45%) threshold




Fig. 5.4 Exe nematodes. MDS (as Fig. 5.1) with clusters indicated at: --- 15%, — 30% similarity

Agreement clearly excellent (because clusters are sharp and MDS stress low). More revealing example provided by the data of Fig. 4.2:

Example: Mesocosm nematodes, GEEP Workshop.




Fig. 5.5 Mesocosm: 4 replicates from 4 treatments (reduced species and log transform, as Fig. 4.2).  
a), c) Group-average clustering from Bray-Curtis similarities; clusters formed at 3 (arbitrary) levels superimposed on the MDS obtained from the same similarities (stress = 0.19).  
b), d) Group average clustering from "Euclidean distance" (dis)similarities superimposed on the PCA (Fig. 4.2). (Euclidean distance is the dissimilarity measure implicit in a PCA ordination)



NOTE:

- 1) Though no natural groupings are apparent from the MDS, the Bray-Curtis cluster and MDS analyses (a and c) are not really inconsistent.
- 2) The PCA and its corresponding cluster analysis (d and b) are in disagreement, indicating that the 2-d PC axis is a distorted representation of the true "distances" between samples.

ORDINATION v CLUSTERING: Strength of ordination is in displaying GRADATION (rather than categorisation) of community composition in a set of samples.

Example: Celtic Sea zooplankton (Collins, pers. comm.)




Fig. 5.6 MDS of zooplankton samples at a single site (22/9/78), from 14 depths (5 m to 70 m, denoted A,B,...,N) for night (circles) and day-time hauls

**MDS STRENGTHS:**

- 1) SIMPLE in concept.
- 2) BASED ON RELEVANT INFORMATION. It can be used with the most appropriate measure of (dis)similarity for the particular data.
- 3) SPECIES DELETIONS UNNECESSARY for an ordination of samples (any exclusion dividing line is inevitably arbitrary). The similarity measure can automatically weight rarer species appropriately (and can be chosen to ignore joint absences).
- 4) GENERALLY APPLICABLE. Since MDS uses only rank order of dissimilarities it makes the weakest possible assumptions about quality of the data.
- 5) SIMILARITIES CAN BE GIVEN UNEQUAL WEIGHT in constructing the MDS plot (e.g. some samples may be more reliable, perhaps because they are based on combining more replicates).

**MDS WEAKNESSES:**

- 1) COMPUTATIONALLY DEMANDING; much more than  $n = 100$  samples is prohibitive (fewer on a PC; CPU time is proportional to  $n^2$ ).
- 2) CONVERGENCE to the correct solution (the global minimum of stress) is NOT GUARANTEED, since MDS is an iterative procedure; the necessary repeats add to the computational burden.
- 3) ALGORITHM PLACES MOST WEIGHT ON LARGE DISTANCES. For detailed structure within large clusters it is sometimes necessary to ordinate clusters separately (same constraint applies to most methods, eg. PCA).

**RECOMMENDATIONS:**

- 1) MDS RECOMMENDED as one of the best (perhaps the best) ordination technique (e.g. Everitt, 1978; Kenkel and Orłoci, 1986). Preferable to PCA because of its flexibility and (lack of) assumptions.
- 2) When sample relationships are simple (e.g. a few strong clusters; one strong gradient) most ordination methods will perform adequately. MDS scores because of its greater ability to REPRESENT MORE COMPLEX RELATIONS in 2-d space.
- 3) If stress is low (say,  $<0.1$ ), an MDS ordination is probably a more useful representation than a cluster analysis, even when the samples are strongly grouped. However, the techniques complement each other, so PERFORM BOTH, AND VIEW THEM IN COMBINATION, especially for higher stress. (In the latter case also try a higher-dimensional ordination).

LECTURE 6

MULTIVARIATE METHODS: TESTING FOR DIFFERENCES BETWEEN GROUPS OF SAMPLES

DISTINGUISHING SITES (or TIMES) by formal significance tests is a necessary first step to INTERPRETING differences (e.g. control v. impacted site) but usually overlooked for multivariate methods (because of unavailability of suitable tests).

(Note: Cluster analysis will always find clusters, even from random data points!)

UNIVARIATE TESTS




Fig. 6.1 Frierfjord macrofauna. Means and 95% confidence intervals for Shannon diversity ( $H'$ ) at 6 field sites

ONE-WAY ANOVA provides a test of the (null) hypothesis:

$H_0$ : No difference between sites

It assumes normality of  $H'$  and constant variance across sites (hence the confidence intervals in Fig. 6.1 use a pooled variance estimate and are of the same widths).

Table 6.1

Frierfjord macrofauna diversity  $H'$ ; ANOVA.

	Sum of squares	Deg. of freedom	Mean Square	F ratio	Sig. level
Treatments	3.938	5	0.788	15.1	<0.1%
Residual	0.937	18	0.052		
Total	4.874	23			

MULTIPLE COMPARISON TESTS are used to follow up a significant F-test with comparison between (all) pairs of sites, e.g.

TUKEY T TEST (i.e. a Least Significant Difference test) shows that the "reference" site A has significantly higher diversity than the rest, and C has a lower  $H'$  than E and G.

NOTE:

- 1) Multiple comparison tests FIX the PROBABILITY of TYPE I ERROR ("reject the null hypothesis when true") at 0.05 (say) over all pairwise comparisons.
- 2) Global F-test is best thought of as a "red light" - unless significant it BARS PROGRESS TO PAIRWISE COMPARISONS and interpretation of differences.
- 3) There are several implications for SAMPLE COLLECTION, which apply equally to the multivariate testing which follows:

IMPLICATIONS FOR DESIGN

- 1) CONTROL (reference) site(s) essential - impact only established by reference to similar unimpacted site(s), or to same site pre-impact. (Preferable to have both spatial and temporal controls).
- 2) REPLICATION at each site essential - should be over appropriate spatial scale (i.e. genuinely representative of that location).
- 3) "BLIND" ANALYSIS desirable - avoids (unconscious) biases, e.g. tendency to uniformity of replicates.

MULTIVARIATE TESTS

INFORMAL: CLUSTER, MDS, etc. assume no knowledge of how samples are divided into sites. So, plots can be inspected for evidence of REPLICATE GROUPING.




Fig. 6.2 Frierfjord macrofauna. MDS plot (Bray-Curtis similarities,  $\sqrt{\sqrt{\quad}}$  transform), for 24 samples, 4 replicates from each of sites A-E,G




Fig. 6.3 Frierfjord macrofauna. Dendrogram for 24 samples (similarities as for Fig. 6.2)

#### PARAMETRIC TESTS

EXACT ANALOGUE OF ONE-WAY ANOVA is multivariate analysis of variance (MANOVA), the F-test being replaced by WILKS'  $\Lambda$  test (e.g. Mardia, 1979). Pairwise differences can be tested by MAHALANOBIS' DISTANCES (e.g. Seber, 1984); but

ASSUMPTIONS RARELY SATISFIED: Tests require multivariate normality of abundances and "large samples" (at each site). For Frierfjord macrofauna, even after reduction to 30 species:

- a) 50% of abundances are zero - normality impossible (even with transform),
- b) ratio of observations to parameters needing estimation is 1.1 - hardly large!

RANDOMISATION/PERMUTATION TESTS:




Fig. 6.4 Frierfjord macrofauna. MDS plot (Bray-Curtis,  $\sqrt{\sqrt{\quad}}$  transform) of 4 replicates from B,C,D

NULL HYPOTHESIS  $H_0$ : no difference between sites. If  $H_0$  false, distances between replicates within sites are less than distances across sites. So:

- 1) COMPUTE STATISTIC reflecting this difference. To derive its sampling distribution, note that when  $H_0$  true, the 12 labels (4 B's, 4 C's, 4 D's) could be allocated at random to the 12 MDS points. So:
- 2) RECOMPUTE STATISTIC under ALL POSSIBLE PERMUTATIONS of the 12 labels between the 12 MDS points, or (since that is prohibitive) under a LARGE NUMBER OF RANDOM ALLOCATIONS of the 12 labels to the points.
- 3) RANDOMISATION/PERMUTATION TEST will reject  $H_0$  AT 5% SIGNIFICANCE LEVEL if observed statistic greater than its value for 95% of the random relabellings.

FORM OF DISPLAY SHOULD BE IRRELEVANT: Desirable that the statistic has exactly the same value whether the representation is:

- a) a dendrogram (Fig. 6.3)
- b) an MDS for all 6 sites (Fig. 6.2) or just a subset of sites (Fig. 6.4)
- c) an MDS in 3-d, say, rather than 2-d.

Bearing in mind that MDS is a function only of rank (dis)similarities, this suggests:

STATISTIC based on DIFFERENCE IN AVERAGE RANK DISSIMILARITIES between and within sites, i.e.

$$R = (\bar{r}_{\text{Between}} - \bar{r}_{\text{Within}})/(M/2) \quad (6.1)$$

where  $M = n(n-1)/2$  ( $n =$  total number of samples) and:

$R = 1$  if all replicates within sites are more similar than any replicates between sites.

$R = 0$  represents the null hypothesis.

( $R < 0$  possible, but only significantly so if experimental design incorrectly specified).

PAIRWISE COMPARISONS OF SITES: If global test rejects  $H_0$ , then same type of test can be carried out on each pair of sites, though note:

- a) These tests must be treated with some caution since NOT true "MULTIPLE COMPARISON" TESTS; overall Type I error not controlled.
- b) Minimum of 4 replicates per site needed for pairwise tests. Can be fewer for global test since NUMBER OF DISTINCT PERMUTATIONS is:


$$(\sum_i n_i)! / (n_1! n_2! \dots n_k! k!) \quad (6.2)$$

where  $\{n_i\}$  replicates at site  $i$  ( $i=1,2,\dots,k$ ).

Example: 2 replicates at each of 2 sites (A,B)

	A	A	B	B
Sample	1	2	3	4
A 1	-			
A 2	2	-		
B 3	4	3	-	
B 4	6	5	1	-

Rank dissimilarities




MDS (1,2 = A; 3,4 = B)



$\bar{r}_{\text{Between}} = 4.5$ ,  $\bar{r}_{\text{Within}} = 1.5$ ,  $M = 6$ , so  $R = 1$ .

Only three possible distinct PERMUTATIONS OF LABELS:



	A	A	B	B
Smp.	1	2	3	4
A 1	-			
A 2	2	-		
B 3	4	3	-	
B 4	6	5	1	-

$R = 1$

	A	A	B	B
Smp.	1	3	2	4
A 1	-			
A 3	4	-		
B 2	2	3	-	
B 4	6	1	5	-

$R = -0.5$

	A	A	B	B
Smp.	1	4	2	3
A 1	-			
A 4	6	-		
B 2	2	5	-	
B 3	4	1	3	-

$R = -0.5$

Observed case ( $R = 1$ ) has 33% probability of occurring by chance, so could not reject the hypothesis of "no difference between sites" (even though the observed case is the most extreme possible, here)

A more realistic example, where there are 12 samples divided between 3 sites (and thus  $12!/(4!4!4!3!) = 5775$  possible permutations) is given by Fig. 6.4:

Example: Frierfjord macrofauna abundances.

Table 6.2

Frierfjord macrofauna. Ranked dissimilarity matrix (Bray-Curtis,  $\sqrt{}$  transform) between the 12 replicates from sites B,C,D.

	B1	B2	B3	B4	C1	C2	C3	C4	D1	D2	D3	D4
B1	-											
B2	33	-										
B3	8	7	-									
B4	22	11	19	-								
C1	66	30	58	65	-							
C2	44	3	15	28	29	-						
C3	23	16	5	38	57	6	-					
C4	9	34	4	32	61	10	1	-				
D1	48	17	42	56	37	55	51	62	-			
D2	14	20	24	39	52	46	35	36	21	-		
D3	59	49	50	64	54	53	63	60	43	41	-	
D4	40	12	18	45	4	27	26	31	25	2	13	-

GLOBAL TEST:

$\bar{r}_{\text{Between}} = 37.54$ ,  $\bar{r}_{\text{Within}} = 22.72$ ,  $M = 66$ , so  $R = 0.45$ .

In 500 random relabellings, none of them gave  $R > 0.45$ , so  $H_0$  rejected at significance level  $p < .002$  (0.2%).

PAIRWISE TESTS:

For each pair of sites, the corresponding subset of the above triangular matrix is extracted, re-ranked and  $R$  computed as above, e.g. for  $B$  v  $C$ ,  $R = 0.23$ . This time,  $R$  can be re-evaluated for all possible relabellings, giving  $p < 12\%$ , so  $B$  &  $C$  not significantly different (only 35 distinct permutations, so the maximum attainable significance level is 3%).

However,  $D$  does differ from  $B$  and  $C$  ( $B$  v  $D$ :  $R = 0.54$ ,  $p < 3\%$ ,  $C$  v  $D$ :  $R = 0.57$ ,  $p < 3\%$ ).

FURTHER FEATURES AND EXTENSIONS:

- 1) PERMUTATION TEST CONCEPT dates to Mantel (1967) and general Monte Carlo (randomisation) tests discussed by Hope (1968). Practicals use a FORTRAN program called ANOSIM (Analysis of Similarities).
- 2) ANOSIM test makes NO ASSUMPTION OF "EQUAL VARIANCE"

Example: Coral communities at South Tikus, Thousand Is., Indonesia (Warwick, Clarke & Suharsono, 1990).




Fig. 6.5 MDS for % cover of coral species (Bray-Curtis, no transform) for 10 replicates in each of 5 years: 1 = 1981 (pre-El Niño), 3 = 1983 etc

ANOSIM test distinguishes the clear difference in initial and impacted conditions (1 and 3), though change is largely in variance rather than location.

- 3) ANOSIM TEST NOT RESTRICTED TO BALANCED REPLICATION at sites (or times); some sites can even have only one replicate provided enough replicates overall to generate sufficient permutations (eqt. (6.2)).
- 4) WIDE APPLICABILITY in that ANOSIM can be used with any (dis)similarity matrix; e.g. for a Euclidean distance matrix (appropriate to a PCA) ANOSIM can be seen as a non-parametric alternative to the parametric Wilks'  $\Delta$  test for a MANOVA, though it:

- 5) LACKS SENSITIVITY (as with many non-parametric tests) in the (unlikely) event that the data is genuinely multivariate normal.
- 6) ANOSIM PROGRAM EXTENDS TO ANALOGUE OF 2-WAY ANOVA:

2-WAY NESTED MODEL:

Example is Oslo Workshop macrofauna data from the mesocosm experiment: 2 cores from each of 4 boxes from each of 4 treatments.

TEST OF "BOX EFFECTS" involves calculating, separately for each treatment, the 1-way ANOSIM statistic for box differences, and then averaging across treatments. The sampling distribution comes from a restricted randomisation, with permutations preserving treatment designations.

The rank dissimilarity matrix is then reformed for a TEST OF TREATMENT EFFECTS by 1-way ANOSIM.

2-WAY CROSSED MODEL:

Example here would be several sites examined at several times. Can test for any overall differences between times (allowing for site differences by restricting permutations within sites). Alternatively test for overall differences between sites (allowing for differences in times).

RECOMMENDATIONS

- 1) USE RANDOMISATION/PERMUTATION TEST (ANOSIM) rather than parametric methods for testing of multivariate differences between previously-defined groups of samples (i.e. sites, times, treatments etc.); its ROBUSTNESS (lack of assumptions) more than makes up for its CONSERVATISM - latter is not so bad anyway. (Note: cannot test if differences between groups of samples are 'significant', if the grouping came from multivariate analysis of that same data).
- 2) USE NORMALITY-BASED TESTS for univariate INDICES, after any necessary transform (see lecture 9).

LECTURE 7

MULTIVARIATE METHODS: SPECIES ANALYSES

SPECIES CLUSTERING

Clustering methods can be applied to SPECIES SIMILARITY matrices (latter defined on pages 85-86).

Example: R. Exe (UK) nematodes, Field *et al.* (1982)




Fig. 7.1 Exe nematodes. Dendrogram (group average link) from Bray-Curtis similarities (standardised abundance data) for 55 species from 19 sites - reduced from 182 species by including those with counts >4% of total at any one site. The 4 to 5 groups indicated correspond closely with sharply defined clusters in the sites analysis (Fig. 5.3)

### SPECIES MDS

A species similarity matrix can also be input to an MDS, in the same way as for samples. In practice, often gives high 2-d stress. As with clustering, works best when samples form strong groups, arising from species sets which tend to be exclusive.




Fig. 7.2 Exe nematodes. MDS of 55 commonest species using Bray-Curtis similarities on standardised abundances. Main groups from cluster analysis (Fig. 7.1) indicated; they correspond closely to groupings of sites (Fig. 5.4)

Note: The LESS-COMMON SPECIES will generate erratic similarities, giving isolated MDS points and an unhelpful plot - they need to be REMOVED initially.

However, SPECIES clustering or ordination is generally less informative than methods which HIGHLIGHT SPECIES contributing to pattern of SAMPLE clustering or ordination:

### DETERMINING DISCRIMINATING SPECIES

Given clear CLUSTERING of SAMPLES, what methods will determine SPECIES RESPONSIBLE for groupings? Hard to see patterns in the original data matrix, so:



From (2.11), contribution to  $\delta_{jk}$  from ith species is:

$$\delta_{jk}(i) = 100 \cdot |y_{ij} - y_{ik}| / \sum_{l=1}^P (y_{ilj} + y_{ilk}) \quad (7.1)$$

$\delta_{jk}(i)$  then averaged over all pairs (with j in 1st and k in 2nd group), to give AVERAGE CONTRIBUTION  $\bar{\delta}_i$  from ith species (& its standard deviation SD ( $\delta_i$ )).

DISCRIMINATING SPECIES are those with HIGH  $\bar{\delta}_i$  and HIGH ratio  $\bar{\delta}_i/SD(\delta_i)$  (this implies CONSISTENCY of contributions across all jk pairs).

Table 7.1

Bristol Channel zooplankton ( $\sqrt{}$  counts). Species contributions  $\bar{\delta}_i$  to total average dissimilarity ( $\bar{\delta} = \sum \bar{\delta}_i = 59.5$ ) between site groups 1 & 2;  $\sum \bar{\delta}_i$  % is cumulative % contribution to  $\bar{\delta}$ . \* denotes good discriminators of groups 1 & 2.

Sp.	Name	$\bar{\delta}_i$	SD( $\delta_i$ )	$\bar{\delta}_i/SD(\delta_i)$	$\sum \bar{\delta}_i$ %
6	<i>Eurytemora affinis</i>	7.7	2.8	2.7*	13.0
4	<i>Centropages hamatus</i>	7.3	4.4	1.7*	25.2
3	<i>Calanus helgolandicus</i>	6.8	4.0	1.7*	36.7
1	<i>Acartia bifilosa</i>	5.7	4.0	1.4*	46.3
23	<i>Temora longicornis</i>	5.6	3.3	1.7*	55.6
18	<i>Pseudocalanus elongatus</i>	4.7	1.5	3.1*	63.5
13	<i>Paracalanus parvus</i>	3.3	4.2	0.8	69.1
15	<i>Pleurobrachia pileus</i> jv	3.1	2.8	1.1	74.3
20	<i>Sagitta elegans</i> jv	2.9	1.9	1.6*	79.1
19	<i>Sagitta elegans</i> jv	2.1	1.6	1.3	82.5
8	<i>Gastrosaccus spinifer</i>	2.0	1.8	1.1	85.9
14	<i>Pleurobrachia pileus</i>	1.9	1.6	1.2	89.0
10	<i>Mesopodopsis slabberi</i>	1.7	1.4	1.3	91.9
21	<i>Schistomysis spiritus</i>	1.6	1.4	1.1	94.5
17	<i>Polychaete larvae</i>	1.5	1.3	1.2	97.1
2	<i>Acartia clausi</i>	0.7	1.8	0.4	98.3
.	.....	...	...	...	....

Can similarly compute the contribution of the ith species ( $\bar{S}_i$ ) to the AVERAGE SIMILARITY WITHIN A GROUP ( $\bar{S}$ ), using the 2nd form of (2.1). This highlights species consistently prominent in that group (i.e. HIGH  $\bar{S}_i$ , HIGH ratio  $\bar{S}_i/SD(S_i)$ ).



Table 7.2

Zooplankton. Species contribution ( $\bar{S}_i$ ) to average similarity ( $\bar{S} = 66.3$ ) within site group 1.

Sp.	Name	$\bar{S}_i$	SD( $S_i$ )	$\bar{S}_i/SD(S_i)$	$\Sigma \bar{S}_i$ %
6	<i>Eurytemora affinis</i>	19.3	6.3	3.1*	29.1
18	<i>Pseudocalanus elongatus</i>	14.7	2.7	5.4*	51.3
1	<i>Acartia bifilosa</i>	12.2	6.4	1.9*	69.6
17	<i>Polychaete larvae</i>	3.9	3.1	1.2	75.5
14	<i>Pleurobrachia pileus</i>	3.4	3.8	0.9	80.7
21	<i>Schistomysis spiritus</i>	3.3	3.6	0.9	85.7
15	<i>Pleurobrachia pileus</i> jv	3.3	4.7	0.7	90.7
.	.....	...	...	...	....

**RECOMMENDATION**

USE SIMILARITY % BREAKDOWN (program SIMPER) or a SHADE MATRIX to INDICATE (not test) which species are mainly responsible for an observed clustering of the samples into groups (or for a confirmed difference between previously-defined groups).

## LECTURE 8

### UNIVARIATE AND DISTRIBUTIONAL METHODS: DIVERSITY MEASURES, DOMINANCE CURVES AND OTHER GRAPHICAL ANALYSES

#### INDICES OF DIVERSITY AND EVENNESS

A single index of species (or higher taxon) diversity is commonly employed in community studies, and is amenable to simple statistical analysis. A bewildering variety of diversity indices has been used, and it is not appropriate here to discuss their relative merits and disadvantages. A good account can be found in Heip *et al.* (1988).

Two different aspects contribute to the concept of community diversity:

**SPECIES RICHNESS** - A measure related to the total number of species present.

**EQUITABILITY** - Expresses how evenly the individuals are distributed among different species.

The most commonly used diversity measure is the SHANNON-WIENER INDEX:

$$H' = - \sum_i p_i (\log p_i)$$

This incorporates both the species richness and equitability components. Note that logarithms to the base 2 are often used in the calculation, giving the diversity units as 'bits per individual'.  $\log_e$  is also frequently used, so care should be exercised when comparing published indices.

**SPECIES RICHNESS** is often given simply as the total number of species (S), which is obviously very dependent on sample size, but more commonly as MARGALEF'S INDEX d, which also incorporates the total number of individuals (N):

$$d = (S-1) / \log N$$

**EQUITABILITY** is most commonly expressed as PIELOU'S EVENNESS INDEX:

$$J' = H'(\text{observed}) / H'_{\max}$$

where  $H'_{\max}$  is the maximum possible diversity ( $\log S$ ).

#### UNITS OF MEASUREMENT

Numbers of individuals belonging to each species are the most common units. For internal comparative purposes other units could be used, e.g. biomass or total cover of each species along a transect (e.g. for hard-bottom epifauna).

REPRESENTING COMMUNITIES

Data usually presented as plots of means and confidence intervals for each site or time.

Example 1: Benthos from Hamilton Harbour, Bermuda.




Fig. 8.1 Diversity ( $H'$ ) and 95% confidence intervals for macrobenthos (left) and meiobenthic nematodes (right) at six stations

Example 2: Reef-corals from South Tikus Island, Indonesia.




Fig. 8.2 Total number of species ( $S$ ), Diversity ( $H'$ ) and Evenness ( $J'$ ) based on coral species cover data along transects, spanning the 1982-3 El Niño. Note dramatic decline and partial recovery of  $S$  and  $H'$ , but no obvious changes in  $J'$

### DISCRIMINATING SITES OR TIMES

The significance of differences in diversity indices between sampling sites or times can be tested by one-way analysis of variance (ANOVA).

### DETERMINING STRESS LEVELS

Increasing levels of environmental stress are generally considered to:

- DECREASE diversity (e.g.  $H'$ )
- DECREASE species richness (e.g.  $d$ )
- DECREASE evenness (e.g.  $J'$ ), i.e. INCREASE dominance

Comparisons of measured indices can be made:

- with reference to comparative stations along a spatial contamination gradient (e.g. Fig. 8.1).
- with reference to comparative historical data (e.g. Fig. 8.2).
- with reference to some theoretical expectation of diversity, given the number of individuals and species present. Comparisons of observed diversity have been compared with predictions from CASWELL'S NEUTRAL MODEL (Caswell, 1976), which assumes certain community assembly rules and no interactions between species. A value of zero for the V statistic indicates neutrality, positive values indicate greater diversity than predicted and negative values lower diversity. Values  $>+2$  or  $<-2$  indicate significant departures from neutrality. The computer program of Goldman & Lamshead (1989) is useful.

Example: V statistics for summed replicates of macrobenthos and meiobenthic nematode samples at six stations in Hamilton Harbour, Bermuda (cf. Fig. 8.1)

STATION	MACROBENTHOS	NEMATODES
H6	-1.3	-0.4
H2	+0.5	-0.1
H7	-0.2	-0.4
H4	-4.5	-0.5
H3	-5.4	+0.4
H5	-1.9	0.0

Note diversity of macrobenthos at H4 and H3 is significantly below neutral model predictions, but nematodes are close to neutrality at all stations.

## GRAPHICAL DISTRIBUTION PLOTS

The purpose of graphical/distributional representations is to extract information on patterns of relative species abundances without reducing that information to a single summary statistic, such as a diversity index. This class of techniques can be thought of as intermediate between *univariate* summaries and full *multivariate* analyses. Unlike multivariate methods, these distributions may extract universal features of community structure which are not a function of the specific taxa present, and may therefore be related to levels of biological stress.

### RAREFACTION CURVES

Rarefaction curves (Sanders, 1968) were among the earliest to be used in marine studies. They are plots of the number of individuals on the x-axis against the number of species on the y-axis. The more diverse the community is, the steeper and more elevated is the rarefaction curve.

Example: Polychaete/bivalve fraction of macrobenthos.




Fig. 8.3 Rarefaction curves comparing North Sea and Friday Harbor stations (from Buchanan & Warwick, 1974)

RANKED SPECIES ABUNDANCE (DOMINANCE) CURVES

These are based on the ranking of species (or higher taxa) in decreasing order of their importance in terms of abundance or biomass. The ranked abundances, expressed as a percentage of the total abundance of all species, are plotted against the relevant species rank. Log transformations of one or both axes have frequently been used to emphasise or downweight different sections of the curves.




Fig. 8.4 The same (hypothetical) species abundance data plotted as ranked species abundance curves with none, one or both axes on a log scale (from Heip *et al.*, 1988)

### k-DOMINANCE AND LORENZ CURVES

As an alternative to the simple dominance curves above, cumulative ranked abundances may be plotted against species rank, or log species rank, to produce k-DOMINANCE CURVES (Lamshead et al., 1983). This has a smoothing effect on the curves. Ordering of curves on a plot will obviously be the reverse of rarefaction curves, with the most elevated curve having the lowest diversity. To compare dominance separately from the number of species, the x-axis (species rank) can be rescaled from 0-100 (relative species rank), to produce LORENZ CURVES.

Example: Nematodes from Loch Ewe, Scotland.




Fig. 8.5 k-dominance curves (left) and Lorenz curves (right) for 20 cm deep cores taken from experimental sand columns 20 days (A) and 77 days (B) after initial setup, and from intertidal (F) and subtidal (S) sand from the study site (from Lamshead et al., 1983)

### ABUNDANCE / BIOMASS COMPARISON (ABC) PLOTS

The advantage of distribution plots such as k-dominance curves is that the distribution of species abundances among individuals and the distribution of species biomasses among individuals can be compared on the same terms. Since the two have different units of measurement, this is not possible with diversity indices.

This is the basis of the ABUNDANCE / BIOMASS COMPARISON (ABC) method of determining levels of disturbance (pollution-induced or otherwise) on benthic macrofauna communities. Both empirical evidence and theoretical considerations suggest that the k-dominance curve for biomass will fall above the curve for abundance in undisturbed (or unpolluted) communities, and *vice versa* for disturbed (or polluted) communities.




Fig. 8.6 Hypothetical k-dominance curves for species biomass and numbers, showing unpolluted, moderately polluted and grossly polluted conditions (from Warwick, 1986)

Example 1: Time series of macrobenthos in Loch Linnhe, Scotland in response to increasing and decreasing levels of organic enrichment (pulp-mill effluent). See Lecture 1, Figs. 1.3 and 1.4

Example 2: Transect across sewage-sludge dumping ground at Garroch Head, Firth of Clyde, Scotland.






Fig. 8.7 Map showing location of dumping-ground. Centre of dump-site denoted by dashed circle: positions of sampling stations (P1 - P12) identified by asterisks

\*




Fig. 8.8 ABC plots for macrobenthos on Garroch Head transect in 1983. Abundance = squares, biomass = crosses (From Warwick *et al.*, 1987)

TRANSFORMATIONS OF k-DOMINANCE CURVES

PROBLEM: It is difficult to distinguish differences between k-dominance curves when cumulative frequencies are near 100% (sometimes after the first 2 or 3 spp.).

SOLUTION: Transform y-axis so that cumulative values are close to linearity. Clarke (1990) suggests the modified *logistic* transformation:

$$y_i' = \log[(1 + y_i)/(101 - y_i)]$$

Example: Macrobenthos from Frierfjord / Langesundfjord, Norway (IOC/GEEP Oslo Workshop).




Fig. 8.9 a), b) Standard ABC plots for sites A (reference) and C (potentially impacted). c), d) ABC plots for sites A and C with the y-axis subjected to modified logistic transformation. Abundance = continuous line, biomass = dashed line

### PARTIAL DOMINANCE CURVES

**PROBLEM:** Visual information presented by k-dominance (and ABC) curves is over dependent on single most dominant species. Unpredictable presence of large numbers of small biomass species, or heavy spatfall of young of one species, may give false impression of disturbance.

**SOLUTION:** With genuine disturbance, patterns of ABC curves should be unaffected by successive removal of most dominant species in terms of abundance or biomass. PARTIAL DOMINANCE CURVES (Clarke, 1990) compute the dominance of the second most dominant species over the remainder, the same with the third most dominant etc.

Example 1: Macrobenthos from Frierfjord/Langesundfjord, Norway (IOC/GEEP Oslo Workshop).




Fig. 8.10 Partial dominance curves (abundance/biomass comparison) for reference station A (c.f. Figs 8.9a and c for corresponding standard and transformed ABC plots). This illustrates the typically undisturbed condition)

Example 2: Loch Linnhe macrobenthos, 1966-68, 1970-72.




Fig. 8.11 a)-f) ABC curves (logistic transform). g)-l) Partial dominance curves for abundance (solid line) and biomass (dashed line) for the same years

SIGNIFICANCE TESTING FOR GRAPHICAL METHODS

Given replicate curves (k-dominance, ABC, 'individuals amongst species' etc.) at 2 or more sites (or times etc.), need a TEST FOR SIGNIFICANT DIFFERENCE.

Example: Hamilton Harbour macrofauna.




Fig. 8.12 Abundance k-dominance curves for four replicates at site H4 (solid) and H6 (dashed line)

Is the apparent difference for H4 and H6, in initial slope of curves, borne out statistically?

Also, testing for difference between sets of ABC CURVES at two (or more) sites reduces to a comparison of two (or more) sets of replicate curves by computing the DIFFERENCE CURVE B-A for each sample, e.g. Fig. 8.13.




Fig. 8.13 Difference (B-A) between k-dominance curves for biomass and abundance for four replicate samples at H2 (solid) and H4 (dashed line)

FIRST APPROACH:

Reduce each replicate curve to a SINGLE SUMMARY STATISTIC. E.g. if  $\{A_i\}$  and  $\{B_i\}$  are the cumulative abundance and biomass values from an ABC plot ( $i=1, \dots, S$  species), define:

$$W = \sum_{i=1}^S (B_i - A_i) / [50(S-1)]$$

W takes values in  $(-1, 1)$ , with  $W=1$  for totally even abundances across species but biomass dominated by a single species, and  $W=-1$  for the converse case.

Similarly, for k-dominance curves of cumulative  $\{A_i\}$ :

$$K_A = [(\sum_{i=1}^S A_i) - 50(S+1)]/[50(S-1)]$$

where extremes are K=0 (evenness) and K=1 (dominance).  $K_b$  defined similarly for biomass.

Now, PERFORM ANOVA on SUMMARY STATISTICS (W or K) from each replicate (e.g. as for diversity indices). Works well in cases like Fig. 8.13 (H2 & H4 differ significantly) but poorly for Fig. 8.12 where difference is in slope not mean area. Need more GENERAL TEST with power to detect any CONSISTENT DIFFERENCE between 2 (or more) sets of curves, so

#### SECOND APPROACH:

Define 'dissimilarity' between any pair of curves  $\{A_{i1}; i=1, \dots, S_1\}$ ,  $\{A_{i2}; i=1, \dots, S_2\}$ , as their total (absolute) distance apart:

$$d = \sum_{i=1}^{S_{\max}} |A_{i1} - A_{i2}|$$

where  $S_{\max} = \max(S_1, S_2)$ . Or better reflection of visual difference in two k-dominance curves is:

$$d' = \sum_{i=1}^{S_{\max}} |A_{i1} - A_{i2}| \log(1+i^{-1})$$

Compute d (or d') for every pair of replicate curves, to give lower triangular dissimilarity matrix, and CALCULATE ANOSIM STATISTIC R, eqt. (6.1).

PERMUTATION/RANDOMISATION TEST of difference between sites/times etc. then carried out exactly as in lecture 6. (ANOSIM on Fig. 8.12 distinguishes H4 and H6, whereas ANOVA on  $K_A$  does not).

Details in Clarke (1990). Note that principle EXTENDS TO OTHER GRAPHICAL METHODS, e.g. partial dominance, 'individuals amongst species' curves etc.



LECTURE 9  
TRANSFORMATIONS

There are two distinct roles for transformations in community analysis:

- a) to validate assumptions for parametric analyses - applies to UNIVARIATE tests
- b) to weight the contributions of common and rare species in a MULTIVARIATE representation.

UNIVARIATE

Example: Frierfjord macrofauna. Indicator species.

Table 9.1

*Thyasira* sp. numbers in 4 replicate grabs at 6 sites.

Site:	A	B	C	D	E	G
Replicate						
1	1	7	0	1	62	66
2	4	0	0	8	102	68
3	3	3	0	5	93	52
4	11	2	3	13	69	36
Mean	4.8	3.0	0.8	6.8	81.8	55.5
Stand.dev.	4.3	2.9	1.5	5.1	18.7	14.8

NOTE:

- 1) The replicates are not symmetrically distributed (they tend to be right-skewed), so normality assumptions are dubious.
- 2) More importantly (for test validity), the variance increases strongly with the mean - this invalidates "constant variance" assumptions of ANOVA.

Both problems can be tackled by:

POWER TRANSFORMATION

Individual replicates  $y$  are transformed to  $y^*$ , given by:

$$y^* = (y^\lambda - 1) / \lambda \quad (9.1)$$

where, in order of INCREASING SEVERITY,

- $\lambda = 1$  - no transform
- $\lambda = 0.5$  - square root ( $\sqrt{\quad}$ )
- $\lambda = 0.25$  - 4th root ( $\sqrt[4]{\quad}$ )
- $\lambda \rightarrow 0$  - log transform ( $y^* = \log_e y$ )

Possible to determine best  $\lambda$ , anywhere in (0,1), for each separate data set (Box and Cox, 1964), but unnecessarily precise - better just to choose between above 4 cases, using:

TAYLOR'S POWER LAW:

If:  $\text{var}(y) \propto (\text{mean } y)^v \quad (9.2)$

then:  $\text{var}(y^\lambda) \propto (\text{mean})^{2(\lambda-1)+v}$  (approx.)  $(9.3)$

Choose  $\lambda = 1 - (v/2)$  to get  $\text{var}(y) = \text{constant}$ .

Find  $v$  by regressing  $\log(\text{stand.dev.})$  on  $\log(\text{mean})$ , because:


$$\log(\text{sd}(y)) = (v/2)\log(\text{mean } y) + \text{constant} \quad (9.4)$$

So  $\lambda = 1 - (\text{slope of regression})$ , thus if:

- slope = 0 - no transform
- = 0.5 - use  $\sqrt{\quad}$
- = 0.75 - use  $\sqrt[4]{\quad}$
- = 1 - use  $\log_e$

$$(9.5)$$

Example: *Thyasira* numbers at 6 sites



Plot indicates  $\sqrt{\quad}$  appropriate. After transform:

Site	A	B	C	D	E	G
Mean( $y^*$ )	2.01	1.45	0.43	2.42	9.00	7.40
Sd ( $y^*$ )	0.97	1.10	0.87	1.10	1.04	1.04

VARIANCE STABILISED so ANOVA and follow-up tests VALID (show E,G different from the rest, clearly). Means and confidence intervals should be back-transformed to original scales (intervals not symmetric but then data was not symmetric).

CAUTION: Beware of doing multiple ANOVAs on a range of indicator species (each runs a 5% risk of error and this compounds). Alright if performed (at higher significance) on a few species selected a priori.

AVOID "SNOOPING" in a large data array for likely species to do an ANOVA on; certain to find some which are significant, even in a random array!

**MULTIVARIATE**

TRANSFORMS can be used for the same reason as in univariate analyses - to induce (multivariate) normality, eg. for MANOVA tests (lecture 6), but:

- a) Insufficient to demonstrate univariate normality and constant variance (for each variable) to prove multivariate normality and constant covariance.
- b) Rarely possible to achieve (marginal) normality for species abundance/biomass data (though possible for, say, a matching set of diversity indices).

MORE IMPORTANT USE OF TRANSFORMS IN COMMUNITY DATA is in WEIGHTING rare and common species in forming similarities between sites, eg. Bray-Curtis:

$$S_{jk} = 100 \left\{ 1 - \frac{\sum_{i=1}^P |y_{ij} - y_{ik}|}{\sum_{i=1}^P (y_{ij} + y_{ik})} \right\} \quad (9.6)$$

Example: Loch Linnhe macrofauna, subset

Sample:	1	2	3	4	<u>UNTRANSFORMED</u>
Species					Sample 1 2 3 4
<i>Echinoca.</i>	9	0	0	0	1 -
<i>Myrioche.</i>	19	0	0	3	2 8 -
<i>Labidopl.</i>	9	37	0	10	3 0 42 -
<i>Amaeana</i>	0	12	144	9	4 39 21 4 -
<i>Capitella</i>	0	128	344	2	
<i>Mytilus</i>	0	0	0	0	
Sample:	1	2	3	4	<u>√√ TRANSFORMED</u>
Species					Sample 1 2 3 4
<i>Echinoca.</i>	1.7	0	0	0	1 -
<i>Myrioche.</i>	2.1	0	0	1.3	2 26 -
<i>Labidopl.</i>	1.7	2.5	0	1.8	3 0 68 -
<i>Amaeana</i>	0	1.9	3.5	1.7	4 52 68 42 -
<i>Capitella</i>	0	3.4	4.3	1.2	
<i>Mytilus</i>	0	0	0	0	

Untransformed similarities are lower (unimportant in itself since MDS is only a function of ranks) but RANK SIMILARITIES ARE TOTALLY CHANGED by transform.

Untransformed similarities are DOMINATED BY THE COMMONEST SPECIES, eg. comparing samples 2 and 4 and omitting each species in turn:

Species omitted:	None	1	2	3	4	5	6
Bray-Curtis (S):	21	21	21	14	13	<u>54</u>	21

By contrast, under a  $\sqrt{y}$  transform, ALL (present) SPECIES MAKE SOME CONTRIBUTION to the similarity:

Species omitted:	None	1	2	3	4	5	6
Bray-Curtis (S):	68	68	75	61	59	76	68

TRANSFORMATION SEQUENCE:

None  $\longrightarrow$   $\sqrt{y}$   $\longrightarrow$   $\sqrt[3]{y}$   $\longrightarrow$   $\log$   $\longrightarrow$  Presence/absence

puts PROGRESSIVELY LESS WEIGHT on common species and increasingly takes account of rarer ones.

Logical end-point is REDUCTION of the data array to one of PRESENCE OR ABSENCE OF SPECIES (this is a transformation to the numbers 0 or 1), where all species contribute equally.

Example: Loch Linnhe macrofauna, subset.

Sample:	1	2	3	4	<u>PRESENCE/ABSENCE</u>
Species					
<i>Echino.</i>	1	0	0	0	Sample 1 2 3 4
<i>Myrioc.</i>	1	0	0	1	1 -
<i>Labido.</i>	1	1	0	1	2 33 -
<i>Amaeana</i>	0	1	1	1	3 0 80 -
<i>Capite</i>	0	1	1	1	4 57 86 67 -
<i>Mytilus</i>	0	0	0	0	

NOTE: 1) NEED TO USE  $\log(1+y)$  not  $\log y$  which DISTORTS TRANSFORM SEQUENCE.  $\log(1+y)$  intermediate between  $\sqrt[3]{y}$  and presence/absence for moderate or large counts but less severe than  $\sqrt[3]{y}$  for small counts.

- 2)  $\sqrt[3]{y}$  preferred to  $\log(1+y)$  because Bray-Curtis is INVARIANT TO A SCALE CHANGE (eg. for biomass) if  $\sqrt[3]{y}$  is used. (Little difference in practice though).
- 3) As severity of transform increases, more species contribute, so sample relationships are expressed in higher-dimensional space, and ordination in 2-d is harder (eg. Fig. 9.1). So, WRONG to assume that TRANSFORMS GIVING LOWER STRESS ARE BETTER; the converse is true if added species are important.




Fig. 9.1 GEEP mesocosm nematodes (Warwick et al., 1988). MDS of 4 boxes from 4 treatments (C,L,M,H). Bray-Curtis similarities from transformed counts: a) no transform, b)  $\sqrt{\quad}$ , c)  $\sqrt{\sqrt{\quad}}$ , d) presence/absence. Stress: a) 0.08, b) 0.14, c) 0.19, d) 0.19

- 4) SAME TRANSFORM SEQUENCE APPLIES TO PCA (and other ordinations) with much the same consequences.
- 5) Log (or  $\sqrt{\quad}$ ) transforms effectively REDUCE DATA TO A 6 POINT SCALE, i.e. 0 = absent, 1 = one individual, 2 = handful, 3 = sizeable, 4 = abundant, 5 = very abundant; replacing data by this scale will make no real difference to the multivariate displays. This may appear crude but often genuinely reflects inherent variability, so greater accuracy in counting may be unnecessary.

CONCLUDE:

- 1) CHOICE OF TRANSFORM often has a bigger effect on conclusions than the CHOICE OF ORDINATION method.
- 2) "What is the RIGHT TRANSFORM for a multivariate analysis?" is largely a BIOLOGICAL rather than a STATISTICAL question (unlike the use of transforms for validating assumptions); the choice of transform determines how the similarity of two samples is defined.

RECOMMEND:

Use INTERMEDIATE transform (eg.  $\sqrt{\quad}$ ,  $\sqrt[3]{\quad}$  or LOG) rather than either of the two EXTREMES:

- a) NO TRANSFORM - MDS reflects only 2 or 3 commonest species, so INTERPRETATION is likely to be SHALLOW.
- b) PRESENCE/ABSENCE - CHANCE OCCURRENCES of rare species DOMINATE the SAMPLE RELATIONSHIPS in high dimensions and make it difficult to get an interpretable low-dimensional ordination.

## LECTURE 10

### SPECIES REMOVAL AND AGGREGATION

#### SPECIES REMOVAL

Two reasons for ELIMINATING SPECIES discussed earlier:

- a) For sample PCA (not MDS) ordination, must reduce to (say) <50 species, else problems with eigenvalues.
- b) For species ordinations, though MDS and CLUSTER are possible for all species, rarer (chance) species must be excluded for an interpretable outcome.

RECOMMEND RETAINING SPECIES ACCOUNTING FOR >p% of total score (abundance or biomass) in ANY ONE SAMPLE (p chosen to reduce to required number, typically p = 3 or 4). Allows for high diversity/ low abundance samples which could have all species eliminated by simple selection of the top q% most abundant species over all samples.

SPECIES REDUNDANCY: Since sample relationships can often be well summarised in a 2-d ordination (from, say, a 100-d species space), many SPECIES MUST BE INTERCHANGEABLE in the way they characterise the samples. This can be seen by performing MDS on a randomly chosen subset (say 20%) of species:




Fig. 10.1 Frierfjord macrofauna counts. Sample MDS (Bray-Curtis,  $\sqrt{\sqrt{\quad}}$ ) for: a) all 110 species, b) 19 random species. (Stress: a) 0.14, b) 0.13)



Above example of no practical interest, but suggests:

SPECIES AGGREGATION to higher taxonomic levels.

If results from identifications to higher taxonomic levels are comparable to a full species analysis:

- a) a great deal of LABOUR CAN BE SAVED;
- b) LESS FAUNAL EXPERTISE NEEDED - major factor in parts of the world where fauna is poorly described.

METHODS AMENABLE TO AGGREGATION:

1) MULTIVARIATE:

All ordination/clustering techniques.

Empirical evidence is increasing that identification only to family level makes little difference.

2) DISTRIBUTIONAL:

- a) Aggregation for ABC curves is possible; family level analyses are often identical to species level analyses (see Figs. 10.6 and 10.7).
- b) Untried for other methods (eg. Individuals amongst species curves).

3) UNIVARIATE:

- a) Concept of "indicator groups" is well-established (eg. nematode/copepod ratios).
- b) Can define diversity indices at hierarchical taxonomic levels (though not commonly used in practice).

Warwick (1988) hypothesises further motivation: that pollution may change community composition at higher taxonomic levels (eg. phyla) whereas natural variables (grain size, water depth etc.) modify it more by species replacement (within phyla). Thus, distribution of higher taxa may even relate more closely to the contamination gradient than species data, the latter being more complicated by effects of confounding natural variables.

MULTIVARIATE EXAMPLES




Fig. 10.2 Mesocosm copepod counts - 3 levels of nutrient enrichment (Gee et al., 1985). Sample MDS plot (Bray-Curtis,  $\sqrt{\sqrt{\quad}}$  transform); species data aggregated into genera and families (Warwick, 1988)




Fig. 10.3 Loch Linnhe macrofauna (Pearson, 1975). MDS (Bray-Curtis) of 11 years samples for  $\sqrt{\sqrt{}}$  transform (left) and no transform (right), based on abundances from 115 species (top), aggregated into 45 families (middle) and 9 phyla (bottom), Warwick (1988). Note more linear configuration for phyla




Fig. 10.4 MDS for macrobenthos at station "Pierre Noire". Species data (left) aggregated into phyla (right). Sampling months are A:4/77, B:8/77, C:9/77, D:12/77, E:2/78, F:4/78, G:8/78, H:11/78, I:2/79, J:5/79, K:7/79, L:10/79, M:2/80, N:4/80, O:8/80, P:10/80, Q:1/81, R:4/81, S:8/81, T:11/81, U:2/82. Oil-spill was during 3/78, i.e. between E and F. Note more linear configuration for phyla




Fig. 10.5 MDS for coral species (n=75) and genus (n=24) cover data at South Pari Island, Indonesia. El Niño occurred in 1982-3. 1=1981, 3=1983, etc

GRAPHICAL/DISTRIBUTIONAL EXAMPLES




Fig. 10.6 Loch Linnhe macrofauna. (A) Diversity  $H'$ , (B)-(L) "ABC" curves for 11 years, of biomass (crosses) and abundance (squares). Analysis at species level, Warwick (1986)




Fig. 10.7 Loch Linnhe macrofauna. (A) Diversity  $H'$ , (B)-(L) "ABC" curves for 11 years, of biomass (crosses) and abundance (squares) for data aggregated to families, Warwick (1986)

UNIVARIATE EXAMPLE




Fig. 10.8 Plots of number of taxa and Shannon diversity for reef corals at South Tikus Island, Indonesia, showing impact and partial recovery from 1982-3 El Niño. Species data (upper) have been aggregated into genera (lower). Note similarity of patterns

## LECTURE 11

### LINKING MULTIVARIATE AND UNIVARIATE COMMUNITY ANALYSES TO ENVIRONMENTAL VARIABLES

#### APPROACH

- 1) FAUNAL AND ENVIRONMENTAL ANALYSIS SEPARATED initially, i.e. the biota is allowed to "tell its own story", without the use of physical or chemical data:
  - a) to DEMONSTRATE the RELATIONSHIPS between samples and differences (if any) between sites (/times),
  - b) to INFER COMMUNITY DISTURBANCE at some sites.
- 2) ENVIRONMENTAL VARIABLES ANALYSED ON THEIR OWN, for similar reasons. Two classes of variables:

"NATURAL" PHYSICAL (or "background") VARIABLES, such as depth of the water column, sediment granulometry, salinity, etc. and

CONTAMINANT VARIABLES, measuring chemical impact.

Analysis attempts:

  - a) to DEMONSTRATE DIFFERENCES (if any) in physical or chemical variables between the sites,
  - b) to REDUCE the COMPLEXITY of the environmental measures, particularly the chemical data, so the nature of the impact (if any) can be summarised by a few key variables.
- 3) SUMMARY REPRESENTATIONS of both biological and environmental analyses are VIEWED TOGETHER:
  - a) to examine whether changes between sites (/times) seem to be the product of differences in "natural" environmental variables, or
  - b) are correlated with inferred or measured contaminant impact.

#### ANALYSIS OF ENVIRONMENTAL DATA

UNIVARIATE: Background (physical) variables are typically univariate, with little variability between replicates within a site (e.g. water depth).

Where there is variability, and it is helpful to establish site differences, use ANOVA and confidence intervals (e.g. as for diversity).



MULTIVARIATE: Chemical measurements can often be highly multivariate (e.g. wide range of PAH compounds, PCB congeners, heavy metals etc.)

Example: Frierfjord sediment - heavy metals.

Table 11.1

Frierfjord sediments. Metal concentrations ( $\mu\text{g g}^{-1}$  dry wt, Fe as %) in top 2 cm from 3 replicate cores at sites A-E,G. Abdullah & Steffenak (1988).

Site	Cu	Zn	Pb	Ni	Cr	Cd	Mn	Fe
A	28	141	73	33	40	0.8	454	3.5
	26	139	71	30	40	(0.6)	653	3.3
	27	147	67	29	35	(0.6)	503	3.1
B	48	238	134	33	50	(0.6)	1050	3.5
	47	228	130	32	50	1.1	2880	3.5
	64	297	167	32	40	1.1	664	3.1
C	44	228	135	35	51	0.8	1500	4.1
	42	216	126	35	60	0.8	3570	4.2
	42	208	117	33	45	1.1	5880	4.0
D	48	241	142	37	56	0.9	1720	4.3
	39	205	114	33	50	0.8	8480	4.4
	44	238	141	35	34	1.1	5440	4.1
E	38	199	160	22	40	0.8	484	2.2
	40	241	156	25	40	1.1	925	2.1
	107	275	184	28	45	1.1	1400	2.5
F	48	328	118	32	35	3.6	10380	3.1
	44	296	110	30	35	3.1	5880	3.0
	47	320	118	32	35	3.4	7430	3.0
G	67	349	212	35	61	2.2	1060	2.8
	70	357	229	35	66	2.5	638	2.7
	77	417	267	38	70	4.5	619	2.6

SAME RANGE OF MULTIVARIATE METHODS AVAILABLE as for faunal analyses (replace species by chemical "species"). However, type of data is different:

- a) ZEROS do NOT predominate.
- b) distribution NOT highly RIGHT-SKEWED.
- c) REDUNDANCY can be very extreme, i.e. similar chemical compounds correlate very closely with each other along a spatial contaminant gradient.

So, possibly after (mild) TRANSFORMATION (e.g.  $\sqrt{\quad}$ ),

- a) MULTIVARIATE NORMAL assumptions often justified;
- b) PCA is useful, a 2-d ordination often giving a good representation of site chemistry,
- c) TESTING of site differences can either be by MANOVA (e.g. Wilks'  $\Lambda$ ) or by ANOSIM on a Euclidean distance dissimilarity matrix.

Example: Frierfjord sediment metals.




Fig. 11.1 Frierfjord sediments. 2-d PCA of metal data of Table 11.1 ( $\sqrt{\quad}$  - transformed and normalised)

NOTE, in Fig. 11.1:

- 1) First 2 PCs ACCOUNT FOR 69% OF VARIABILITY, so 2-d ordination is not too bad a representation.
- 2) Some DIFFERENCES BETWEEN SITES ( $p < 0.001$  in ANOSIM test), principally between A, G and the rest.
- 3) PC1 represents an AXIS OF INCREASING CONTAMINANT LOAD, the weights given to the (normalised) Cu, Zn, Pb, Ni, Cr, Cd, Mn, Fe levels being 0.41, 0.48, 0.46, 0.30, 0.35, 0.35, -0.05 and -0.21.
- 4) PC1 AXIS is thus a UNIVARIATE descriptor of the overall metal load, useful in relating this chemistry to faunal descriptions.
- 5) Though it exists, the CONTAMINANT GRADIENT is WEAK, no more than a factor of 2 or 3 between the extremes, A and G. (PAH gradient weaker still).

#### RELATION TO FAUNAL ANALYSES - FIRST APPROACH

SELECT at most 2 or 3 DESCRIPTORS of the CONTAMINANT GRADIENT (eg. one for metals, one for hydrocarbons) - even 2 or 3 could be ambitious if the different classes of contaminants are well-correlated.

The two cases considered below are when the biological data are UNIVARIATE (eg. diversity indices) and when they are MULTIVARIATE (eg. ordinations).

#### UNIVARIATE

REGRESSION is a possible technique: either SIMPLE LINEAR REGRESSION (1 environmental variable)

or MULTIPLE LINEAR REGRESSION (for 2 or more)

or NON-LINEAR REGRESSION (if there is a range of contaminant values and sufficient replicates to justify a more complex "dose-response" curve.)

Example: Frierfjord macrofauna.




Fig. 11.2 Frierfjord macrofauna abundances. Shannon diversity  $H'$  regressed on an overall measure of sediment metal concentration (latter is mean PCI at each of the 6 sites, from the PCA of Fig. 11.1). x - replicate grabs, — fitted regression line, --- 95% confidence "funnel" for the mean  $H'$  at any metal concentration

NOTE: Simple linear regression of  $H'$  on metal levels is not convincing!

- a) Slope just fails to differ significantly from zero, at 5%.
- b) Linear relation is not adequate (but data does not justify more complex fit).
- c) Most prominent feature (clear from the earlier ANOVA also - Fig. 6.1) is the general drop in diversity from the "reference" site (A).

MULTIVARIATE

SUPERIMPOSITION OF ENVIRONMENTAL VARIABLES OF FAUNAL ORDINATION: an effective visual technique performed separately for each environmental variable.

This may allow a GRADIENT in the ENVIRONMENTAL VARIABLE to be matched visually to a GRADIENT of change in the COMMUNITY structure.

Example: Bristol Channel zooplankton, April 1978.




Fig. 11.3 MDS of 57 sites (from Bray-Curtis similarities, on  $\sqrt{v}$  - transformed counts; stress = 0.11). For map of sites and corresponding cluster analysis, see Figs. 3.2 and 3.3

Though clear evidence of clusters (from Fig. 3.3), overall pattern is one of GRADATION of COMMUNITY STRUCTURE across the plot (note characteristic "arching", common for strong gradation).



Example: Frierfjord macrofauna counts ( $\sqrt{}$  - transformed)




Fig. 11.5 MDS of sites A-E,G with superimposed values of (a) water depth (22-113 m), (b) sediment median grain size (7.8-16.5  $\mu\text{m}$ ), (c) metal levels (PC1 in Fig. 11.1) and (d) "total" PAH (4.4-14.8  $\mu\text{g g}^{-1}$ )

- 1) Site grouping on the MDS bears LITTLE RELATION to the (weak) metal and PAH CONTAMINANT GRADIENTS.
- 2) Sediment granulometry is NOT A DETERMINANT of COMMUNITY DIFFERENCES here (B & C span the range of grain sizes but have the same communities).
- 3) DEPTH-RELATED differences between the sites appear to be the major CORRELATE of COMMUNITY DIFFERENCES. (Seasonal anoxia in the deeper parts of the fjord is likely to be a significant "stress" factor.)

Sometimes MORE THAN ONE AXIS OF CHANGE MAY BE SEEN, correlating with different environmental variables.

Example: Exe nematode abundances, Field et al. (1982)




Fig. 11.6 MDS of 19 sites (Fig. 5.1), with values of: (a) mean salinity of interstitial water (10-90% of standard seawater), (b) median sediment particle size (0.06-1.14 mm), superimposed at each site

Grain size forms a gradient from bottom left to top right, whereas salinity distinguishes the "middle" from the "end" sites along the first MDS axis.

Though the visual approach is generally more helpful, FORMAL TESTING of gradients can be performed by:

- a) REGRESSING each environmental variable on the (x,y) CO-ORDINATES of the SAMPLE LOCATIONS on the MDS; this would be multiple linear regression (and not appropriate for a curvilinear gradient).
- b) Using 2-WAY ANOSIM on sites (treated as replicates), which are categorised by, say, 2 environmental variables at 2 levels, e.g. deep/shallow, high/low contaminant loads. This would need a reasonable number of sites (with some in all 4 combinations).



RELATION TO FAUNAL ANALYSES - SECOND APPROACH

First approach designed mainly to show COMMUNITY pattern related to ONE ENVIRONMENTAL VARIABLE at a time. Alternative considers ALL environmental variables together and COMPARES ordination of biota to ORDINATION of environmental variables.

Example: Exe nematode abundances.




Fig. 11.7 (a) MDS of 19 sites (as in Fig. 5.1), (b) PCA of 4 environmental variables (salinity, median particle size, % organics, depth of H<sub>2</sub>S layer)

The close match of patterns shows these 4 variables "EXPLAIN" biota clusters (in Fig. 11.7a) well. Two questions: Would subset of environmental variables do as well? Would more variables do better? (e.g. height up shore, water table depth.)

Answer by DEFINING MATCH between two ordinations as some form of RANK CORRELATION ( $\rho$ ) between underlying DISSIMILARITY MATRICES (Bray-Curtis and Euclidean distance, respectively). Then find subset of environmental variables which MAXIMISES  $\rho$ . Here, this is the 4 variables in Fig. 11.7b.

IMPLICATIONS FOR DESIGN

- 1) SITE SELECTION: where there is choice, attempt to select sites such that VARIATION IN "NUISANCE" (physical) VARIABLES IS SMALL, (i.e. small enough not to have a significant affect on community structure).
  
- 2) Where between-site variation in natural variables is considerable, AVOID DESIGNS in which important physical variables are TOTALLY CONFOUNDED (i.e. run in parallel) with contaminant gradients. It may then be possible to DISTINGUISH SEPARATE PHYSICAL AND CONTAMINANT GRADIENTS in an MDS plot.  
  
(Alternatively, choose CONTROL SITES MATCHED to the PHYSICAL VARIABLES for each impacted site.)
  
- 3) Where within-site variation in natural variables is considerable (comparable with between-site), MDS distinction of contaminant and natural gradients is greatly AIDED by separate MEASUREMENT of environmental variables MATCHING EACH COMMUNITY REPLICATE.

## LECTURE 12

### CAUSALITY: COMMUNITY EXPERIMENTS IN THE FIELD AND LABORATORY

In experimental situations we can investigate the effects of a single factor (the TREATMENT) on community structure, while other factors are held constant or controlled. There are three main categories of experiments that can be used:

1. 'NATURAL EXPERIMENTS' - Nature provides the treatment: i.e. we compare places or times which differ in the intensity of the environmental factor in question.
2. FIELD EXPERIMENTS - The experimenter provides the treatment: i.e. environmental factors are manipulated in the field.
3. LABORATORY EXPERIMENTS - Environmental factors are manipulated by the experimenter in laboratory mesocosms or microcosms.

The degree of 'naturalness' (hence realism) decreases from 1-3, but the degree of control which can be exerted over confounding environmental variables increases from 1-3.

In all cases care should be taken to avoid PSEUDOREPLICATION, i.e. the treatments should be replicated, rather than a series of 'replicate' samples taken from a single treatment (pseudoreplicates). This is because other confounding variables, often unknown, may also differ between the treatments. It is also important to run experiments long enough for community changes to occur: this favours components of the fauna with short generation times (see Lecture 13).

#### NATURAL EXPERIMENTS

The obvious logical flaw with this approach is that its validity rests on the assumption that places or times differ only in the intensity of the selected environmental factor (treatment). Experimental design is often a problem, but statistical techniques such as TWO-WAY ANOVA or TWO-WAY ANOSIM, which enable us to examine the treatment effect allowing for differences between sites, are useful.

Example: The effects of disturbance by soldier crabs (*Mictyris platycheles*) on meiobenthic community structure.

LOCATION: Sand-flat at Eaglehawk Neck, S.E. Tasmania.

SAMPLING: Sediment disturbed by crabs in discrete patches.  $4 \times 5 \text{ m}^2$  blocks of 4 samples with each block including 2 disturbed and 2 undisturbed:




Fig. 12.1 Sketch showing the type of sample design. Sample positions (large dots) in relation to disturbed sediment patches (stippled)

UNIVARIATE INDICES:

Table 12.1

Mean values per core sample of univariate measures for nematodes, copepods and total meiofauna (nematodes + copepods) in the disturbed and undisturbed areas. The significance levels for differences are from a two-way ANOVA, i.e. they allow for differences between blocks, although these were not significant at the 5% level.

	Tot.ind.	Tot.sp.	d	H'	J'
<i>Nematodes</i>					
Disturbed	205	14.4	2.6	1.6	0.58
Undisturbed	200	20.1	3.7	2.2	0.74
Significance (%)	91	1	0.3	0.1	1
<i>Copepods</i>					
Disturbed	94	5.4	1.0	0.96	0.59
Undisturbed	146	5.7	1.0	0.84	0.49
Significance (%)	11	52	99	52	38
<i>Total meiofauna</i>					
Disturbed	299	19.8	3.4	2.0	0.66
Undisturbed	346	25.9	4.4	2.3	0.69
Significance (%)	48	1	3	3	16

For NEMATODES: significant reduction in total number of species, Species Richness, Shannon Diversity and Evenness in relation to disturbance.

For COPEPODS: no differences in any of these univariate measures.

GRAPHICAL/DISTRIBUTIONAL PLOTS




Fig. 12.2 Replicate k-dominance curves for NEMATODE abundance in each sampling block. D = disturbed, U = undisturbed

Summary statistics  $K_A$  and R (see Lecture 8) both show significant treatment effect when tested with two-way ANOSIM.

For COPEPODS (figure not given here), k-dominance curves are intermingled and crossing, and there is no significant treatment effect on  $K_A$  and R.

MULTIVARIATE ANALYSIS:




Fig. 12.3 MDS configurations for nematode, copepod and 'meiofauna' (nematode + copepod) abundance.  
Circles = Block 1, Squares 2, Pentagons 3, Diamonds 4.  
Open symbols = disturbed, shaded = undisturbed

Note similarities: both disturbed samples within each block are above both undisturbed; blocks arranged in sequence (left to right) 3,4,2,1.

Table 12.2

Results of the two-way ANOSIM test for treatment (disturbance/no disturbance) and block effects.

	DISTURBANCE		BLOCKS	
	R Statistic	Sig.(%)	R statistic	Sig. (%)
Nematodes	1.0	1.2	0.99	0.2
Copepods	0.56	3.7	0.70	0.2
Meiofauna	0.94	1.2	0.94	0.2

For both nematodes and copepods, two-way ANOSIM shows significant effect of both treatment (disturbance) and blocks, but differences more marked for nematodes (higher values of R statistic).

#### CONCLUSIONS:

Univariate indices and graphical/distributional plot only significantly affected by crab disturbance for nematodes. Multivariate analysis reveals similar response for nematodes and copepods (i.e. seems to be more sensitive). In multivariate analyses, natural variations in species composition across the beach (i.e. between blocks) were about as great as those between treatments within blocks: disturbance effect would not have been clearly evidenced without this block sampling design.

#### FIELD EXPERIMENTS

These include, e.g. caging experiments to exclude or include predators, controlled pollution of experimental plots, big-bag experiments with plankton. Have mostly been used so far for population rather than community studies: not possible to find an example where univariate, graphical/distributional and multivariate techniques have all been applied.

Example: Effect of sediment particle diameter on a harpacticoid copepod community (Hockin, 1982).

LOCATION: Sandy estuarine beach, Ythan estuary, Scotland.

SAMPLING: 2 replicates of 4 grades of glass beads deployed in plastic trays in randomised block design at two tide levels. Left in field for 14 wks, with core sample taken every 5 days.



UNIVARIATE INDICES:




Fig. 12.4 Number of species at upper (solid circles) and lower (open circles) sites




Fig. 12.5 The index of diversity (based on the log-series distribution) for upper (solid circles) and lower (open circles) sites

ANOVA on both the number of species and the species diversity revealed no significant differences with respect to the treatment (sediment particle size).

Table 12.3

Particle diameter of artificial monometric sediments in which the maximum population densities of the numerically dominant harpacticoid copepod species were found.

COPEPOD SPECIES	PARTICLE DIAMETER (MM)
<i>Arenosetella germanica</i>	0.267
<i>Arenosetella tenuissima</i>	0.367
<i>Arenopontia subterranea</i>	0.147
<i>Evansula incerta</i>	0.367
<i>Stenocaris pygmea</i>	0.267
<i>Heterolaophonte minuta</i>	0.485
<i>Heterolaophonte littoralis</i>	0.485
<i>Esola typhlops</i>	0.367
<i>Paronychocamptus curticaudatus</i>	0.485
<i>Huntemannia jadensis</i>	0.147
<i>Nannopus palustris</i>	0.147

Although no MULTIVARIATE ANALYSES were done, different species reached maximum abundance in different sediment grades. This suggests that a multivariate analysis may well have provided discrimination between treatments.

#### LABORATORY EXPERIMENTS

More or less natural communities of some components of the biota can be maintained in laboratory mesocosms or microcosms (also in outdoor mesocosms), and subjected to a variety of manipulations.

Example: Effects of organic enrichment on meiofaunal community structure (Gee et al., 1985).

LOCATION: Sediment from Oslofjord; mesocosm at Solbergstrand, Norway.

SAMPLING: Undisturbed 0.25 m<sup>2</sup> box cores of sediment transferred to mesocosm basin. 4 replicate boxes dosed with high (200 g C m<sup>-2</sup>) and low (50 g C m<sup>-2</sup>) levels of powdered algae (*Ascophyllum*), with 4 undosed controls, in randomised block design. Meiofauna sampled 56 days after dosing: 5 cores from each box combined to give one sample.

UNIVARIATE INDICES:

Nematodes: No significant differences in species richness or diversity between treatments, but evenness significantly higher in enriched boxes than controls.

Copepods: Significant differences in species richness and evenness between treatments, but not in diversity.

Table 12.4

Univariate measures for all replicates at end of experiment, with F-ratio and significance levels from one-way ANOVA.

Treatment	Sample number	Species richness	Shannon-Wiener index	Species evenness
<b>Nematodes</b>				
Control	1	3.023	2.245	0.750
	2	3.739	2.394	0.774
	3	3.357	2.470	0.824
	4	4.589	2.764	0.829
	Total	6.342	2.738	0.747
Low dose	1	4.386	2.856	0.877
	2	2.652	2.474	0.840
	3	4.669	2.885	0.875
	4	2.327	2.268	0.860
	Total	6.153	2.877	0.791
High dose	1	2.856	2.168	0.782
	2	2.824	2.388	0.843
	3	4.302	2.395	0.829
	4	4.088	2.466	0.853
	Total	5.508	2.677	0.759
F-ratio		0.043	1.387	5.131
Significance		ns	ns	P<0.05
<b>Copepods</b>				
Control	1	2.525	1.927	0.927
	2	1.924	1.560	0.969
	3	2.502	1.768	0.908
	4	2.471	1.936	0.931
	Total	2.531	2.102	0.877
Low dose	1	1.804	1.597	0.643
	2	1.661	1.275	0.532
	3	1.655	1.160	0.484
	4	1.786	1.535	0.640
	Total	1.907	1.581	0.584
High dose	1	1.747	1.594	0.767
	2	0.973	0.997	0.620
	3	1.034	0.297	0.165
	4	1.179	1.696	0.872
	Total	1.666	1.683	0.702
F-ratio		17.715	2.654	4.559
Significance		P<0.001	ns	P<0.05

GRAPHICAL/DISTRIBUTIONAL PLOTS:




Fig. 12.6 k-dominance curves for A nematodes, b total copepods and C copepods omitting the 'weed' species of *Tisbe* for summed replicates of each treatment. Circles = control, squares = low dose, triangles = high dose

NEMATODES: No obvious treatment effect.

COPEPODS: Control with highest diversity; when *Tisbe* spp. omitted, sequence of increasing elevation of curves (decreasing diversity) from control to high dose.

MULTIVARIATE ANALYSES:




Fig. 12.7 MDS of double square root transformed abundances of nematodes, copepods and total meiofauna (nematodes + copepods). Circles = control, squares = low dose, triangles = high dose

Table 12.5

Values of the R statistic from the ANOSIM test, in pairwise comparisons between treatments, together with significance levels. C = control, L = low dose, H = high dose.

	TREATMENT	STATISTIC VALUE	% SIG LEVEL
Nematodes	(L, C)	0.27	2.86
	(H, C)	0.22	5.71
	(H, L)	0.28	8.57
Copepods	(L, C)	1.00	2.86
	(H, C)	0.97	2.86
	(H, L)	0.59	2.86

NEMATODES: Only differences between low dose and control treatments are significant at the 5% level.

COPEPODS: Differences between all treatments significant at the 5% level.

Note higher values of the R statistic for copepods in all cases.

CONCLUSIONS: Univariate and graphical/distributional techniques show lowered diversity with increasing dose for copepods, but no effect on nematodes. Multivariate techniques clearly discriminate between treatments for copepods, and still have some discriminating power for nematodes. Changes in nematode community may not have been detectable because of great variability in abundance of nematodes in the high dose boxes.



### LECTURE 13

#### DATA REQUIREMENTS FOR BIOLOGICAL EFFECTS STUDIES: WHICH COMPONENTS AND ATTRIBUTES OF THE BIOTA TO EXAMINE

- COMPONENTS: Pelagos
- plankton
  - fish
- Benthos
- soft-bottom
    - macrobenthos
    - meiobenthos
    - (microbenthos)
  - hard-bottom
    - epifauna
    - motile fauna
      - macrofauna
      - meiofauna
- ATTRIBUTES: Abundance
- species
  - higher taxa
- Biomass
- species
  - higher taxa
- (Production)

#### PLANKTON

##### ADVANTAGES:

- Integrate ecological conditions over areas; useful in monitoring more global changes.
- Taxonomy moderately easy.

##### DISADVANTAGES:

- Not useful for monitoring local effects, due to mobility.

Example: Continuous Plankton Recorder Survey of NE Atlantic.




Fig. 13.1 First principal components for zooplankton and phytoplankton (left) in each of the 12 areas shown in the chart (right). Graphs scaled to zero mean and unit variance

## FISH

### ADVANTAGES:

- Again more useful for general rather than local effects, but demersal spp. may have site-fidelity
- Taxonomy easy (at least in Europe)
- Of immediate commercial/public interest

### DISADVANTAGES:

- Strictly quantitative sampling difficult
- Uncertainty about site-fidelity

Example: Effects of mining activity on coral-reef fish communities in the Maldives.




Fig. 13.2 MDS ordination of fish species abundance data from mined (M) and un-mined (U) reef-tops

### MACROBENTHOS

#### ADVANTAGES:

- Non-mobile, therefore useful for local effects
- Taxonomy relatively easy
- Quantitative sampling easy
- Extensive research literature on community effects

#### DISADVANTAGES:

- Sampling requires relatively large ships
- Sample-processing at sea labour-intensive

- Response time relatively slow (long generation time)
- Unsuitable for causality experiments (slow response time, planktonic larvae).

Example: Amoco Cadiz oil-spill in the Bay of Morlaix.




Fig. 13.3 MDS for macrobenthos at station "Pierre Noire". Sampling months are A:4/77, B:8/77, C:9/77, D:12/77, E:2/78, F:4/78, G:8/78, H:11/78, I:2/79, J:5/79, K:7/79, L:10/79, M:2/80, N:4/80, O:8/80, P:10/80, Q:1/81, R:4/81, S:8/81, T:11/81, U:2/82. Oil-spill was during 3/78, i.e. between E and F

### MEIOBENTHOS

#### ADVANTAGES:

- Useful for local effects studies
- Quantitative sampling easy from small ships
- Samples need not be processed on ship
- Potentially fast response (short generation time)
- Good for causality experiments (direct benthic development, fast response)

DISADVANTAGES:

- Taxonomy considered difficulty
- Community responses not well known or documented

Example: Effects of soldier crab disturbance on nematode assemblages at Eaglehawk Neck, Tasmania.




Fig. 13.4 k-dominance curves for disturbed (D) and undisturbed (U) samples in 4 separate sampling blocks

The macrobenthos & meiobenthos may RESPOND DIFFERENTLY to different kinds of perturbation (e.g. physical disturbance, "pollution") so that a comparative study of both may be indicative of the cause.

Example: Hamilton Harbour, Bermuda.




Fig. 13.5 k-dominance curves for macrobenthos (left) and meiobenthic nematodes (right) at six stations in Hamilton Harbour, Bermuda. Elevated macrofauna curves at stations 3 and 4 suggest that physical disturbance is the cause, since the corresponding meiofauna curves at these sites are not similarly affected

### HARD-BOTTOM EPIFAUNA

#### ADVANTAGES:

- Immobile; good for local effects
- Two dimensional nature permits non-destructive (visual) sampling for determination of temporal changes

#### DISADVANTAGES:

- Remote sampling difficult
- Enumeration of colonial organisms difficult
- Biomass measurements difficult

Example: Effects of the 1982-3 El Niño on Indonesian reef corals.




Fig. 13.6 MDS for coral species percentage cover data for South Pari Island. 1=1981, 3=1983 etc

#### HARD-BOTTOM MOTILE FAUNA

##### DISADVANTAGES:

- Remote sampling difficult
- Quantification difficult
- Responses to perturbation not known
- Suitable habitat (e.g. algae) not always available

Example: Macrofauna and meiofauna of replicated intertidal seaweed samples from the Isles of Scilly.




Fig. 13.7 MDS macrobenthos (left) and meiobenthos (right) from different species of seaweeds: Ch=Chondrus, Lo=Lomentaria, La=Laurencia, Cl=Cladophora, Po=Polysiphonia. Note similarity between the two configurations

#### ABUNDANCE, BIOMASS OR BOTH?

Abundances are easier to measure, but biomass may be a better reflection of the ecological importance of a species within a community. In practice, multivariate analyses of abundance and biomass data give remarkably similar results, despite the fact that the species mainly responsible for discriminating between stations are different.



Example: Frierfjord macrofauna




Fig. 13.8 MDS ordinations for macrofauna abundance and biomass. Note the close similarity

Perturbations of various kinds may affect the distribution of numbers of individuals among species differently from the distribution of biomass among species. This is the basis of the 'ABC' (Abundance Biomass Comparison) method for the assessment of disturbance, which was dealt with in Lecture 8.

SPECIES OR HIGHER TAXA

In a wide variety of pollution-impact studies, it has been found for both graphical-distributional and multivariate analyses that there is surprisingly little loss of information when the species data are aggregated into higher taxa, e.g. genera, families or even phyla. Initial collection of data at the level of higher taxa would result in a considerable saving of time (and cost) in the analysis of samples. This was dealt with in more detail in Lecture 10.

## RECOMMENDATIONS

It is difficult to give firm recommendations as to which components or attributes of the biota should be studied, since this depends on the problem in hand and the expertise and funds available. In general, however, the wider the variety of components and attributes studied, the easier the results will be to interpret. A broad approach at the level of higher taxa is often preferable to a painstakingly detailed analysis of species abundances. If only one component of the fauna is to be studied, then consideration should be given to working up a larger number of stations/replicates at the level of higher taxa in preference to a small number of stations at the species level. Of course, a large number of stations at the species level is always the ideal!

## LECTURE 14

### RELATIVE SENSITIVITIES AND MERITS OF UNIVARIATE, GRAPHICAL/DISTRIBUTIONAL AND MULTIVARIATE TECHNIQUES

Two communities with a completely different taxonomic composition may have identical univariate or graphical/distributional structure, and conversely those comprising the same species may have very different univariate or graphical/distributional structure. Do species dependent and species independent attributes of community structure behave the same or differently in response to environmental changes, and which are the most sensitive? These questions will be addressed by reference to a number of case studies in which a variety of methods of data analysis has been employed.

Example 1: Macrobenthos from Frierfjord/Langesundfjord, Norway (IOC/GEEP Oslo Workshop).

MAP OF SITES: See Fig. 1.1.

UNIVARIATE INDICES:




Fig. 14.1 Means (and 95% CIs) for diversity  $H'$

Site A has higher species diversity ( $H'$ ) and site C the lowest: others not significantly different.

GRAPHICAL/DISTRIBUTIONAL PLOTS:




Fig. 14.2 ABC plots based on totals of 4 replicates. Squares = abundance, crosses = biomass

These indicate C, D and E most stressed, B moderately stressed, A and G unstressed. No tests have been done to determine significance of differences.

MULTIVARIATE ANALYSES:




Fig. 14.3 MDS of 4 replicates at each of sites A-E,G (Bray-Curtis similarities on  $\sqrt{}$ -transformed counts)

Stations B,C and D cluster together (ANOSIM separates B from A and C), E and G together (separated with ANOSIM), A on its own. Clusters correlate with water depth rather than measured levels of anthropogenic variables (see Fig. 11.5).

CONCLUSIONS: Multivariate analysis the most sensitive for discriminating stations (only B and C not significantly different). Univariate and graphical distributions conflict with this. For example, E & G have different ABC plots but cluster together; diversity at E is not significantly different from D, but they are the furthest apart on the MDS plots. However, B, C and D all have low diversity and ABC indicates disturbance. Most likely explanation is that these deep-water stations are affected by seasonal anoxia, rather than anthropogenic pollution.

Example 2: Macrobenthos from Hamilton Harbour, Bermuda (IOC/GEEP Bermuda Workshop).

MAP OF SITES:




Fig. 14.4 Map of Hamilton Harbour showing locations of 6 sampling stations

UNIVARIATE INDICES: See Fig. 8.1. H5 with highest diversity, H3 and H4 with lowest diversity (significantly below neutral model prediction, see Table on page 128).

GRAPHICAL/DISTRIBUTIONAL PLOTS: ABC curves show H2, H6 and H7 undisturbed, H5 moderately disturbed, H3 and H4 moderately/grossly disturbed (Fig. 14.5).

MULTIVARIATE ANALYSES: On MDS (Fig. 14.6) stations ordered (left to right) 5,4,3,2,7,6. ANOSIM gives all sites significantly different from each other. Superimposing values of environmental variables shows close correlation with metals and TBT, not with water depth, sediment type or hydrocarbons.

CONCLUSIONS: MDS most sensitive in discriminating sites, and relates to pollution levels. Diversity not ordered in the same way. Stations with highest pollution levels not the most 'stressed'.




Fig. 14.5 ABC curves for Hamilton Harbour macrobenthos (sum of 4 replicates at each station); A = abundance, B = biomass




Fig. 14.6 A) 2-D MDS configuration for macrofauna standardised root-transformed abundance. B-F) same configuration with symbols representing values of environmental variables superimposed: B) grain size, C) water depth, D) sediment Pb concentration, E) TBT in water, F) sediment PAH



Example 3: Reef corals at South Tikus Island, Indonesia, before and after 1982-3 El Niño.

MAP OF SITES: Not available. Ten sets of 3 x 10 m transects across reef-flat in each year.

UNIVARIATE INDICES: See Fig. 8.2. Immediate post El Niño decline in number of species and  $H'$ , slight recovery in 1984 but no significant change after this. No significant changes in  $J'$ .

GRAPHICAL DISTRIBUTIONAL PLOTS: From 1984 onwards, k-dominance curves lie entirely above that of 1981, indicating no apparent recovery. With ANOSIM, few significant differences between years detectable after 1984.




Fig. 14.7 k-dominance curves for totals of all ten replicates in each year. 1=1981, 2=1982 etc

MULTIVARIATE ANALYSES:




Fig. 14.8 MDS for coral species percentage cover data for South Pari Island. 1=1981, 3=1983 etc

El Niño location shift between 1981 and 1983, with gradual recovery towards the 1981 condition until 1985, then a slight move away again in 1987 and 1988. ANOSIM shows all pairs of years to be significantly different.

CONCLUSIONS: All methods demonstrate the dramatic post El Niño decline in species, though the multivariate techniques were seen to be more sensitive in monitoring the recovery phase in later years.

Example 4: Fish communities from mined and non-mined reef tops in the Maldives.

MAP OF SITES: Not available.

UNIVARIATE INDICES: ANOVA shows no significant effect of mining on  $H'$  or  $J'$ .

GRAPHICAL/DISTRIBUTIONAL PLOTS. k-dominance curves for individual replicates given in Fig. 14.9. ANOSIM shows no significant difference between mined and non-mined sites.




Fig. 14.9 Replicate k-dominance curves for fish communities from mined (top) and non-mined (bottom) reef-tops

MULTIVARIATE ANALYSES:




Fig. 14.10 MDS of fish species abundance data from mined (M) and un-mined (U) reef-tops

Clear separation of mined and non-mined sites, which ANOSIM shows to be significant (though test is unnecessary in such a clear-cut case).

CONCLUSIONS: Clear difference in community composition due to mining activity revealed by multivariate methods, but not detected at all by univariate or graphical/distributional techniques.



UNIVARIATE INDICES: Note that meiofauna and macrofauna show different trends, and for all indices many pairs of weeds are not significantly different from each other.




Fig. 14.12 Species richness (left), Shannon diversity (middle) and evenness (right) for meiofauna (top) and macrofauna (bottom), with 95% confidence intervals. Ch = Chondrus, La = Laurencia, Lo = Lomentaria, Cl = Cladophora, Po = Polysiphonia

GRAPHICAL/DISTRIBUTIONAL PLOTS: k-dominance curves for meiofauna show only *Polysiphonia* with a distinctly lower curve than the other species. For macrofauna, curves not clearly distinguishable from each other.




Fig. 14.13 k-dominance curves for meiofauna (left) and macrofauna (right).  
1 = Chondrus, 2 = Laurencia, 3 = Lomentaria, 4 = Cladophora, 5 = Polysiphonia

MULTIVARIATE ANALYSES: See Fig. 13.7. Two-way ANOSIM (weed species/sites) shows all weed species significantly different for both meiofauna and macrofauna. Note similarity of macrofauna and meiofauna configurations.

CONCLUSIONS: Multivariate methods more sensitive than univariate or graphical/distributional methods for discriminating between weed species. Univariate and graphical/distributional methods give different results for macrobenthos and meiobenthos, whereas for the multivariate methods the results are similar for both.

Example 6: Meiobenthos (nematodes and copepods) from the Tamar estuary, S.W. England (Austen & Warwick, 1989).

MAP OF SITES:




Fig. 14.14 Map of Tamar estuary showing locations of 10 intertidal mud-flat sites

UNIVARIATE INDICES: Not determined.



GRAPHICAL/DISTRIBUTIONAL PLOTS: k-dominance curves for nematodes and copepods do not show similar sequence. For nematodes, sequence does not correspond to the salinity gradient, but for the copepods the agreement with salinity is closer.




Fig. 14.15 k-dominance curves for amalgamated data from 6 replicate cores for nematodes (top) and copepods (bottom)

MULTIVARIATE ANALYSES: Sequence of sites ordered along the salinity gradient for both nematodes and copepods. ANOSIM shows copepod assemblages significantly different at all pairs of sites, nematodes at all pairs except 6/7 and 8/9.




Fig. 14.16 MDS for nematodes (left) and copepods (right) for six replicate cores at each of 10 stations. Note that, allowing for the difference in orientation, the configurations are almost identical

CONCLUSIONS: Multivariate techniques more sensitive in discriminating sites (many sites indistinguishable on basis of k-dominance curves). Multivariate methods give similar patterns for nematodes and copepods; graphical/distributional methods give different patterns for the two taxa. For nematodes, factors other than salinity are more important in determining diversity profiles, but for copepods salinity correlates well with diversity.

Example 7: Meiofauna from Tasmanian sandflat, influenced by burrowing and feeding of soldier crabs.

MAP OF SITES:

UNIVARIATE INDICES:

GRAPHICAL/DISTRIBUTIONAL PLOTS:

See Lecture 12

MULTIVARIATE ANALYSES:




Fig. 14.17 MDS plots for nematode, copepod and 'meiofauna' (nematodes + copepods) abundance. Open symbols = disturbed samples, closed = undisturbed (different shapes denote the flour blocks)

**CONCLUSIONS:** For nematodes, univariate, graphical/distributional and multivariate methods all distinguish disturbed from undisturbed sites. For copepods only the multivariate methods do. Univariate and graphical/distributional methods indicate different responses for nematodes and copepods; multivariate methods indicate a similar response.

## **GENERAL CONCLUSIONS**

Three general conclusions emerge from these examples:

1. Similarity between sites based on their univariate or graphical/distributional properties is usually different from their clustering in multivariate analyses.
2. SPECIES DEPENDENT (multivariate) methods are much more sensitive than SPECIES INDEPENDENT (univariate and graphical/distributional) methods in discriminating between sites.
3. In examples where more than one component of the fauna has been studied, univariate and graphical/distributional methods may give different results for different components, whereas multivariate methods tend to give the same results.

The sensitive multivariate methods are only capable of detecting differences in community composition between sites, although these differences can be correlated with measured levels of stressors such as pollutants. Only the species independent methods of data analysis can be used to determine deleterious (stress) responses. There is a need to develop techniques for determining stress which utilise the full multivariate information contained in a species/sites matrix.

## **RECOMMENDATIONS**

At present, it is important to apply a wide variety of classes of data analysis, as each will give different information and this will aid interpretation. Sensitive multivariate methods will give an 'early warning' that community changes are occurring, but indications that these changes are deleterious are required by environmental managers, and the less sensitive species independent methods must be used.

## FURTHER READING

For general texts on multivariate methods, the two books by Everitt (1978 and 1980) are useful introductions, and Chatfield and Collins (1980) can be recommended (though requires some knowledge of matrix algebra and statistical inference). A more detailed, but still approachable, exposition of MDS is the monograph by Kruskal and Wish (1978). (None of these texts is written from an ecological viewpoint).

Papers which reflect the approach taken in these lectures include Field *et al.* (1982), Warwick (1986), papers from the GEEP Oslo Workshop Proceedings (Mar.Ecol.Prog.Ser.Vol.46), e.g. Gray *et al.* (1988), Warwick *et al.* (1988), Clarke and Green (1988), and from the GEEP Bermuda Workshop Proceedings (to appear in J.Exp.Mar.Biol.Ecol. in July 1990), viz. Clarke (1990) and Warwick *et al.* (1990).

## LITERATURE CITED

- Austen, M.C. and R.M. Warwick (1989), Comparison of univariate and multivariate aspects of estuarine meiobenthic community structure. Est.Cstl.Shelf.Sci., 29:23-42.
- Bayne, B.L., K.R. Clarke, and J.S. Gray (eds.) (1988), Biological effects of pollutants: results of a practical workshop. Mar.Ecol.Prog.Ser., 46 p.
- Box, G.E.P. and D.R. Cox (1964), An analysis of transformations. J.R.Statist.Soc.Ser.B, 26:211-243.
- Bray, J.R. and J.T. Curtis (1957), An ordination of the upland forest communities of Southern Wisconsin. Ecol.Monogr., 27:325-349.
- Buchanana, J.B. and R.M. Warwick (1974), An estimate of benthic macrofauna production in the offshore mud of the Northumberland coast. J.Mar.Biol.Assoc.U.K., 54:197-222.
- Chatfield, C. and A.J. Collins (1980), Introduction to multivariate analysis. London, Chapman and Hall, 1st ed., 246 p.
- Clarke, K.R. (1990), Comparisons of dominance curves. J.Exp.Mar.Biol.Ecol., 138:143-157.
- Clarke, K.R. and R.H. Green (1988), Statistical design and analysis for a 'biological effects' study. Mar.Ecol.Prog.Ser., 46:213-226.
- Collins, N.R. and R. Williams (1982), Zooplankton communities in the Bristol Channel and Severn Estuary. Mar.Ecol.Prog.Ser., 9:1-11.
- Cormack, R.M. (1971), A review of classification. J.R.Statist.Soc.Ser.A, 134:321-367.
- Dawson-Shepherd, A., R.M. Warwick, K.R. Clarke and B.E. Brown (1992), An analysis of fish community responses to coral mining in the Maldives. Env.Biol.Fishes, 33:367-380.

- Everitt, B. (1978), Graphical techniques for multivariate data. London, Heinemann, 117 p.
- Everitt, B. (1980), Cluster analysis, 2nd edn. London, Heinemann, 2nd ed., 136 p.
- Field, J.G., K.R. Clarke and R.M. Warwick (1982), A practical strategy for analysing multispecies distribution patterns. Mar.Ecol.Prog.Ser., 8:37-52.
- Gee, J.M., R.M. Warwick, M. Schaanning, J.A. Berge, and W.G. Ambrose Jr. (1985), Effects of organic enrichment on meiofaunal abundance and community structure in sublittoral soft sediments. J.Exp.Mar.Biol.Ecol., 91:247-262.
- Gower, J.C. (1966), Some distance properties of latent root and vector methods used in multivariate analysis. Biometrika, 53:325-328.
- Gower, J.C. and G.J.S. Ross (1969), Minimum spanning trees and single linkage cluster analysis. Appl.Statist., 18:54-64.
- Gray, J.S. and T.H. Pearson (1982), Objective selection of sensitive species indicative of pollution-induced change in benthic communities. I. Comparative methodology. Mar.Ecol.Prog.Ser., 9:111-119.
- Gray, J.S., M. Aschan, M.R. Carr, K.R. Clarke, R.H. Green, T.H. Pearson, R. Rosenberg and R.M. Warwick (1988), Analysis of community attributes of the benthic macrofauna of Frierfjord/Langesundfjord and in a mesocosm experiment. Mar.Ecol.Prog.Ser., 46:151-165.
- Gray, J.S., K.R. Clarke, R.M. Warwick and G. Hobbs (1990), Detection of initial effects of pollution on marine benthos: an example from the Ekofisk and Eldfisk oilfields, North Sea. Mar.Ecol.Prog.Ser., 66:285-299.
- Heip, C., P.M.J. Herman and K. Soetaert (1988), Data processing, evaluation, and analysis. pp.197-231 in R.P. Higgins and H. Thiel (eds.), Introduction to the study of meiofauna. Smithsonian Institution, Washington DC.
- Hill, M.O. and H.G. Gauch (1980), Detrended correspondence analysis, an improved ordination technique. Vegetatio, 42:47-48.
- Hockin, D.C. (1982), The effects of sediment particle diameter upon the meiobenthic copepod community of an intertidal beach: a field and laboratory experiment. J.Anim.Ecol., 51:555-572.
- Hope, A.C.A. (1968), A simplified Monte Carlo significance test procedure. J.R.Statist.Soc.Ser.B, 30:582-598.
- Kenkel, N.C. and L. Orloci (1986), Applying metric and nonmetric multidimensional scaling to some ecological studies: some new results. Ecology, 67:919-928.

- Kruskal, J.B. and M. Wish (1978), Multidimensional scaling. Beverley Hills, California, Sage Publications, 93 p.
- Lambshead, P.J.D., H.M. Platt and K.M. Shaw (1983), The detection of differences among assemblages of marine benthic species based on an assessment of dominance and diversity. J.Nat.Hist., 17:859-874.
- Lance, G.N. and W.T. Williams (1967), A general theory of classificatory sorting strategies: 1 Hierarchical Systems. Comp.J., 9:373-380.
- McConnaughey, B.H. (1964), The determination and analysis of plankton communities. Marine Research in Indonesia (Spec. No.): 1-25.
- Mantel, N. (1967), The detection of disease clustering and a generalized regression approach. Cancer Res., 27:209-220.
- Mardia, K.V., J.T. Kent and J.M. Bibby (1979), Multivariate analysis. London, Academic Press, 521 p.
- Pearson, T.H. (1975), The benthic ecology of Loch Linnhe and Loch Eil, a sea-loch system on the west coast of Scotland. IV. Changes in the benthic fauna attributable to organic enrichment. J.Exp.Mar.Biol.Ecol., 20:1-41.
- Pearson, T.H., J.S. Gray and P.J. Johannessen (1983), Objective selection of sensitive species indicative of pollution-induced change in benthic communities. 2. Data analyses. Mar.Ecol.Prog.Ser., 12:237-255.
- Sanders, H.L. (1968), Marine benthic diversity: a comparative study. Am.Nat., 102:243-282.
- Seber, G.A.F. (1984), Multivariate observations. New York, Wiley, 686 p.
- Sneath, P.H.A. and R.R. Sokal (1973), Numerical taxonomy. W.H. Freeman, San Francisco.
- Warwick, R.M. (1986), A new method for detecting pollution effects on marine macrobenthic communities. Mar.Biol., 92:557-562.
- Warwick, R.M. (1988), The level of taxonomic discrimination required to detect pollution effects on marine benthic communities. Mar.Pollut.Bull., 19:259-268.
- Warwick, R.M. and K.R. Clarke (1991), A comparison of methods for analysing changes in benthic community structure. J.Mar.Biol.Assoc.U.K., 71:225-244.
- Warwick, R.M., T.H. Pearson and Ruswahyuni (1987), Detection of pollution effects on marine macrobenthos: further evaluation of the species abundance/biomass method. Mar.Biol., 95:193-200.
- Warwick, R.M., M.R. Carr, K.R. Clarke, J.M. Gee and R.H. Green (1988), A mesocosm experiment on the effects of hydrocarbon and copper pollution on a sublittoral soft-sediment meiobenthic community. Mar.Ecol.Prog.Ser., 46:181-191.

Warwick, R.M., K.R. Clarke and Suharsono (1990), A statistical analysis of coral community responses to the 1982-1983 El Niño in the Thousand Islands, Indonesia. *Coral Reefs*. 8:171-179.

Warwick, R.M., K.R. Clarke and J.M. Gee (1990), The effect of disturbance by soldier crabs, Mictyris platycheles H. Milne Edwards, on meiobenthic community structure. J.Exp.Mar.Biol.Ecol., 135:19-33.

Warwick, R.M., H.M. Platt, K.R. Clarke, J. Agard and J. Gobin (1990), Analysis of macrobenthic and meiobenthic community structure in relation to pollution and disturbance in Hamilton Harbour, Bermuda. J.Exp.Mar.Biol.Ecol., (in press).

## **ACKNOWLEDGEMENTS**

We are grateful to Martin Carr, Jane Addy, John Hall, Ralph Bruno, Martin Budge, Mike Ainsworth, Robert Pritchard, Roger Carter, Mike Gee and Mel Austen who in various ways contributed to the programs and analyses featured in this series of lectures.

K.R. Clarke  
R.M. Warwick



## PUBLICATIONS OF THE MAP TECHNICAL REPORTS SERIES

1. UNEP/IOC/WMO: Baseline studies and monitoring of oil and petroleum hydrocarbons in marine waters (MED POL I). MAP Technical Reports Series No. 1. UNEP, Athens, 1986 (96 pages) (parts in English, French or Spanish only).
2. UNEP/FAO: Baseline studies and monitoring of metals, particularly mercury and cadmium, in marine organisms (MED POL II). MAP Technical Reports Series No. 2. UNEP, Athens, 1986 (220 pages) (parts in English, French or Spanish only).
3. UNEP/FAO: Baseline studies and monitoring of DDT, PCBs and other chlorinated hydrocarbons in marine organisms (MED POL III). MAP Technical Reports Series No. 3. UNEP, Athens, 1986 (128 pages) (parts in English, French or Spanish only).
4. UNEP/FAO: Research on the effects of pollutants on marine organisms and their populations (MED POL IV). MAP Technical Reports Series No. 4. UNEP, Athens, 1986 (118 pages) (parts in English, French or Spanish only).
5. UNEP/FAO: Research on the effects of pollutants on marine communities and ecosystems (MED POL V). MAP Technical Reports Series No. 5. UNEP, Athens, 1986 (146 pages) (parts in English or French only).
6. UNEP/IOC: Problems of coastal transport of pollutants (MED POL VI). MAP Technical Reports Series No. 6. UNEP, Athens, 1986 (100 pages) (English only).
7. UNEP/WHO: Coastal water quality control (MED POL VII). MAP Technical Reports Series No. 7. UNEP, Athens, 1986 (426 pages) (parts in English or French only).
8. UNEP/IAEA/IOC: Biogeochemical studies of selected pollutants in the open waters of the Mediterranean (MED POL VIII). MAP Technical Reports Series No. 8. UNEP, Athens, 1986 (42 pages) (parts in English or French only).
8. Add. UNEP: Biogeochemical studies of selected pollutants in the open waters of the Mediterranean (MED POL VIII). Addendum, Greek Oceanographic Cruise 1980. MAP Technical Reports Series No. 8, Addendum. UNEP, Athens, 1986 (66 pages) (English only).
9. UNEP: Co-ordinated Mediterranean pollution monitoring and research programme (MED POL - PHASE I). Final report, 1975-1980. MAP Technical Reports Series No. 9. UNEP, Athens, 1986 (276 pages) (English only).
10. UNEP: Research on the toxicity, persistence, bioaccumulation, carcinogenicity and mutagenicity of selected substances (Activity G). Final reports on projects dealing with toxicity (1983-85). MAP Technical Reports Series No. 10. UNEP, Athens, 1987 (118 pages) (English only).
11. UNEP: Rehabilitation and reconstruction of Mediterranean historic settlements. Documents produced in the first stage of the Priority Action (1984-1985). MAP Technical Reports Series No. 11. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1986 (158 pages) (parts in English or French only).
12. UNEP: Water resources development of small Mediterranean islands and isolated coastal areas. Documents produced in the first stage of the Priority Action (1984-1985). MAP Technical Reports Series No. 12. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (162 pages) (parts in English or French only).

13. UNEP: Specific topics related to water resources development of large Mediterranean islands. Documents produced in the second phase of the Priority Action (1985-1986). MAP Technical Reports Series No. 13. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (162 pages) (parts in English or French only).
14. UNEP: Experience of Mediterranean historic towns in the integrated process of rehabilitation of urban and architectural heritage. Documents produced in the second phase of the Priority Action (1986). MAP Technical Reports Series No. 14. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (500 pages) (parts in English or French only).
15. UNEP: Environmental aspects of aquaculture development in the Mediterranean region. Documents produced in the period 1985-1987. MAP Technical Reports Series No. 15. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (101 pages) (English only).
16. UNEP: Promotion of soil protection as an essential component of environmental protection in Mediterranean coastal zones. Selected documents (1985-1987). MAP Technical Reports Series No. 16. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (424 pages) (parts in English or French only).
17. UNEP: Seismic risk reduction in the Mediterranean region. Selected studies and documents (1985-1987). MAP Technical Reports Series No. 17. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (247 pages) (parts in English or French only).
18. UNEP/FAO/WHO: Assessment of the state of pollution of the Mediterranean Sea by mercury and mercury compounds. MAP Technical Reports Series No. 18. UNEP, Athens, 1987 (354 pages) (English and French).
19. UNEP/IOC: Assessment of the state of pollution of the Mediterranean Sea by petroleum hydrocarbons. MAP Technical Reports Series No. 19. UNEP, Athens, 1988 (130 pages) (English and French).
20. UNEP/WHO: Epidemiological studies related to environmental quality criteria for bathing waters, shellfish-growing waters and edible marine organisms (Activity D). Final report on project on relationship between microbial quality of coastal seawater and health effects (1983-86). MAP Technical Reports Series No. 20. UNEP, Athens, 1988 (156 pages) (English only).
21. UNEP/UNESCO/FAO: Eutrophication in the Mediterranean Sea: Receiving capacity and monitoring of long-term effects. MAP Technical Reports Series No. 21. UNEP, Athens, 1988 (200 pages) (parts in English or French only).
22. UNEP/FAO: Study of ecosystem modifications in areas influenced by pollutants (Activity I). MAP Technical Reports Series No. 22. UNEP, Athens, 1988 (146 pages) (parts in English or French only).
23. UNEP: National monitoring programme of Yugoslavia, Report for 1983-1986. MAP Technical Reports Series No. 23. UNEP, Athens, 1988 (223 pages) (English only).
24. UNEP/FAO: Toxicity, persistence and bioaccumulation of selected substances to marine organisms (Activity G). MAP Technical Reports Series No. 24. UNEP, Athens, 1988 (122 pages) (parts in English or French only).
25. UNEP: The Mediterranean Action Plan in a functional perspective: A quest for law and policy. MAP Technical Reports Series No. 25. UNEP, Athens, 1988 (105 pages) (English only).

26. UNEP/IUCN: Directory of marine and coastal protected areas in the Mediterranean Region. Part I - Sites of biological and ecological value. MAP Technical Reports Series No. 26. UNEP, Athens, 1989 (196 pages) (English only).
27. UNEP: Implications of expected climate changes in the Mediterranean Region: An overview. MAP Technical Reports Series No. 27. UNEP, Athens, 1989 (52 pages) (English only).
28. UNEP: State of the Mediterranean marine environment. MAP Technical Reports Series No. 28. UNEP, Athens, 1989 (225 pages) (English only).
29. UNEP: Bibliography on effects of climatic change and related topics. MAP Technical Reports Series No. 29. UNEP, Athens, 1989 (143 pages) (English only).
30. UNEP: Meteorological and climatological data from surface and upper measurements for the assessment of atmospheric transport and deposition of pollutants in the Mediterranean Basin: A review. MAP Technical Reports Series No. 30. UNEP, Athens, 1989 (137 pages) (English only).
31. UNEP/WMO: Airborne pollution of the Mediterranean Sea. Report and proceedings of a WMO/UNEP Workshop. MAP Technical Reports Series No. 31. UNEP, Athens, 1989 (247 pages) (parts in English or French only).
32. UNEP/FAO: Biogeochemical cycles of specific pollutants (Activity K). MAP Technical Reports Series No. 32. UNEP, Athens, 1989 (139 pages) (parts in English or French only).
33. UNEP/FAO/WHO/IAEA: Assessment of organotin compounds as marine pollutants in the Mediterranean. MAP Technical Reports Series No. 33. UNEP, Athens, 1989 (185 pages) (English and French).
34. UNEP/FAO/WHO: Assessment of the state of pollution of the Mediterranean Sea by cadmium and cadmium compounds. MAP Technical Reports Series No. 34. UNEP, Athens, 1989 (175 pages) (English and French).
35. UNEP: Bibliography on marine pollution by organotin compounds. MAP Technical Reports Series No. 35. UNEP, Athens, 1989 (92 pages) (English only).
36. UNEP/IUCN: Directory of marine and coastal protected areas in the Mediterranean region. Part I - Sites of biological and ecological value. MAP Technical Reports Series No. 36. UNEP, Athens, 1990 (198 pages) (French only).
37. UNEP/FAO: Final reports on research projects dealing with eutrophication and plankton blooms (Activity H). MAP Technical Reports Series No. 37. UNEP, Athens, 1990 (74 pages) (parts in English or French only).
38. UNEP: Common measures adopted by the Contracting Parties to the Convention for the Protection of the Mediterranean Sea against pollution. MAP Technical Reports Series No. 38. UNEP, Athens, 1990 (100 pages) (English, French, Spanish and Arabic).
39. UNEP/FAO/WHO/IAEA: Assessment of the state of pollution of the Mediterranean Sea by organohalogen compounds. MAP Technical Reports Series No. 39. UNEP, Athens, 1990 (224 pages) (English and French).
40. UNEP/FAO: Final reports on research projects (Activities H,I and J). MAP Technical Reports Series No. 40. UNEP, Athens, 1990 (125 pages) (English and French).
41. UNEP: Wastewater reuse for irrigation in the Mediterranean region. MAP Technical Reports Series No. 41. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1990 (330 pages) (English and French).

42. UNEP/IUCN: Report on the status of Mediterranean marine turtles. MAP Technical Reports Series No. 42. UNEP, Athens, 1990 (204 pages) (English and French).
43. UNEP/IUCN/GIS Posidonia: Red Book "Gérard Vuignier", marine plants, populations and landscapes threatened in the Mediterranean. MAP Technical Reports Series No. 43. UNEP, Athens, 1990 (250 pages) (French only).
44. UNEP: Bibliography on aquatic pollution by organophosphorus compounds. MAP Technical Reports Series No. 44. UNEP, Athens, 1990 (98 pages) (English only).
45. UNEP/IAEA: Transport of pollutants by sedimentation: Collected papers from the first Mediterranean Workshop (Villefranche-sur-Mer, France, 10-12 December 1987). MAP Technical Reports Series No. 45. UNEP, Athens, 1990 (302 pages) (English only).
46. UNEP/WHO: Epidemiological studies related to environmental quality criteria for bathing waters, shellfish-growing waters and edible marine organisms (Activity D). Final report on project on relationship between microbial quality of coastal seawater and rotavirus-induced gastroenteritis among bathers (1986-88). MAP Technical Reports Series No.46, UNEP, Athens, 1991 (64 pages) (English only).
47. UNEP: Jellyfish blooms in the Mediterranean. Proceedings of the II workshop on jellyfish in the Mediterranean Sea. MAP Technical Reports Series No.47. UNEP, Athens, 1991 (320 pages) (parts in English or French only).
48. UNEP/FAO: Final reports on research projects (Activity G). MAP Technical Reports Series No. 48. UNEP, Athens, 1991 (126 pages) (parts in English or French only).
49. UNEP/WHO: Biogeochemical cycles of specific pollutants. Survival of pathogens. Final reports on research projects (Activity K). MAP Technical Reports Series No. 49. UNEP, Athens, 1991 (71 pages) (parts in English or French only).
50. UNEP: Bibliography on marine litter. MAP Technical Reports Series No. 50. UNEP, Athens, 1991 (62 pages) (English only).
51. UNEP/FAO: Final reports on research projects dealing with mercury, toxicity and analytical techniques. MAP Technical Reports Series No. 51. UNEP, Athens, 1991 (166 pages) (parts in English or French only).
52. UNEP/FAO: Final reports on research projects dealing with bioaccumulation and toxicity of chemical pollutants. MAP Technical Reports Series No. 52. UNEP, Athens, 1991 (86 pages) (parts in English or French only).
53. UNEP/WHO: Epidemiological studies related to environmental quality criteria for bathing waters, shellfish-growing waters and edible marine organisms (Activity D). Final report on epidemiological study on bathers from selected beaches in Malaga, Spain (1988-1989). MAP Technical Reports Series No. 53. UNEP, Athens, 1991 (127 pages) (English only).
54. UNEP/WHO: Development and testing of sampling and analytical techniques for monitoring of marine pollutants (Activity A): Final reports on selected microbiological projects. MAP Technical Reports Series No. 54. UNEP, Athens, 1991 (83 pages) (English only).
55. UNEP/WHO: Biogeochemical cycles of specific pollutants (Activity K): Final report on project on survival of pathogenic organisms in seawater. MAP Technical Reports Series No. 55. UNEP, Athens, 1991 (95 pages) (English only).
56. UNEP/IOC/FAO: Assessment of the state of pollution of the Mediterranean Sea by persistent synthetic materials which may float, sink or remain in suspension. MAP Technical Reports Series No. 56. UNEP, Athens, 1991 (113 pages) (English and French).

57. UNEP/WHO: Research on the toxicity, persistence, bioaccumulation, carcinogenicity and mutagenicity of selected substances (Activity G): Final reports on projects dealing with carcinogenicity and mutagenicity. MAP Technical Reports Series No. 57. UNEP, Athens, 1991 (59 pages) (English only).
58. UNEP/FAO/WHO/IAEA: Assessment of the state of pollution of the Mediterranean Sea by organophosphorus compounds. MAP Technical Reports Series No. 58. UNEP, Athens, 1991 (122 pages) (English and French).
59. UNEP/FAO/IAEA: Proceedings of the FAO/UNEP/IAEA Consultation Meeting on the Accumulation and Transformation of Chemical contaminants by Biotic and Abiotic Processes in the Marine Environment (La Spezia, Italy, 24-28 September 1990), edited by G.P. Gabrielides. MAP Technical Reports Series No. 59. UNEP, Athens, 1991 (392 pages) (English only).
60. UNEP/WHO: Development and testing of sampling and analytical techniques for monitoring of marine pollutants (Activity A): Final reports on selected microbiological projects (1987-1990). MAP Technical Reports Series No. 60. UNEP, Athens, 1991 (76 pages) (parts in English or French only).
61. UNEP: Integrated Planning and Management of the Mediterranean Coastal Zones. Documents produced in the first and second stage of the Priority Action (1985-1986). MAP Technical Reports Series No. 61. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1991 (437 pages) (parts in English or French only).
62. UNEP/IAEA: Assessment of the State of Pollution of the Mediterranean Sea by Radioactive Substances. MAP Technical Reports Series No. 62, UNEP, Athens, 1992 (133 pages) (English and French).
63. UNEP/WHO: Biogeochemical cycles of specific pollutants (Activity K) - Survival of Pathogens - Final reports on Research Projects (1989-1991). MAP Technical Reports Series No. 63, UNEP, Athens, 1992 (86 pages) (French only).
64. UNEP/WMO: Airborne Pollution of the Mediterranean Sea. Report and Proceedings of the Second WMO/UNEP Workshop. MAP Technical Reports Series No. 64, UNEP, Athens, 1992 (246 pages) (English only).
65. UNEP: Directory of Mediterranean Marine Environmental Centres. MAP Technical Reports Series No. 65, UNEP, Athens, 1992 (351 pages) (English and French).
66. UNEP/CRU: Regional Changes in Climate in the Mediterranean Basin Due to Global Greenhouse Gas Warming. MAP Technical Reports Series No. 66, UNEP, Athens, 1992 (172 pages) (English only).
67. UNEP/IOC: Applicability of Remote Sensing for Survey of Water Quality Parameters in the Mediterranean. Final Report of the Research Project. MAP Technical Reports Series No. 67, UNEP, Athens, 1992 (142 pages) (English only).

### **PUBLICATIONS "MAP TECHNICAL REPORTS SERIES"**

1. PNUE/COI/OMM: Etudes de base et surveillance continue du pétrole et des hydrocarbures contenus dans les eaux de la mer (MED POL I). MAP Technical Reports Series No. 1. UNEP, Athens, 1986 (96 pages) (parties en anglais, français ou espagnol seulement).
2. PNUE/FAO: Etudes de base et surveillance continue des métaux, notamment du mercure et du cadmium, dans les organismes marins (MED POL II). MAP Technical Reports Series No. 2. UNEP, Athens, 1986 (220 pages) (parties en anglais, français ou espagnol seulement).
3. PNUE/FAO: Etudes de base et surveillance continue du DDT, des PCB et des autres hydrocarbures chlorés contenus dans les organismes marins (MED POL III). MAP Technical Reports Series No. 3. UNEP, Athens, 1986 (128 pages) (parties en anglais, français ou espagnol seulement).
4. PNUE/FAO: Recherche sur les effets des polluants sur les organismes marins et leurs peuplements (MED POL IV). MAP Technical Reports Series No. 4. UNEP, Athens, 1986 (118 pages) (parties en anglais, français ou espagnol seulement).
5. PNUE/FAO: Recherche sur les effets des polluants sur les communautés et écosystèmes marins (MED POL V). MAP Technical Reports Series No. 5. UNEP, Athens, 1986 (146 pages) (parties en anglais ou français seulement).
6. PNUE/COI: Problèmes du transfert des polluants le long des côtes (MED POL VI). MAP Technical Reports Series No. 6. UNEP, Athens, 1986 (100 pages) (anglais seulement).
7. PNUE/OMS: Contrôle de la qualité des eaux côtières (MED POL VII). MAP Technical Reports Series No. 7. UNEP, Athens, 1986 (426 pages) (parties en anglais ou français seulement).
8. PNUE/AIEA/COI: Etudes biogéochimiques de certains polluants au large de la Méditerranée (MED POL VIII). MAP Technical Reports Series No. 8. UNEP, Athens, 1986 (42 pages) (parties en anglais ou français seulement).
8. Add. PNUE: Etudes biogéochimiques de certains polluants au large de la Méditerranée (MED POL VIII). Addendum, Croisière Océanographique de la Grèce 1980. MAP Technical Reports Series No. 8, Addendum. UNEP, Athens, 1986 (66 pages) (anglais seulement).
9. PNUE: Programme coordonné de surveillance continue et de recherche en matière de pollution dans la Méditerranée (MED POL -PHASE I). Rapport final, 1975-1980. MAP Technical Reports Series No. 9. UNEP, Athens, 1986 (276 pages) (anglais seulement).
10. PNUE: Recherches sur la toxicité, la persistance, la bioaccumulation, la cancérogénicité et la mutagénicité de certaines substances (Activité G). Rapports finaux sur les projets ayant trait à la toxicité (1983-85). MAP Technical Reports Series No. 10. UNEP, Athens, 1987 (118 pages) (anglais seulement).
11. PNUE: Réhabilitation et reconstruction des établissements historiques méditerranéens. Textes rédigés au cours de la première phase de l'action prioritaire (1984-1985). MAP Technical Reports Series No. 11. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1986 (158 pages) (parties en anglais ou français seulement).
12. PNUE: Développement des ressources en eau des petites îles et des zones côtières isolées méditerranéennes. Textes rédigés au cours de la première phase de l'action prioritaire (1984-1985). MAP Technical Reports Series No. 12. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (162 pages) (parties en anglais ou français seulement).

13. PNUE: Thèmes spécifiques concernant le développement des ressources en eau des grandes îles méditerranéennes. Textes rédigés au cours de la deuxième phase de l'action prioritaire (1985-1986). MAP Technical Reports Series No. 13. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (162 pages) (parties en anglais ou français seulement).
14. PNUE: L'expérience des villes historiques de la Méditerranée dans le processus intégré de réhabilitation du patrimoine urbain et architectural. Documents établis lors de la seconde phase de l'Action prioritaire (1986). MAP Technical Reports Series No. 14. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (500 pages) (parties en anglais ou français seulement).
15. PNUE: Aspects environnementaux du développement de l'aquaculture dans la région méditerranéenne. Documents établis pendant la période 1985-1987. MAP Technical Reports Series No. 15. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (101 pages) (anglais seulement).
16. PNUE: Promotion de la protection des sols comme élément essentiel de la protection de l'environnement dans les zones côtières méditerranéennes. Documents sélectionnés (1985-1987). MAP Technical Reports Series No. 16. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (424 pages) (parties en anglais ou français seulement).
17. PNUE: Réduction des risques sismiques dans la région méditerranéenne. Documents et études sélectionnés (1985-1987). MAP Technical Reports Series No. 17. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1987 (247 pages) (parties en anglais ou français seulement).
18. PNUE/FAO/OMS: Evaluation de l'état de la pollution de la mer Méditerranée par le mercure et les composés mercuriels. MAP Technical Reports Series No. 18. UNEP, Athens, 1987 (354 pages) (anglais et français).
19. PNUE/COI: Evaluation de l'état de la pollution de la mer Méditerranée par les hydrocarbures de pétrole. MAP Technical Reports Series No. 19. UNEP, Athens, 1988 (130 pages) (anglais et français).
20. PNUE/OMS: Etudes épidémiologiques relatives aux critères de la qualité de l'environnement pour les eaux servant à la baignade, à la culture de coquillages et à l'élevage d'autres organismes marins comestibles (Activité D). Rapport final sur le projet sur la relation entre la qualité microbienne des eaux marines côtières et les effets sur la santé (1983-86). MAP Technical Reports Series No. 20. UNEP, Athens, 1988 (156 pages) (anglais seulement).
21. PNUE/UNESCO/FAO: Eutrophisation dans la mer Méditerranée: capacité réceptrice et surveillance continue des effets à long terme. MAP Technical Reports Series No. 21. UNEP, Athens, 1988 (200 pages) (parties en anglais ou français seulement).
22. PNUE/FAO: Etude des modifications de l'écosystème dans les zones soumises à l'influence des polluants (Activité I). MAP Technical Reports Series No. 22. UNEP, Athens, 1988 (146 pages) (parties en anglais ou français seulement).
23. PNUE: Programme national de surveillance continue pour la Yougoslavie, Rapport pour 1983-1986. MAP Technical Reports Series No. 23. UNEP, Athens, 1988 (223 pages) (anglais seulement).
24. PNUE/FAO: Toxicité, persistance et bioaccumulation de certaines substances vis-à-vis des organismes marins (Activité G). MAP Technical Reports Series No. 24. UNEP, Athens, 1988 (122 pages) (parties en anglais ou français seulement).

25. PNUE: Le Plan d'action pour la Méditerranée, perspective fonctionnelle; une recherche juridique et politique. MAP Technical Reports Series No. 25. UNEP, Athens, 1988 (105 pages) (anglais seulement).
26. PNUE/UICN: Répertoire des aires marines et côtières protégées de la Méditerranée. Première partie - Sites d'importance biologique et écologique. MAP Technical Reports Series No. 26. UNEP, Athens, 1989 (196 pages) (anglais seulement).
27. PNUE: Implications des modifications climatiques prévues dans la région méditerranéenne: une vue d'ensemble. MAP Technical Reports Series No. 27. UNEP, Athens, 1989 (52 pages) (anglais seulement).
28. PNUE: Etat du milieu marin en Méditerranée. MAP Technical Reports Series No. 28. UNEP, Athens, 1989 (225 pages) (anglais seulement).
29. PNUE: Bibliographie sur les effets des modifications climatiques et sujets connexes. MAP Technical Reports Series No. 29. UNEP, Athens, 1989 (143 pages) (anglais seulement).
30. PNUE: Données météorologiques et climatologiques provenant de mesures effectuées dans l'air en surface et en altitude en vue de l'évaluation du transfert et du dépôt atmosphériques des polluants dans le bassin méditerranéen: un compte rendu. MAP Technical Reports Series No. 30. UNEP, Athens, 1989 (137 pages) (anglais seulement).
31. PNUE/OMM: Pollution par voie atmosphérique de la mer Méditerranée. Rapport et actes des Journées d'étude OMM/PNUE. MAP Technical Reports Series No. 31. UNEP, Athens, 1989 (247 pages) (parties en anglais ou français seulement).
32. PNUE/FAO: Cycles biogéochimiques de polluants spécifiques (Activité K). MAP Technical Reports Series No. 32. UNEP, Athens, 1989 (139 pages) (parties en anglais ou français seulement).
33. PNUE/FAO/OMS/AIEA: Evaluation des composés organostanniques en tant que polluants du milieu marin en Méditerranée. MAP Technical Reports Series No. 33. UNEP, Athens, 1989 (185 pages) (anglais et français).
34. PNUE/FAO/OMS: Evaluation de l'état de la pollution de la mer Méditerranée par le cadmium et les composés de cadmium. MAP Technical Reports Series No. 34. UNEP, Athens, 1989 (175 pages) (anglais et français).
35. PNUE: Bibliographie sur la pollution marine par les composés organostanniques. MAP Technical Reports Series No. 35. UNEP, Athens, 1989 (92 pages) (anglais seulement).
36. PNUE/UICN: Répertoire des aires marines et côtières protégées de la Méditerranée. Première partie - Sites d'importance biologique et écologique. MAP Technical Reports Series No. 36. UNEP, Athens, 1990 (198 pages) (français seulement).
37. PNUE/FAO: Rapports finaux sur les projets de recherche consacrés à l'eutrophisation et aux efflorescences de plancton (Activité H). MAP Technical Reports Series No. 37. UNEP, Athens, 1990 (74 pages) (parties en anglais ou français seulement).
38. PNUE: Mesures communes adoptées par les Parties Contractantes à la Convention pour la protection de la mer Méditerranée contre la pollution. MAP Technical Reports Series No. 38. UNEP, Athens, 1990 (100 pages) (anglais, français, espagnol et arabe).
39. PNUE/FAO/OMS/AIEA: Evaluation de l'état de la pollution par les composés organohalogénés. MAP Technical Reports Series No. 39. UNEP, Athens, 1990 (224 pages) (anglais et français).



40. PNUE/FAO: Rapports finaux sur les projets de recherche (Activités H, I et J). MAP Technical Reports Series No. 40. UNEP, Athens, 1990 (125 pages) (anglais et français).
41. PNUE: Réutilisation agricole des eaux usées dans la région méditerranéenne. MAP Technical Reports Series No. 41. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1990 (330 pages) (anglais et français).
42. PNUE/UICN: Rapport sur le statut des tortues marines de Méditerranée. MAP Technical Reports Series No. 42. UNEP, Athens, 1990 (204 pages) (anglais et français).
43. PNUE/UICN/GIS Posidonie: Livre rouge "Gérard Vuignier" des végétaux, peuplements et paysages marins menacés de Méditerranée. MAP Technical Reports Series No. 43. UNEP, Athens, 1990 (250 pages) (français seulement).
44. PNUE: Bibliographie sur la pollution aquatique par les composés organophosphorés. MAP Technical Reports Series No. 44. UNEP, Athens, 1990 (98 pages) (anglais seulement).
45. PNUE/AIEA: Transfert des polluants par sédimentation: Recueil des communications présentées aux premières journées d'études méditerranéennes (Villefranche-sur-Mer, France, 10-12 décembre 1987). MAP Technical Reports Series No. 45. UNEP, Athens, 1990 (302 pages) (anglais seulement).
46. PNUE/OMS: Etudes épidémiologiques relatives aux critères de la qualité de l'environnement pour les eaux servant à la baignade, à la culture de coquillages et à l'élevage d'autres organismes marins comestibles (Activité D). Rapport final sur le projet sur la relation entre la qualité microbienne des eaux marines côtières et la gastroentérite provoquée par le rotavirus entre les baigneurs (1986-88). MAP Technical Reports Series No.46. UNEP, Athens, 1991 (64 pages) (anglais seulement).
47. PNUE: Les proliférations de méduses en Méditerranée. Actes des 11èmes journées d'étude sur les méduses en mer Méditerranée. MAP Technical Reports Series No.47. UNEP, Athens, 1991 (320 pages) (parties en anglais ou français seulement).
48. PNUE/FAO: Rapports finaux sur les projets de recherche (Activité G). MAP Technical Reports Series No. 48. UNEP, Athens, 1991 (126 pages) (parties en anglais ou français seulement).
49. PNUE/OMS: Cycles biogéochimiques de polluants spécifiques. Survie des Pathogènes. Rapports finaux sur les projets de recherche (activité K). MAP Technical Reports Series No. 49. UNEP, Athens, 1991 (71 pages) (parties en anglais ou français seulement).
50. PNUE: Bibliographie sur les déchets marins. MAP Technical Reports Series No. 50. UNEP, Athens, 1991 (62 pages) (anglais seulement).
51. PNUE/FAO: Rapports finaux sur les projets de recherche traitant du mercure, de la toxicité et des techniques analytiques. MAP Technical Reports Series No. 51. UNEP, Athens, 1991 (166 pages) (parties en anglais ou français seulement).
52. PNUE/FAO: Rapports finaux sur les projets de recherche traitant de la bioaccumulation et de la toxicité des polluants chimiques. MAP Technical Reports Series No. 52. UNEP, Athens, 1991 (86 pages) (parties en anglais ou français seulement).
53. PNUE/OMS: Etudes épidémiologiques relatives aux critères de la qualité de l'environnement pour les eaux servant à la baignade, à la culture de coquillages et à l'élevage d'autres organismes marins comestibles (Activité D). Rapport final sur l'étude épidémiologique menée parmi les baigneurs de certaines plages à Malaga, Espagne (1988-1989). MAP Technical Reports Series No. 53. UNEP, Athens, 1991 (127 pages) (anglais seulement).

54. PNUE/OMS: Mise au point et essai des techniques d'échantillonnage et d'analyse pour la surveillance continue des polluants marins (Activité A): Rapports finaux sur certains projets de nature microbiologique. MAP Technical Reports Series No. 54. UNEP, Athens, 1991 (83 pages) (anglais seulement).
55. PNUE/OMS: Cycles biogéochimiques de polluants spécifiques (Activité K): Rapport final sur le projet sur la survie des microorganismes pathogènes dans l'eau de mer. MAP Technical Reports Series No. 55. UNEP, Athens, 1991 (95 pages) (anglais seulement).
56. PNUE/COI/FAO: Evaluation de l'état de la pollution de la mer Méditerranée par les matières synthétiques persistantes qui peuvent flotter, couler ou rester en suspension. MAP Technical Reports Series No. 56. UNEP, Athens, 1991 (113 pages) (anglais et français).
57. PNUE/OMS: Recherches sur la toxicité, la persistance, la bioaccumulation, la cancérogénicité et la mutagénicité de certaines substances (Activité G). Rapports finaux sur les projets ayant trait à la cancérogénicité et la mutagénicité. MAP Technical Reports Series No. 57. UNEP, Athens, 1991 (59 pages) (anglais seulement).
58. PNUE/FAO/OMS/AIEA: Evaluation de l'état de la pollution de la mer Méditerranée par les composés organophosphorés. MAP Technical Reports Series No. 58. UNEP, Athens, 1991 (122 pages) (anglais et français).
59. PNUE/FAO/AIEA: Actes de la réunion consultative FAO/PNUE/AIEA sur l'accumulation et la transformation des contaminants chimiques par les processus biotiques et abiotiques dans le milieu marin (La Spezia, Italie, 24-28 septembre 1990), publié sous la direction de G.P. Gabrielides. MAP Technical Reports Series No. 59. UNEP, Athens, 1991 (392 pages) (anglais seulement).
60. PNUE/OMS: Mise au point et essai des techniques d'échantillonnage et d'analyse pour la surveillance continue des polluants marins (Activité A): Rapports finaux sur certains projets de nature microbiologique (1987-1990). MAP Technical Reports Series No. 60. UNEP, Athens, 1991 (76 pages) (parties en anglais ou français seulement).
61. PNUE: Planification intégrée et gestion des zones côtières méditerranéennes. Textes rédigés au cours de la première et de la deuxième phase de l'action prioritaire (1985-1986). MAP Technical Reports Series No. 61. UNEP, Priority Actions Programme, Regional Activity Centre, Split, 1991 (437 pages) (parties en anglais ou français seulement).
62. PNUE/AIEA: Evaluation de l'état de la pollution de la mer Méditerranée par les substances radioactives. MAP Technical Reports Series No. 62, UNEP, Athens, 1992 (133 pages) (anglais et français).
63. PNUE/OMS: Cycles biogéochimiques de polluants spécifiques (Activité K) - Survie des pathogènes - Rapports finaux sur les projets de recherche (1989-1991). MAP Technical Reports Series No. 63, UNEP, Athens, 1992 (86 pages) (français seulement).
64. PNUE/OMM: Pollution par voie atmosphérique de la mer Méditerranée. Rapport et actes des deuxièmes journées d'études OMM/PNUE. MAP Technical Reports Series No. 64, UNEP, Athens, 1992 (246 pages) (anglais seulement).
65. PNUE: Répertoire des centres relatifs au milieu marin en Méditerranée. MAP Technical Reports Series No. 65, UNEP, Athens, 1992 (351 pages) (anglais et français).
66. PNUE/CRU: Modifications régionales du climat dans le bassin méditerranéen résultant du réchauffement global dû aux gaz à effet de serre. MAP Technical Reports Series No. 66, UNEP, Athens, 1992 (172 pages) (anglais seulement).

67. PNUE/COI: Applicabilité de la télédétection à l'étude des paramètres de la qualité de l'eau en Méditerranée. Rapport final du projet de recherche. MAP Technical Reports Series No. 67, UNEP, Athens, 1992 (142 pages) (anglais seulement).



Issued and printed by:

Mediterranean Action Plan  
United Nations Environment Programme

Additional copies of this and other publications issued by  
the Mediterranean Action Plan of UNEP can be obtained from:

Coordinating Unit for the Mediterranean Action Plan  
United Nations Environment Programme  
Leoforos Vassileos Konstantinou, 48  
P.O.Box 18019  
11610 Athens  
GREECE



Publié et imprimé par:

Plan d'action pour la Méditerranée  
Programme des Nations Unies pour l'Environnement

Des exemplaires de ce document ainsi que d'autres  
publications du Plan d'action pour la Méditerranée  
du PNUE peuvent être obtenus de:

Unité de coordination du Plan d'action pour la Méditerranée  
Programme des Nations Unies pour l'Environnement  
Leoforos Vassileos Konstantinou, 48  
B.P. 18019  
11610 Athènes  
GRECE